

Submitted 3/20/2012

1.8111 3.5434

03771600



NO INK
NO VARNISH

Drug Facts (continued)

Frequent chest pain

- frequent, wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking warfarin, clopidogrel or cimetidine (blood-thinning medicines)
- taking prescription antifungal or anti-yeast medicines
- taking diazepam (anxiety medicine)
- taking digoxin (heart medicine)
- taking sacrolimus (immune system medicine)
- taking prescription antiretrovirals (medicines for HIV infection)
- taking any other prescription drugs. Sodium bicarbonate may interact with certain prescription drugs.

Stop use and ask a doctor if

- you get diarrhea
- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Other Information

- each capsule contains sodium 300 mg
- read the directions, warnings and accompanying label information before use
- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if the blue band around the capsule is missing or broken. Do not use if foil liner seal imprinted with "Sealed for your protection" is missing, open or broken.
- keep product out of high heat and humidity

Inactive Ingredients

croscarmellose sodium, FD&C blue No. 1, gelatin, magnesium stearate, pharmaceutical ink, polysorbate 80, titanium dioxide

Questions or comments?
Call 1-888-4-ZEG-OTC (1-888-483-4822) between 8:00 AM and 5:00 PM Central Time, Monday through Friday or visit www.ZegeridOTC.com

© Copyright ©, Distributed by MSD Consumer Care, Inc., PO Box 377, Memphis, TN 38151 USA.
A subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA.
All rights reserved. Product of Spain.
37716-01PKG00866
157/6566
each

FREE STARTER PACK

Treats Frequent Heartburn

Zegerid OTC

• Omeprazole 20 mg/Acid Reducer
• Sodium Bicarbonate 1100 mg/
Allows Absorption of this Omeprazole Product

2 CAPSULES
Follow Samples with a 14-Day Course of Treatment

SAMPLE-NOT FOR SALE

Drug Facts

Active Ingredients (in each capsule)

Omeprazole 20 mg
Sodium Bicarbonate 1100 mg

Purpose

Acid reducer

Always absorption of this omeprazole product

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.

Warnings

Alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food
- vomiting with blood
- bloody or black stools

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Tips for Managing Heartburn

- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some fruits and vegetables
- Do not lie flat or bend over soon after eating
- Do not eat, take or use before bedtime
- Raise the head of your bed
- Wear loose-fitting clothing around your stomach

NDC 1152-7265-5

PRINTSIDE GLUE

FOLD

PRINTSIDE GLUE

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic

Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd. Bold Italic (continued) - 8 pt Helvetica Cnd.

Headings - 8 pt Helvetica Cnd. Bold Italic

Sub Heads - 6 pt Helvetica Cnd. Bold

Text - 6 pt Helvetica Cnd, minimum leading 6.5

Square Bullets - 3.5 pt Zapf Dingbats no compression

Border - 1 pt rule, 3/64" gap all around

Barline - 1 pt rule

Hairline - .5 pt rule

Certified by: eb
APO #: 2012-93465
02.23.12 rp

LOT/EXP
Area
No ink
No Varnish


NO INK
NO VARNISH

1.497


Submitted 3/20/2012



A - Top Panel




B - Back Panel




Base Panel


Printable Text Area




C - Insert 1 Top




D - Insert 1 Back




E - Insert 2 Top




F - Insert 2 Back




G - Insert 3 Top




H - Insert 3 Back




I - Insert 4 Top




J - Insert 4 Back




K - Insert 5 Top



L - Insert 5 Back



M - Insert 6 Top



N - Insert 6 Back

Submitted 3/20/2012

1.8111 3.5434

03658905



NO INK
NO VARNISH

Drug Facts (continued)

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking:

- warfarin, clopidogrel or cloisazoli (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- diazepam (anxiety medicine)
- digoxin (heart medicine)
- tacrolimus (immune system medicine)
- prescription antiretrovirals (medicines for HIV infection)

Slip use and ask a doctor if:

- you get diarrhea
- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water at least 1 hour before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- do not chew or crush the capsule
- do not open capsule and sprinkle on food
- do not use for more than 14 days unless directed by your doctor

Repeat 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor.

Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- each capsule contains: sodium 303 mg
- read the directions, warnings and accompanying label information before use
- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if the blue band around the capsule is missing or broken. Do not use if foil inner seal imprinted with "Sealed for your protection" is missing, open or broken.
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients

croscarmellose sodium, FD&C blue No. 1, gelatin, magnesium stearate, pharmaceutical ink, polysorbate 80, titanium dioxide

Questions or comments?
Call 1-888-4-ZEG-OTC (1-888-482-4882) between 8:00 AM and 5:00 PM Central Time, Monday through Friday or visit www.ZegeridOTC.com

© Copyright & distributed by MSD Consumer Care, Inc., PO Box 377, Memphis, TN 38151 USA, a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA. All rights reserved. Product of Spain. 36589-05PKG01272

recyclable carton

Treats Frequent Heartburn

Zegerid[®] OTC

• Omeprazole 20 mg/Acid Reducer
• Sodium Bicarbonate 1100 mg/
Allows Absorption of this Omeprazole Product

14 CAPSULES
One 14-Day Course of Treatment

Drug Facts

Active Ingredients (in each capsule)

Omeprazole 20 mg.....Acid reducer
Sodium Bicarbonate 1100 mg.....Allows absorption of this omeprazole product

Use

- treat frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.

Warnings

Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food
- vomiting with blood
- bloody or black stools

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

LOT/EXP Area
No ink
No Varnish

0 141100 80673 4

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic

Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd. Bold Italic (continued) - 8 pt Helvetica Cnd.

Headings - 8 pt Helvetica Cnd. Bold Italic

Sub Heads - 6 pt Helvetica Cnd. Bold

Text - 6 pt Helvetica Cnd. minimum leading 6.5

Square Bullets - 3.5 pt Zaph Dingbats no compression

Border - 1 pt rule, 3/64" gap all around

Barline - 1 pt rule

Hairline - .5 pt rule

Certified by: eb

AP0 #: 2012-93465

02.23.12 - rp

3.1497

1.8111 [46.00]

4.7638 [121.00]

3.5434 [90.00]

Submitted 5/30/2012

This label may not be the latest approved by FDA. For current labeling information, please visit <https://www.fda.gov/drugsatfda>



03658506

905859E0



NO INK
NO VARNISH

NO INK
NO VARNISH

Drug Facts
Active Ingredients (in each capsule)
Omeprazole 20 mg.....Acid reducer
Sodium bicarbonate 1100 mg.....Allows absorption of this omeprazole product.

Use
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.

Warnings
Allergy alert: Do not use if you are allergic to omeprazole.
Do not use if you have:
■ trouble or pain swallowing food
■ vomiting with blood
■ bloody or black stools
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with **lightheadedness, swelling or dizziness**
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
■ frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Ask a doctor or pharmacist before use if you are taking
■ warfarin, clopidogrel or clostrazolil (blood-thinning medicines)
■ prescription antifungal or anti-yeast medicines

Drug Facts (continued)
■ diazepam (anxiety medicine)
■ digoxin (heart medicine)
■ tacrolimus (immune system medicine)
■ prescription antiretrovirals (medicines for HIV infection)
■ any other prescription drugs. Sodium bicarbonate may interact with certain prescription drugs.

Strip uses and ask a doctor if
■ you get diarrhea
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you are pregnant or breastfeeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ for adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
■ swallow 1 capsule with a glass of water at least 1 hour before eating in the morning
■ take every day for 14 days
■ do not take more than 1 capsule a day
■ do not chew or crush the capsule
■ do not open capsule and sprinkle on food
■ do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Drug Facts (continued)
■ children under 18 years of age: ask a doctor.
Heartburn in children may sometimes be caused by a serious condition.

Other information
■ each capsule contains sodium 303 mg
■ read the directions, warnings and accompanying label information before use
■ store at 20°-25°C (68°-77°F)
■ tamper-evident: do not use if the blue band around the capsule is missing or broken. Do not use if foil inner seal imprinted with "Sealed for your protection" is missing, open or broken.
■ keep product out of high heat and humidity
■ protect product from moisture

Inactive ingredients
croscarmellose sodium, FD&C blue No. 1, gelatin, indigotine, polyethylene glycol, polyethylene glycol 80, titanium dioxide

Questions or comments?
Call 1-888-4-TEC-OTC (1-888-483-4862) between 8:00 AM and 5:00 PM Central Time, Monday through Friday or visit www.ZegeridOTC.com

© Copyright & Distributed by MSD Consumer Care, Inc., PO Box 377, Memphis, TN 38151 USA, a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA. All rights reserved. Product of Spain.
36585-06/PKG01450
recyclable carton

Treats Frequent Heartburn

Zegerid[®] OTC



42 CAPSULES

NDX 11523-7265-3

Treats Frequent Heartburn

Zegerid[®] OTC

• Omeprazole 20 mg/Acid Reducer
• Sodium Bicarbonate 1100 mg!

Allows Absorption of this Omeprazole Product

42 CAPSULES

Three 14-Day Courses of Treatment

Lot/EXP Area
No Varnish
No Ink



0 141100 80677 2

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd, Bold Italic
Drug Facts (continued)

Drug Facts - 8 pt Helvetica Cnd, Bold Italic (continued) - 8 pt Helvetica Cnd.
Headings - 8 pt Helvetica Cnd, Bold Italic
Sub-Headings - 6 pt Helvetica Cnd, Bold
Text - 6 pt Helvetica Cnd, minimum leading 6.5
Square Bullets - 3.5 pt Zaph Dingbats no compression
Border - 1 pt rule, 3/64" gap all around
Barline - 1 pt rule
Hairline - .5 pt rule

Certified by: eb
APO #: 2012-93628
03.27.12 - ju

Submitted 3/20/2012


ATTACH LEGEND ON ALL
DIGITAL MECHANICALS
STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 37 characters
per inch
Drug Facts - 9 pt Helvetica Cond. Bold Italic

Treats Frequent Heartburn

Zegerid

OTC[®]

- Omeprazole 20 mg/Acid Reducer
- Sodium Bicarbonate 1100 mg/
Allows Absorption of this Omeprazole Product

<p>Drug Facts</p> <p>Active ingredients (in each capsule) Omeprazole 20 mg..... Sodium Bicarbonate 1100 mg..... Allows absorption of this omeprazole product</p> <p>Purpose Acid reducer</p>	<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you get diarrhea ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.
<p>Use</p> <ul style="list-style-type: none"> ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect. 	<p>Directions</p> <ul style="list-style-type: none"> ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ swallow 1 capsule with a glass of water at least 1 hour before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ do not chew or crush the capsule ■ do not open capsule and sprinkle on food ■ do not use for more than 14 days unless directed by your doctor <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months 	<p>Other information</p> <ul style="list-style-type: none"> ■ each capsule contains: sodium 383 mg ■ read the directions, warnings and accompanying label information before use ■ store at 20°-25°C (68°-77°F) ■ tamper-evident: do not use if the blue band around the capsule is missing or broken. Do not use if foil inner seal imprinted with "Sealed for your protection" is missing, open or broken. ■ keep product out of high heat and humidity ■ protect product from moisture
<p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to omeprazole</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food ■ vomiting with blood ■ bloody or black stools <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ a sodium-restricted diet 	<p>Inactive ingredients</p> <p>croscarmellose sodium, FD&C blue No. 1, gelatin, magnesium stearate, pharmaceutical ink, polysorbate 80, titanium dioxide</p> <p>Questions or comments? Call 1-888-4-ZEG-OTC (1-888-483-4862) between 8:00 AM and 5:00 PM Central Time, Monday through Friday or visit www.ZegeridOTC.com</p>	<p>Repeat 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months
<p>Ask a doctor or pharmacist before use if you are taking</p> <ul style="list-style-type: none"> ■ warfarin, clopidogrel or cilostazol (blood-thinning medicines) ■ prescription antifungal or anti-yeast medicines ■ diazepam (anxiety medicine) ■ digoxin (heart medicine) ■ tacrolimus (immune system medicine) ■ prescription antiretrovirals (medicines for HIV infection) ■ any other prescription drugs. Sodium bicarbonate may interact with certain prescription drugs. 	<p>NO INK NO VARNISH</p>	 <p>0 41100 80717 5</p>

© Copyright & Distributed by MSD Consumer Care, Inc., PO Box 377, Memphis, TN 38151 USA, a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA. All rights reserved. Product of Spain.

37821-04



42
CAPSULES
Three 14-Day
Courses of Treatment

Treats Frequent Heartburn

Zegerid

OTC[®]

- Omeprazole 20 mg/Acid Reducer
- Sodium Bicarbonate 1100 mg/
Allows Absorption of this Omeprazole Product

3 Pack

ITM# 807175

NDC 11523-7265-3

7.5000

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER
09/18/2012