


ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch
Drug Facts - 9 pt Helvetica Cnd. Bold Italic
Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd. Bold Italic (continued) - 8 pt Helvetica Cnd.
Headings - 8 pt Helvetica Cnd. Bold Italic
Sub Heads - 6 pt Helvetica Cnd. Bold
Text - 6 pt Helvetica Cnd, minimum leading 6.5
Square Bullets - 3.5 pt Zaph Dingbats no compression
Border - 1 pt rule, 3/64" gap all around
Barline - 1 pt rule _____
Hairline - .5 pt rule _____


Certified by: _____ (b) (4)
APO: 2017-XXXXXX
01.20.14 (b) (4)
04.24.17 (b) (4)




A - Top Panel




B - Back Panel




Base Panel




C - Insert 1 Top




D - Insert 1 Back




E - Insert 2 Top




F - Insert 2 Back




G - Insert 3 Top




H - Insert 3 Back




I - Insert 4 Top




J - Insert 4 Back




K - Insert 5 Top



L - Insert 5 Back



M - Insert 6 Top



N - Insert 6 Back

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic

Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd. Bold Italic (continued) - 8 pt Helvetica Cnd.

Headings - 8 pt Helvetica Cnd. Bold Italic

Sub Heads - 6 pt Helvetica Cnd. Bold

Text - 6 pt Helvetica Cnd, minimum leading 6.5

Square Bullets - 3.5 pt Zaph Dingbats no compression

Border - 1 pt rule, 3/64" gap all around


Barline - 1 pt rule _____

Hairline - .5 pt rule _____

Certified by: (b) (4)

APO #: 2017-XXXX
01.19.15 (b) (4)
04.25.17

80685950



NO INK
NO VARNISH

How Zegerid OTC Works For Your Frequent Heartburn

Zegerid OTC works differently from other OTC heartburn products, such as antacids and other acid reducers. Zegerid OTC stops acid production at the source - the acid pump that produces stomach acid. Zegerid OTC is to be used once a day (every 24 hours), every day for 14 days.

TIPS FOR MANAGING HEARTBURN

- Do not lie flat or bend over soon after eating
- Do not eat late at night or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose-fitting clothing around your stomach

Bayer

Drug Facts (continued)

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking

- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- diazepam (anxiety medicine)
- dioxin (heart medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)
- any other prescription drugs. Sodium bicarbonate may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- for adults 18 years of age and older
 - this product is to be used once a day (every 24 hours), every day for 14 days
 - it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- 14-Day Course of Treatment**
- swallow 1 capsule with a glass of water at least 1 hour before eating in the morning
 - take every day for 14 days
 - do not take more than 1 capsule a day
 - do not chew or crush the capsule
 - do not open capsule and sprinkle on food
 - do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information

- each capsule contains: sodium 303 mg
- read the directions, warnings and accompanying label information before use
- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if the blue band around the capsule is missing or broken. Do not use if foil inner seal imprinted with "Sealed for your protection" is missing, open or broken.
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients

croscarmellose sodium, FD&C blue No. 1, gelatin, magnesium stearate, pharmaceutical ink, polysorbate 80, titanium dioxide

Questions or comments?

Call 1-888-4-ZEG-OTC (1-888-493-4682) between 9:00 AM and 5:00 PM Eastern Standard Time, Monday through Friday or visit www.ZegeridOTC.com

Bayer and the Bayer Cross are registered trademarks of Bayer. Zegerid is a registered trademark of Salix Pharmaceuticals, Inc. used under license.

© 2015 Bayer. Dist by: Bayer HealthCare LLC, Whippany, NJ 07981 Product of Spain.

recyclable carton 36589-08/PKG01979

Treats Frequent Heartburn

Zegerid
OTC

- Omeprazole 20 mg/Acid Reducer
- Sodium Bicarbonate 1100 mg/ Allows Absorption of this Omeprazole Product



14
CAPSULES
One 14-Day Course of Treatment

Drug Facts

Active ingredients (in each capsule)

Omeprazole 20 mg..... Acid reducer
Sodium Bicarbonate 1100 mg.....
.....Allows absorption of this omeprazole product

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.

Warnings

Allergy alert: Do not use if you are allergic to omeprazole.

Do not use if you have:

- trouble or pain swallowing food
- vomiting with blood
- bloody or black stools

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

PRINTSIDE GLUE

FOLD

PRINTSIDE GLUE

NO INK
NO VARNISH



LOT/EXP
Area
No ink
No Varnish

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
10/30/2017