

Drug Facts (continued)

neck or shoulders, or lightheadedness

- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water at least 1 hour before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- do not chew or crush the capsule
- do not open capsule and sprinkle on food
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor.

Heartburn in children may sometimes be caused by a serious condition.

Other information

- each capsule contains: sodium 303 mg
- read the directions, warnings and accompanying label information before use
- store at 20°-25°C (68°-77°F)

Drug Facts (continued)

- tamper-evident: do not use if the blue band around the capsule is missing or broken. Do not use if full inner seal imprinted with "Sealed for your protection" is missing, open or broken
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients
FD&C blue no. 1, FD&C blue no. 2 aluminum lake, FD&C red no. 40, gelatin, polysorbate 80, sodium lauryl sulfate, sodium starch glycolate, sodium stearyl fumarate, titanium dioxide

Questions or comments?
Call 1-888-4-ZEG-OTC (1-888-493-4682) between 9:00 AM and 5:00 PM Eastern Standard Time, Monday through Friday

Dist. by: Riley Consumer Care
New York, New York 10151

recyclable carton

HOW ZEGERID OTC WORKS FOR YOUR FREQUENT HEARTBURN

Zegerid OTC works differently from other OTC heartburn products, such as antacids and other acid reducers. Zegerid OTC stops acid production at the source - the acid pump that produces stomach acid. Zegerid OTC is to be used once a day (every 24 hours), every day for 14 days.

TIPS FOR MANAGING HEARTBURN

- Do not lie flat or bend over soon after eating
- Do not eat late at night or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose-fitting clothing around your stomach

Drug Facts

Active ingredients (in each capsule)	Purpose
Omeprazole 20 mg	Acid reducer
Sodium Bicarbonate 1100 mg Allows absorption of this omeprazole product

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.

Warnings

Allergy alert:

- do not use if you are allergic to omeprazole.
- omeprazole may cause severe skin reactions. Symptoms may include:
 - skin redness
 - blisters
 - rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms,

CLINICALLY PROVEN TO
Treat Frequent Heartburn

Zegerid
OTC

- Omeprazole 20mg/Acid Reducer
- Sodium Bicarbonate 1100mg /Allows Absorption of this Omeprazole Product

14
CAPSULES

24 HOUR

One 14-Day Course of Treatment
MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT



0 4 1100 80673 4

1H9D5 BY C3

Drug Facts (continued)

- neck or shoulders; or lightheadedness
- frequent chest pain**
These may be signs of a serious condition. See your doctor.
- Ask a doctor before use if you have**
 - had heartburn over 3 months. This may be a sign of a more serious condition.
 - frequent wheezing, particularly with heartburn
 - unexplained weight loss
 - nausea or vomiting
 - stomach pain
 - a sodium-restricted diet
- Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.
- Stop use and ask a doctor if**
 - your heartburn continues or worsens
 - you need to take this product for more than 14 days
 - you need to take more than 1 course of treatment every 4 months
 - you get diarrhea
 - you develop a rash or joint pain
- If pregnant or breast-feeding,** ask a health professional before use.
- Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water at least 1 hour before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- do not chew or crush the capsule
- do not open capsule and sprinkle on food
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor.
- Heartburn in children may sometimes be caused by a serious condition.

Other information

- each capsule contains: **sodium 303 mg**
- read the directions, warnings and accompanying label information before use
- store at 20°-25°C (68°-77°F)

Drug Facts (continued)

- tamper-evident: do not use if the blue band around the capsule is missing or broken. Do not use if foil inner seal imprinted with "Sealed for your protection" is missing, open or broken.
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients
FD&C blue no. 1, FD&C blue no. 2 aluminum lake, FD&C red no. 40, gelatin, polysorbate 80, sodium lauryl sulfate, sodium starch glycolate, sodium stearyl fumarate, titanium dioxide

Questions or comments?
Call 1-888-4-ZEG-OTC (1-888-493-4682) between 9:00 AM and 5:00 PM Eastern Standard Time, Monday through Friday

Dist. by: Riley Consumer Care
New York, New York 10151

recyclable carton

HOW ZEGERID OTC WORKS FOR YOUR FREQUENT HEARTBURN

Zegerid OTC works differently from other OTC heartburn products, such as antacids and other acid reducers. Zegerid OTC stops acid production at the source - the acid pump that produces stomach acid. Zegerid OTC is to be used once a day (every 24 hours), every day for 14 days.

TIPS FOR MANAGING HEARTBURN

- Do not lie flat or bend over soon after eating
- Do not eat late at night or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some fruits and vegetables
- Est slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose-fitting clothing around your stomach

Drug Facts

Active ingredients (in each capsule)	Purpose
Omeprazole 20 mg	Acid reducer
Sodium Bicarbonate 1100 mg Allows absorption of this omeprazole product

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.

Warnings

Allergy alert:

- do not use if you are allergic to omeprazole.
- omeprazole may cause severe skin reactions. Symptoms may include:
 - skin reddening
 - blisters
 - rash
- If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms.

CLINICALLY PROVEN TO
Treat Frequent Heartburn

Zegerid

OTC

- Omeprazole 20mg/Acid Reducer
- Sodium Bicarbonate 1100mg
- Allows Absorption of this Omeprazole Product

28
CAPSULES

Two 14-Day Courses of Treatment
MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

24 HOUR

1H9D7 BY C3

FPO UPC

0 00000 00000 0

0 00000 00000 0

Drug Facts	Purpose
Active ingredients (in each capsule)	
Omeprazole 20 mg.....	Acid reducer
Sodium Bicarbonate 1100 mg.....	Allows absorption of this omeprazole product
Use	
<ul style="list-style-type: none"> ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect. 	
Warnings	
Allergy alert:	
<ul style="list-style-type: none"> ■ do not use if you are allergic to omeprazole. ■ omeprazole may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> ■ skin reddening ■ blisters ■ rash 	
If an allergic reaction occurs, stop use and seek medical help right away.	
Do not use if you have:	
<ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain 	
These may be signs of a serious condition. See your doctor.	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn 	

Drug Facts (continued)
<ul style="list-style-type: none"> ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ a sodium-restricted diet
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
Stop use and ask a doctor if
<ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
Directions
<ul style="list-style-type: none"> ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
14-Day Course of Treatment
<ul style="list-style-type: none"> ■ swallow 1 capsule with a glass of water at least 1 hour before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ do not chew or crush the capsule ■ do not open capsule and sprinkle on food ■ do not use for more than 14 days unless directed by your doctor

Drug Facts (continued)
Repeated 14-Day Courses (if needed)
<ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
<ul style="list-style-type: none"> ■ children under 18 years of age: ask a doctor. ■ Heartburn in children may sometimes be caused by a serious condition.
Other information
<ul style="list-style-type: none"> ■ each capsule contains: sodium 303 mg ■ read the directions, warnings and accompanying label information before use ■ store at 20°-25°C (68°-77°F) ■ tamper-evident: do not use if the blue band around the capsule is missing or broken. Do not use if foil inner seal imprinted with "Sealed for your protection" is missing, open or broken. ■ keep product out of high heat and humidity ■ protect product from moisture
Inactive ingredients
FD&C blue no. 1, FD&C blue no. 2 aluminum lake, FD&C red no. 40, gelatin, polysorbate 80, sodium lauryl sulfate, sodium starch glycolate, sodium stearyl fumarate, titanium dioxide
Questions or comments?
Call 1-888-4-ZEG-OTC (1-888-493-4682) between 9:00 AM and 5:00 PM Eastern Standard Time, Monday through Friday
Dist. by: Riley Consumer Care New York, New York 10151

CLINICALLY PROVEN TO

Treat Frequent Heartburn

Zegerid

OTC

42 CAPSULES

CLINICALLY PROVEN TO

Treat Frequent Heartburn

Zegerid

OTC

- Omeprazole 20mg/Acid Reducer
- Sodium Bicarbonate 1100mg /Allows Absorption of this Omeprazole Product

24 HOUR

42 CAPSULES

Three 14-Day Courses of Treatment

MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

HOW ZEGERID OTC WORKS FOR YOUR FREQUENT HEARTBURN

Zegerid OTC works differently from other OTC heartburn products, such as antacids and other acid reducers. Zegerid OTC stops acid production at the source - the acid pump that produces stomach acid. Zegerid OTC is to be used once a day (every 24 hours), every day for 14 days.

TIPS FOR MANAGING HEARTBURN

- Do not lie flat or bend over soon after eating
- Do not eat late at night or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose-fitting clothing around your stomach

0 41100 80677 2

1H9D7 BY C3

CLINICALLY PROVEN TO
Treat Frequent Heartburn

Zegerid[®] OTC

• Omeprazole 20 mg/Acid Reducer
• Sodium Bicarbonate 1100 mg
/Allows Absorption of this
Omeprazole Product

14
CAPSULES
ONE 14-DAY COURSE OF TREATMENT

Tamper-evident: Do not use if the band around the capsule is missing or broken. Do not use if printed seal under cap is broken or missing.

Distributed by
Riley Consumer Care
New York, New York 10151

: 1H9D5 XL F1

Open to read directions and additional drug facts information. Keep outer carton for complete warnings and product information.

Active ingredients (in each capsule)
Omeprazole 20 mg Acid reducer
Sodium bicarbonate 1100 mg Allows absorption of this omeprazole product

Purpose
Zegerid OTC (omeprazole) reduces the amount of acid in the stomach. It helps relieve heartburn (occurs 2 or more days a week). Zegerid OTC does not provide relief of heartburn. This drug may take 1 to 4 days for full effect.

Warnings
Allergy alert: Do not use if you are allergic to omeprazole.

PEEL BACK HERE →

■ omeprazole may cause severe side effects. Symptoms may include: ■ swollen, red, itchy skin ■ rash ■ hives ■ difficulty breathing ■ difficulty swallowing ■ dizziness ■ fainting ■ blurred vision ■ watery eyes ■ headache ■ dizziness ■ ringing in ears ■ changes in voice ■ changes in taste ■ changes in smell ■ changes in vision ■ changes in hearing ■ changes in touch ■ changes in taste ■ changes in smell ■ changes in vision ■ changes in hearing ■ changes in touch

Do not use if you have:
■ trouble or pain swallowing food, or weight loss, or bloating, or heartburn, or ■ heartburn with ■ lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders or lightheadedness ■ frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ loss of appetite ■ loss of energy ■ a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you feel more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or your pain ■ if present or breast-feeding, ask a health professional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.
■ for adults 18 years of age and older:
■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief during the first 2 to 3 days of treatment

14-Day Course of Treatment
■ swallow 1 capsule with a glass of water at least 1 hour before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ do not chew or crush the capsule ■ do not open capsule and sprinkle on food ■ do not use for more than 14 days unless directed by your doctor

Other information
■ each capsule contains sodium 303 mg

Questions? Call 1-888-493-4892

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
04/01/2024 09:33:46 AM