




Modified format

- Characters per inch: < 39
- Headings: 6pt
- Subheadings: 5.5pt
- Text: 5.5pt
- Text leading: 6pt
- Square bullets: 3.5pt
- Borders: 1pt, 3/64" gap all around
- Barline: 1pt



DRUG FACTS INFO	
CATEGORY	TYPE SIZE
Drug Facts Title	9 pt
Drug Facts Continued	9 pt
Headings	8 pt
Subheadings	6 pt
Body Text	6 pt
Bullets	5 pt
Hairlines	0.5 pt rule
Box	1 pt rule

DRUG FACTS INFO	
CATEGORY	TYPE SIZE
Drug Facts Title	9.5g
Drug Facts Download	8.5g
Headings	8.5g
Subheadings	8.5g
Body Text	8.5g
Images	3.5g
Warnings	0.5g (Full)
Box	1.0g (Full)

14

TREATS FREQUENT HEARTBURN

Zegerid^{OTC}

• Omeprazole 20mg/Acid Reducer
• Sodium Bicarbonate 1680mg/Allows
Absorption of this Omeprazole Product
Powder for Oral Suspension

14

TREATS FREQUENT HEARTBURN

14

TREATS FREQUENT HEARTBURN

Zegerid^{OTC}

• Omeprazole 20mg/Acid Reducer
• Sodium Bicarbonate 1680mg/Allows
Absorption of this Omeprazole Product
Powder for Oral Suspension

14

TREATS FREQUENT HEARTBURN

14

TREATS FREQUENT HEARTBURN

Zegerid^{OTC}

• Omeprazole 20mg/Acid Reducer
• Sodium Bicarbonate 1680mg/Allows
Absorption of this Omeprazole Product
Powder for Oral Suspension

14

TREATS FREQUENT HEARTBURN

14

TREATS FREQUENT HEARTBURN

Zegerid^{OTC}

• Omeprazole 20mg/Acid Reducer
• Sodium Bicarbonate 1680mg/Allows
Absorption of this Omeprazole Product
Powder for Oral Suspension

14

TREATS FREQUENT HEARTBURN

14

TREATS FREQUENT HEARTBURN

Zegerid^{OTC}

• Omeprazole 20mg/Acid Reducer
• Sodium Bicarbonate 1680mg/Allows
Absorption of this Omeprazole Product
Powder for Oral Suspension

14

TREATS FREQUENT HEARTBURN

14

TREATS FREQUENT HEARTBURN

Zegerid^{OTC}

• Omeprazole 20mg/Acid Reducer
• Sodium Bicarbonate 1680mg/Allows
Absorption of this Omeprazole Product
Powder for Oral Suspension

14

TREATS FREQUENT HEARTBURN

Drug Facts

Active Ingredients (in each powder packet)
Omeprazole 20 mg Acid reducer
Sodium Bicarbonate 1680 mg
Allows absorption of this omeprazole product

Purpose
Acid reducer

Use
• treats frequent heartburn (occurs 2 or more days a week)
• not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.

Warnings
Allergy alert: do not use if you are allergic to omeprazole
• omeprazole may cause severe skin reactions. Symptoms may include:
• skin reddening
• blisters
• rash.
If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:
• trouble or pain swallowing food, vomiting with blood, or bloody or black stools
• heartburn with lightheadedness, sweating or dizziness
• chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
• frequent chest pain
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
• had heartburn over 3 months. This may be a sign of a more serious condition.

Drug Facts (continued)

• frequent wheezing, particularly with heartburn
• unexplained weight loss
• nausea or vomiting
• stomach pain
• a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
• your heartburn continues or worsens
• you need to take this product for more than 14 days
• you need to take more than 1 course of treatment every 4 months
• you get diarrhea
• you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

14-Day Course of Treatment
• product should be taken at least 1 hour before eating in the morning
• take every day for 14 days
• do not take more than once a day
• empty packet contents into a small cup containing 2 tablespoons of WATER
• DO NOT USE WITH OTHER LIQUIDS OR FOOD
• stir well and drink immediately
• refill cup with water and drink
• do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
• you may repeat a 14-day course every 4 months
• do not take for more than 14 days or more often than every 4 months unless directed by a doctor

• children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Other information
• each powder packet contains sodium 450 mg
• read the directions and warnings before use
• keep the carton. It contains important information.
• store between 20° to 25°C (68° to 77°F)
• tamper-evident: do not use if foil is open or broken
• keep product out of high heat and humidity
• protect product from moisture


Inactive ingredients flavor, sucralose, sucrose, xanthan gum, xylitol

Questions or comments?
Call 1-888-4-ZEG-OTC (1-888-483-4682) between 9:00 AM and 5:00 PM Eastern Standard Time, Monday through Friday or visit www.ZegeridOTC.com

Dist. by Riley Consumer Care
New York, New York 10151

recyclable carton

LOT
EXP
NO INK NO VARNISH



HOW ZEGERID OTC WORKS FOR YOUR FREQUENT HEARTBURN

Zegerid OTC works differently from other OTC heartburn products, such as antacids and other acid reducers. Zegerid OTC stops acid production at the source - the acid pump that produces stomach acid. Zegerid OTC is to be used once daily every 24 hours every day for 14 days.

TIPS FOR MANAGING HEARTBURN

- Do not lie flat or bend over soon after eating
- Do not eat late at night or just before bedtime
- Certain foods, drinks or items (such as spicy foods, heartburn, such as soft, spicy, fatty and fried foods, chocolate, caffeine, alcohol) can cause heartburn and even some flat and legumes
- Eat slowly and do not eat too much
- If you are overweight, lose weight
- Raise the head of your bed
- Wear loose-fitting clothing around your stomach

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

JODY E GREEN
11/23/2022 08:00:30 AM