

NDC 0078-0684-15

Rx only

Promacta[®] (eltrombopag) Tablets 12.5 mg

Swallow tablets whole. Do not split, chew, or crush tablets.

Dispense with Medication Guide attached or provided separately.

 NOVARTIS 30 Tablets

Store at room temperature between 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). Dispense in original bottle.

Do not use if printed safety seal under cap is broken or missing.

Dosage: See accompanying Prescribing Information.

Important: Use safety closures when dispensing this product unless otherwise directed by physicians or requested by purchaser.

KEEP OUT OF THE REACH OF CHILDREN

Each tablet contains eltrombopag olamine equivalent to 12.5 mg of eltrombopag free acid.

Mfd. by: Novartis d.o.o.

Ljubljana, 1000, Slovenia

Dist. by: Novartis Pharmaceuticals Corp.

East Hanover, New Jersey 07936

Product of Slovenia

© Novartis

300107500 US



GTIN 12345678901234
SN 00000000000000
LOT ZZ9999
EXP YYYY-MM-DD



NDC 0078-0685-15

Rx only

Promacta[®] (eltrombopag) Tablets 25 mg

Swallow tablets whole. Do not split, chew, or crush tablets.

Dispense with Medication Guide attached or provided separately.

 NOVARTIS 30 Tablets

Store at room temperature between 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). Dispense in original bottle.

Do not use if printed safety seal under cap is broken or missing.

Dosage: See accompanying Prescribing Information.

Important: Use safety closures when dispensing this product unless otherwise directed by physicians or requested by purchaser.

KEEP OUT OF THE REACH OF CHILDREN
Each tablet contains eltrombopag olamine equivalent to 25 mg of eltrombopag free acid.

Mfd. by: Novartis d.o.o.
Ljubljana, 1000, Slovenia
Dist. by: Novartis Pharmaceuticals Corp.
East Hanover, New Jersey 07936
Product of Slovenia



© Novartis
300107491 US

GTIN 12345678901234
SN 00000000000000
LOT ZZ9999
EXP YYYY-MM-DD



Sample-Not for Sale R only
NDC 0078-9686-55

Promacta[®]
(eltrombopag) Tablets
50 mg

Swallow tablets whole. Do not split, chew, or crush tablets.

Dispense with Medication Guide attached or provided separately.

 **NOVARTIS** 14 Tablets

Store at room temperature between 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F).

Dispense in original bottle.

Do not use if printed safety seal under cap is broken or missing.

Dosage: See accompanying Prescribing Information.

Important: Use safety closures when dispensing this product unless otherwise directed by physicians or requested by purchaser.

KEEP OUT OF THE REACH OF CHILDREN

Each tablet contains eltrombopag olamine equivalent to 50 mg of eltrombopag free acid.

Mfd. by: Novartis d.o.o.
Ljubljana, 1000, Slovenia
Dist. by: Novartis Pharmaceuticals Corp.
East Hanover, New Jersey 07936
Product of Slovenia

© Novartis
300107502 US



LOI XXXXXX
EXP YYY-MM-DD

Sample-Not for Sale
NDC 0078-9686-55

Rx only

Promacta[®]
(eltrombopag) Tablets
50 mg

Swallow tablets whole. Do not split, chew, or crush tablets.

Dispense with Medication Guide attached or provided separately.

 NOVARTIS 14 Tablets

Store at room temperature between 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F).

Dispense in original bottle.

Do not use if printed safety seal under cap is broken or missing.

Dosage: See accompanying Prescribing Information.

Important: Use safety closures when dispensing this product unless otherwise directed by physicians or requested by purchaser.

KEEP OUT OF THE REACH OF CHILDREN

Each tablet contains eltrombopag olamine equivalent to 50 mg of eltrombopag free acid.

Mfd. by: Siegfried Barbera, S.L.
Barberá del Vallés, Spain
Dist. by: Novartis Pharmaceuticals Corp.
East Hanover, New Jersey 07936

Product of Slovenia

© Novartis
300107488 US



LOT XXXXXX
EXP YYY-YY-YY

NDC 0078-0686-55

Rx only

Promacta[®] (eltrombopag) Tablets 50 mg

Swallow tablets whole. Do not split, chew, or crush tablets.

Dispense with Medication Guide attached or provided separately.

 NOVARTIS 14 Tablets

Store at room temperature between 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). Dispense in original bottle.

Do not use if printed safety seal under cap is broken or missing.

Dosage: See accompanying Prescribing Information.

Important: Use safety closures when dispensing this product unless otherwise directed by physicians or requested by purchaser.

KEEP OUT OF THE REACH OF CHILDREN
Each tablet contains eltrombopag olamine equivalent to 50 mg of eltrombopag free acid.

Mfd. by: Novartis d.o.o.,
Ljubljana, 1000, Slovenia
Dist. by: Novartis Pharmaceuticals Corp.,
East Hanover, New Jersey 07936
Product of Slovenia

© Novartis
300107504 US



GTIN 12345678901234
SN 00000000000000
LOT ZZ9999
EXP YYYY-MM-DD



NDC 0078-0686-55

Rx only

Promacta[®] (eltrombopag) Tablets 50 mg

Swallow tablets whole. Do not split, chew, or crush tablets.

Dispense with Medication Guide attached or provided separately.

 NOVARTIS 14 Tablets

Store at room temperature between 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). Dispense in original bottle.

Do not use if printed safety seal under cap is broken or missing.

Dosage: See accompanying Prescribing Information.

Important: Use safety closures when dispensing this product unless otherwise directed by physicians or requested by purchaser.

KEEP OUT OF THE REACH OF CHILDREN

Each tablet contains eltrombopag olamine equivalent to 50 mg of eltrombopag free acid.

Mfd. by: Siegfried Barbera, S.L.
Barberà del Valles, Spain
Dist. by: Novartis Pharmaceuticals Corp.
East Hanover, New Jersey 07936
Product of Slovenia

© Novartis

300107490 US



GTIN 12345678901234
SN 00000000000000
LOT ZZ9999
EXP YYYY-MM-DD



NDC 0078-0686-15

Rx only

Promacta[®] (eltrombopag) Tablets 50 mg

Swallow tablets whole. Do not split, chew, or crush tablets.

Dispense with Medication Guide attached or provided separately.

 NOVARTIS 30 Tablets

Store at room temperature between 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). Dispense in original bottle.

Do not use if printed safety seal under cap is broken or missing.

Dosage: See accompanying Prescribing Information.

Important: Use safety closures when dispensing this product unless otherwise directed by physicians or requested by purchaser.

KEEP OUT OF THE REACH OF CHILDREN

Each tablet contains eltrombopag olamine equivalent to 50 mg of eltrombopag free acid.

Mfd. by: Novartis d.o.o.
Ljubljana, 1000, Slovenia
Dist. by: Novartis Pharmaceuticals Corp.
East Hanover, New Jersey 07936
Product of Slovenia



© Novartis
300107501 US

GTIN 12345678901234
SN 00000000000000
LOT ZZ9999
EXP YYYY-MM-DD



NDC 0078-0686-15

Rx only

Promacta[®] (eltrombopag) Tablets 50 mg

Swallow tablets whole. Do not split, chew, or crush tablets.

Dispense with Medication Guide attached or provided separately.

 NOVARTIS 30 Tablets

Store at room temperature between 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). Dispense in original bottle.

Do not use if printed safety seal under cap is broken or missing.

Dosage: See accompanying Prescribing Information.

Important: Use safety closures when dispensing this product unless otherwise directed by physicians or requested by purchaser.

KEEP OUT OF THE REACH OF CHILDREN
Each tablet contains eltrombopag olamine equivalent to 50 mg of eltrombopag free acid.

Mfd. by: Siegfried Barbra, S.L.,
Barberá del Valles, Spain
Dist. by: Novartis Pharmaceuticals Corp.,
East Hanover, New Jersey 07936
Product of Slovenia

© Novartis
300107489 US



GTIN 1324578901234
SN 0000000000000
LOT Z2999
EXP YYYT-WWH-DD



NDC 0078-0687-15

Rx only

Promacta[®] (eltrombopag) Tablets 75 mg

Swallow tablets whole. Do not split, chew, or crush tablets.

Dispense with Medication Guide attached or provided separately.

 NOVARTIS 30 Tablets

Store at room temperature between 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F). Dispense in original bottle.

Do not use if printed safety seal under cap is broken or missing.

Dosage: See accompanying Prescribing Information.

Important: Use safety closures when dispensing this product unless otherwise directed by physicians or requested by purchaser.

KEEP OUT OF THE REACH OF CHILDREN
Each tablet contains eltrombopag oleamine equivalent to 75 mg of eltrombopag free acid.

Mfd. by: Novartis d.o.o., Ljubljana, 1000, Slovenia
Dist. by: Novartis Pharmaceuticals Corp., East Hanover, New Jersey 07936
Product of Slovenia

© Novartis
300107503 US



GTIN 12345678901234
SN 00000000000000
LOT ZZ9999
EXP YYYY-MM-DD





Ramesh
Raghavachari

Digitally signed by Ramesh Raghavachari

Date: 12/27/2024 04:27:19PM

GUID: 502d0913000029f375128b0de8c50020