

PREVACID[®] 24HR
 Lansoprazole delayed-release capsules 15 mg acid reducer
14 CAPSULES
 ONE 14-DAY COURSE
 OF TREATMENT

1 BOTTLE INSIDE
14 CAPSULES TOTAL

PREVACID[®] 24HR
 Lansoprazole delayed-release capsules 15 mg / acid reducer
Actual Size
 ONE 14-DAY COURSE OF TREATMENT

SAVE \$X.XX NOW
 Sodium Free

PREVACID[®] 24HR
 Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- Clinically Proven To Treat Frequent Heartburn

14 CAPSULES
 ONE 14-DAY COURSE OF TREATMENT

Drug Facts (continued)

Other information

- read the directions, warnings and package insert, before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
 1-800-452-0051
 Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Drug Facts

Active ingredient (in each capsule)
 Lansoprazole 15 mg.....Acid reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)

Drug Facts (continued)

- theophylline (asthma medicine)
- tacrolimus (immune system medicine)
- atazanavir (medicine for HIV infection)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses. (If needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

NOVARTIS

Distributed by Novartis Consumer Health, Inc.
 Parsippany, NJ 07054-0022 ©2010 XXXXX
 PREVACID[®] is a registered trademark of Novartis Pharmaceuticals North America, Inc. and is used under license.

FPO

3 0067-6286-14 3



(b) (4)



SAVE \$X NOW

PREVACID[®] 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer

• May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

• Clinically Proven To Treat Frequent Heartburn

14 CAPSULES
ONE 14-DAY COURSE OF TREATMENT

Sodium Free

NDC 0067-6286-14

1 BOTTLE INSIDE
14 CAPSULES TOTAL

PREVACID[®] 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer

Actual Size
ONE 14-DAY COURSE OF TREATMENT

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Drug Facts (continued)

- theophylline (asthma medicine)
- lactulose (immune system medicine)
- atazanavir (medicine for HIV infection)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses. (If needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts

Active ingredient (in each capsule) Lansoprazole 15 mg, Acid reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)

NOVARTIS

Distributed by Novartis Consumer Health, Inc.
 Parsippany, NJ 07054-0822 ©2010 XXXXX
 PREVACID[®] is a registered trademark of Takeda Pharmaceuticals North America, Inc. and is used under license.

FPO



3 0067-6286-14 3



(b) (4)



3 0067-6286-14 3



Drug Facts

Active ingredient (in each capsule)	Purpose
Lansoprazole 15 mg	Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus (immune system medicine)
- atazanavir (medicine for HIV infection)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

Drug Facts (continued)

Directions

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeat 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of light and humidity
- protect product from moisture

PREVACID 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer

• May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

• Clinically Proven To Treat Frequent Heartburn

14 CAPSULES

ONE 14-DAY COURSE OF TREATMENT

SAVE \$X.XX NOW

Sodium Free

PER HERE

ND-C-0067-6286-14

Drug Facts (continued)

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

Visit us at www.prevacid24hr.com

Distributed by: Novartis Consumer Health, Inc.
Parsippany, NJ 07054-0622

©2010 XXXXX

PREVACID® is a registered trademark of Takeda Pharmaceuticals North America, Inc. and is used under license.

LOT:

EXP:

MADE IN:

Drug Facts (continued)

Directions

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeat 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of light and humidity
- protect product from moisture

PREVACID 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer

• May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

• Clinically Proven To Treat Frequent Heartburn

14 CAPSULES

ONE 14-DAY COURSE OF TREATMENT

SAVE \$X.XX NOW

Sodium Free

PER HERE

ND-C-0067-6286-14

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE

DO NOT USE IF INNER FOL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

FPO


3 0067-6286-15 0



(b) (4)

3 0067-6286-15 0

(b) (4)



KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAPE-REMOVED BOTTLE:
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Drug Facts

Active ingredient (in each capsule) Purpose
Lansoprazole 15 mg Acid reducer

Use
• treats frequent heartburn (occurs 2 or more days a week)
• not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to lansoprazole

Do not use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools; these may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
• liver disease
• had heartburn over 3 months. This may be a sign of a more serious condition.

Drug Facts (continued)

- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss
- nausea or vomiting • stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine) • prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus (immune system medicine)
- atazanavir (medicine for HIV infection)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

• children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information. • store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture



Drug Facts (continued)

Inactive ingredients: colloidal silicon dioxide, D&C red No. 28, FD&C Blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, polyethylene 80, starch, sucrose, sugar spheres, etc., titanium dioxide

Questions or comments? 1-800-452-0051

Visit us at www.prevacid24hr.com

PREVACID 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer



- May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- Clinically Proven To Treat Frequent Heartburn

14 CAPSULES
ONE 14-DAY COURSE OF TREATMENT

SAVE \$X.XX NOW

PER HERE

Sodium Free



NOVARTIS

Distributed by: Novartis Consumer Health, Inc.
Parsippany, NJ 07054-0222 ©2010 NOVARTIS

PREVACID[®] is a registered trademark of Novartis Pharmaceuticals (Novartis Consumer Health, Inc.) and is used under license.


PP0

3 0067-6286-15 0

LOT: :dXK

THE TITLE CTN PREV 14CT CLPSTRP HUM/CAT

NDC 0067-6286-14



KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Warnings
 • Do not use if you are allergic to lansoprazole

Do not use
 • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools; these may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
 • liver disease
 • had heartburn over 3 months. This may be a sign of a more serious condition.

Drug Facts
Active ingredient (in each capsule) Purpose
 Lansoprazole 15 mg Acid reducer

Use
 • treats frequent heartburn (occurs 2 or more days a week)
 • not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.

Drug Facts (continued)
 • heartburn with **lightheadedness, sweating or dizziness**
 • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 • frequent **chest pain** • frequent wheezing, particularly with heartburn • unexplained weight loss
 • nausea or vomiting • stomach pain

Ask a doctor or pharmacist before use if you are taking
 • warfarin (blood-thinning medicine) • prescription antifungal or anti-yeast medicines
 • digoxin (heart medicine)
 • theophylline (asthma medicine)
 • tacrolimus (immune system medicine)
 • atazanavir (medicine for HIV infection)

Stop use and ask a doctor if
 • your heartburn continues or worsens
 • you need to take this product for more than 14 days
 • you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 • adults 18 years of age and older
 • this product is to be used once a day (every 24 hours), every day for 14 days
 • it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 • swallow 1 capsule with a glass of water before eating in the morning
 • take every day for 14 days
 • do not take more than 1 capsule a day
 • swallow whole. Do not crush or chew capsules.
 • do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
 • you may repeat a 14-day course every 4 months
do not take for more than 14 days or more often than every 4 months unless directed by a doctor

• children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information
 • read the directions, warnings and package insert before use
 • keep the carton and package insert. They contain important information. • store at 20-25°C (68-77°F)
 • keep product out of high heat and humidity
 • protect product from moisture

1 BOTTLE INSIDE 14 CAPSULES TOTAL

PREV/ACID^{24HR}
 lansoprazole delayed-release capsules 15 mg / acid reducer

Actual Size
 ONE 14-DAY COURSE OF TREATMENT

SAVE \$X NOW!

PREV/ACID^{24HR}
 lansoprazole delayed-release capsules 15 mg / acid reducer

• May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
 • Clinically Proven To Treat Frequent Heartburn

14 CAPSULES
 ONE 14-DAY COURSE OF TREATMENT

PREV/ACID^{24HR}
 Sodium Free

Drug Facts (continued)
Inactive ingredients: colloidal silicon dioxide, D&C red No. 28, FD&C Blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, polymeric 80, starch, sucrose, sugar sphere, etc., titanium dioxide

Questions or comments? 1-800-452-0051

Visit us at www.lansoprazole.com

NOVARTIS
 Distributed by: Novartis Consumer Health, Inc.
 Parsippany, NJ 07654-0222 ©2010 NOVARTIS
 PREV/ACID^{24HR} is a registered trademark of Novartis Pharmaceuticals (Novartis Consumer Health, Inc. and its user) under license.

PP0

3 0067-6286-15 0


LOT:
 EX:
 IN: 031910

NDC 0067-6286-14

THE TITLE CTN PREV 14CT CLPSTRP HUM/CAT



(b) (4)



Drug Facts

Active ingredient (in each capsule)	Purpose
Lansoprazole 15 mg	Acid reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus (immune system medicine)
- atazanavir (medicine for HIV infection)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

Drug Facts (continued)

Directions

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor
- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition

Other Information

- read the directions, warnings and package insert before use
- keep the carton and package insert
- these contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- product protected from moisture



Drug Facts (continued)

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

Visit us at www.prevacid24hr.com
Distributed by: Novartis Consumer Health, Inc. Parsippany, NJ 07054-0622
©2010 XXXXX
PREVACID® is a registered trademark of Takeda Pharmaceuticals North America, Inc. and is used under license.

ONE 14-DAY COURSE OF TREATMENT



• May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

• Clinically Proven To Treat Frequent Heartburn

PREVACID 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer

SAVE \$X NOW!

Sodium Free

NDC 0087-6286-14

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF THIS POLY SEAL, IMPRINTED IN RED AND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.



PREVACID[®] 24HR
Lansoprazole delayed-release capsules 15 mg acid reducer
28 CAPSULES
TWO 14-DAY COURSES
OF TREATMENT

PREVACID[®] 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer
Actual Size
ONE 14-DAY COURSE OF TREATMENT

28 CAPSULES
TWO 14-DAY COURSES OF TREATMENT

SAVE \$X.XX NOW
Sodium Free

May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

Clinically Proven To Treat Frequent Heartburn

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051
Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051
Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051
Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051
Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051
Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051
Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.



3 0067-6286-28 0

3 PACK

SAVE \$X.XX NOW

NDC 0067-6286-42

PREVACID[®] / 24HR

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- Clinically Proven To Treat Frequent Heartburn

3 14-DAY COURSES OF TREATMENT

42 CAPSULES

Sodium Free

(b) (4)

The image shows a stylized graphic of a Prevacid 24HR packaging box. The box is primarily teal and pink. At the top left, a yellow banner says "3 PACK". To its right, a yellow banner says "SAVE \$X NOW". In the top right corner, the NDC number "NDC 0067-6286-42" is printed. The product name "PREVACID" is written in large, bold, white letters with a black outline, and "24HR" is written in smaller white letters above it. Below the product name, the text "Lansoprazole delayed-release capsules 15 mg / acid reducer" is written in a smaller font. A list of two bullet points is positioned below the product name: "• May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours" and "• Clinically Proven To Treat Frequent Heartburn". A large, empty rectangular box is outlined in the center of the packaging. At the bottom left, a large white number "3" is followed by the text "14-DAY COURSES OF TREATMENT". Below that, the text "42 CAPSULES" is written. In the bottom right corner, the text "Sodium Free" is visible. A small vertical text "XXXX" is located near the bottom right corner of the box graphic.



Walmart
Save money. Live better.
1-800-WAL-MART

If this product is found for sale at a location other than Walmart, please call 1-800-WAL-MART

PREV/ACID[®]
Lansoprazole delayed-release capsules 15 mg / acid reducer

42 CAPSULES
THREE 14-DAY COURSES OF TREATMENT

3 BOTTLES INSIDE

42 CAPSULES TOTAL

PREV/ACID[®]
Lansoprazole delayed-release capsules 15 mg / acid reducer

Actual Size
ONE 14-DAY COURSE OF TREATMENT

NDC 0067-6286-42

PREV/ACID

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- Clinically Proven To Treat Frequent Heartburn

42 CAPSULES

THREE 14-DAY COURSES OF TREATMENT

Sodium Free

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C, red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-462-0851

Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE.
DO NOT USE IF THE FOL SCAL IMPRINTED WITH THE PREV/ACID LOGO IS CRACKED OR BLACK GRIND IS FOUND ON THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN

Drug Facts (continued)

- theophylline (asthma medicine)
- tacrolimus (immune system medicine)
- atazanavir (medicine for HIV infection)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts

Active ingredient (in each capsule) Lansoprazole 15 mg Acid reducer

Purpose Acid reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)

MADE IN: _____

EXP: _____

LOT: _____

NOVARTIS

Distributed by Novartis Consumer Health, Inc.
Persimmon, NJ 07754-0622 ©2010 XXXX
PREV/ACID is a registered trademark of Novartis Pharmaceuticals North America, Inc. and is used under license.

3 0067-6286-42 6

P0

PREV/ACID[®]
Lansoprazole delayed-release capsules 15 mg / acid reducer

42 CAPSULES
THREE 14-DAY COURSES OF TREATMENT

3 BOTTLES INSIDE



(b) (4)



3 0067-6286-42 6



PREVACID[®] 24HR
Lansoprazole delayed-release capsules 15 mg / acid reducer

Actual Size
ONE 14-DAY COURSE OF TREATMENT

42 CAPSULES
THREE 14-DAY COURSES OF TREATMENT

Sodium Free

SAVE \$X NOW

• May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

• Clinically Proven To Treat Frequent Heartburn

3 BOTTLES INSIDE
42 CAPSULES TOTAL

3 BOTTLES INSIDE
42 CAPSULES TOTAL

Walmart
Save Money Live Better.
1-800-WAL-MART

If this product is found for sale at a location other than Walmart, please call 1-800-WAL-MART

LOT: _____
EXP: _____
MADE IN: _____

NOVARTIS
Distributed by Novartis Consumer Health, Inc.
Persimmon, NJ 07754-0622 ©2010 XXXX
PREVACID[®] is a registered trademark of Novartis Pharmaceuticals North America, Inc. and is used under license.

Drug Facts

Active ingredient (in each capsule) Lansoprazole 15 mg Acid reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)

Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-462-0851
Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE.
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH PREVACID[®] IS MISSING OR IF THE BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN



(b) (4)



(b) (4)





Suggest "peel here" corner, on folded edge of visible panel.

FRONT TOP PLY



BACK TOP PLY



BASE PLY

