


PRESSURE SENSITIVE BASE PLY

DRUG FACTS TEXT (ABBREVIATED)	
	TYPE SIZE/STYLE
DRUG FACTS TITLE	N/A / N/A
HEADINGS	8 pt. / Helvetica Condensed Bold Obl.
SUBHEADS	6.0 pt. / Helvetica Condensed Bold
BODY TEXT	6.0 pt. / Helvetica Condensed
TRACKING: 0	HORIZONTAL SCALE: 85% LEADING: 6.5 pt.



Drug Facts (continued)

Other information

- read the directions, warnings and package insert before use
- keep the carton and package insert. They contain important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted methacrylic acid copolymer, polyethylene glycol, polyacrylate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
1-800-452-0051

Visit us at www.prevacid24hr.com

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE. DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

NDC 0067-6286-14

PREVACID 24HR
 Lansoprazole delayed-release capsules 15 mg / acid reducer

• May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

• Clinically Proven To Treat Frequent Heartburn

14 CAPSULES
 ONE 14-DAY COURSE OF TREATMENT

Sodium Free



Drug Facts

Active ingredient (in each capsule) Lansoprazole 15 mg
Purpose Acid reducer

Uses

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss • nausea or vomiting • stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)

Drug Facts (continued)

- atazanavir (medicine for HIV infection)
- methotrexate (arthritis medication)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

• children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

3 0067-6286-14 3

FPD

NOVARTIS

Produced by: Novartis Consumer Health, Inc.
 Parsippany, NJ 07054-0522 ©2014 14121-14
 PREVACID® is a registered trademark of Takeda
 Pharmacia outside North America, Inc. and is used under license.

LOT:
 EXP:
 MADE IN:

DRUG FACTS TEXT	
ORIG. FACTS TITLE	TYPE SIZE STYLE
HEADINGS	9.0 pt / Helvetica Condensed Bold OHL
SUBHEADS	8.0 pt / Helvetica Condensed Bold OHL
BODY TEXT	6.0 pt / Helvetica Condensed
BULLETS	5.0 pt / Helvetica Condensed
TRAILING 0	HORIZONTAL SCALE: 85% LEADING: 6.5 pt
CHARACTER SPACING	-28
BARCODES, HORIZONTALS	1.5 pt, 0.5 pt



DRUG FACTS TEXT	
	TYPE SIZE/STYLE
DRUG FACTS TITLE	9.0 pt. / Helvetica Condensed Bold Obl.
HEADINGS	8.0 pt. / Helvetica Condensed Bold Obl.
SUBHEADS	6.0 pt. / Helvetica Condensed Bold
BODY TEXT	6.0 pt. / Helvetica Condensed
BULLETS	5.0 pt. / Helvetica Condensed
TRACKING: 0	HORIZONTAL SCALE: 85% LEADING: 6.5 pt.
CHARACTERS/INCH: <39	
BARLINES, HAIRLINES: 1.5 pt., 0.5 pt.	



Drug Facts (continued)

Other information

- Read the directions, warnings and package insert before use.
- Keep the carton and package insert. They contain important information.
- Store at 20-25°C (68-77°F).
- Keep product out of high heat and humidity.
- Protect product from moisture.

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, polyacrylic acid copolymer, polyethylene glycol, polyoxate 40, starch, sucrose, sugar spheres, talc, titanium dioxide

Questions or comments?
 Visit us at www.drugsatfda.com
 1-800-432-0061

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE
 DO NOT USE IF THE BOTTLE CAP HAS BEEN OPENED. CHECK FOR A BLACK BAND AROUND THE CENTER OF EACH CAPSULE. IF MISSING OR BROWN.

Drug Facts (continued)

Other information

- azacitidine medicine for HIV infection
- methotrexate (antimetabolite medication)

Sup use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get dizziness

Directions

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts

Active ingredient (in each capsule)
 Lansoprazole 15 mg

Purpose
 Acid reducer

Uses

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.


Ask a doctor before use if you have:

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with light-headedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or light-headedness
- frequent chest pain
- inquiet wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking:

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or cyclosporine (immunosuppressant medicines)

DRUG FACTS TEXT	
	TYPE SIZE/STYLE
DRUG FACTS TITLE	9.0 pt. / Helvetica Condensed Bold Obl.
HEADINGS	8.0 pt. / Helvetica Condensed Bold Obl.
SUBHEADS	6.0 pt. / Helvetica Condensed Bold
BODY TEXT	6.0 pt. / Helvetica Condensed
BULLETS	5.0 pt. / Helvetica Condensed
TRACKING: 0	HORIZONTAL SCALE: 85% LEADING: 6.5 pt.
CHARACTERS/INCH: <39	
BARLINES, HAIRLINES: 1.5 pt., 0.5 pt.	



6⁷/₁₆

TAMPER-EVIDENT BOTTLE.
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION"
OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

Drug Facts	Purpose
Active ingredient (in each capsule) Lansoprazole 15 mg	Acid reducer
Uses	
<ul style="list-style-type: none"> treats frequent heartburn (occurs 2 or more days a week) not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect 	
Warnings	
Allergy alert: Do not use if you are allergic to lansoprazole	
Do not use	
<ul style="list-style-type: none"> if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. 	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> liver disease had heartburn over 3 months. This may be a sign of a more serious condition. heartburn with lightheadedness, sweating or dizziness chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain unexplained weight loss 	<ul style="list-style-type: none"> frequent wheezing, particularly with heartburn nausea or vomiting stomach pain
Ask a doctor or pharmacist before use if you are taking	
<ul style="list-style-type: none"> warfarin (blood-thinning medicine) digoxin (heart medicine) tacrolimus or mycophenolate mofetil (immune system medicines) atazanavir (medicine for HIV infection) 	<ul style="list-style-type: none"> prescription antifungal or anti-yeast medicines theophylline (asthma medicine) immune system medicines methotrexate (arthritis medication)
Stop use and ask a doctor if	
<ul style="list-style-type: none"> your heartburn continues or worsens you need to take this product for more than 14 days you need to take more than 1 course of treatment every 4 months you get diarrhea 	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	

Drug Facts (continued)

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions, warnings and all information before use
- keep the package. It contains important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture

Inactive ingredients colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, gelatin, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, polysorbate 80, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments? 1-800-452-0051



Visit us at www.prevacid24hr.com

What you should know about PREVACID® 24HR

Please read all information before taking PREVACID® 24HR.
Save for future reference.

How PREVACID® 24HR Treats Your Frequent Heartburn

PREVACID® 24HR stops acid production at the source – the pumps that release acid into the stomach. PREVACID® 24HR is taken once a day (every 24 hours), every day for 14 days.

What You Can Expect When Taking PREVACID® 24HR

Frequent heartburn can occur anytime during the 24-hour period (day or night). Take PREVACID® 24HR in the morning before eating. PREVACID® 24HR is clinically proven to treat frequent heartburn. Although some people get complete relief of symptoms within 24 hours, it may take 1 to 4 days for full effect. Make sure you take PREVACID® 24HR every day for 14 days to treat your frequent heartburn.

Who Should Take PREVACID® 24HR

Adults (18 years and older) with frequent heartburn – when you have heartburn 2 or more days a week.

Who Should NOT Take PREVACID® 24HR

People who have one episode of heartburn a week or less, or who want immediate relief of heartburn.

How to Take PREVACID® 24HR

14-DAY Course of Treatment

- Swallow 1 capsule with a glass of water before eating in the morning.
- Take every day for 14 days.
- Do not take more than 1 capsule a day.
- Swallow whole. Do not crush or chew capsules.
- Do not use for more than 14 days unless directed by your doctor.

When to Take PREVACID® 24HR Again

You may repeat a 14-day course of therapy every 4 months.

When to Talk to Your Doctor

Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.

Warnings and When to Ask Your Doctor

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- atazanavir (medicine for HIV infection)
- methotrexate (arthritis medication)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Clinical studies prove PREVACID® 24HR effectively treats frequent heartburn

In three clinical studies, PREVACID® 24HR was shown to be significantly better than placebo in treating frequent heartburn.

How PREVACID® 24HR is Sold

PREVACID® 24HR is available in 14 capsule, 28 capsule and 42 capsule sizes. These sizes contain one, two and three 14-day courses of treatment, respectively. Do not use for more than 14 days in a row unless directed by your doctor. For the 28 count (two 14-day courses) and the 42 count (three 14-day courses), you may repeat a 14-day course every 4 months.

For Questions or Comments About PREVACID® 24HR
Call 1-800-452-0051 or visit us at www.prevacid24hr.com

LOT:
EXP:
MADE IN:



Distributed by: **Novartis Consumer Health, Inc.**, Parsippany, NJ 07054-0622 ©2014 141121-CB
PREVACID® is a registered trademark of Takeda Pharmaceuticals North America, Inc., and is used under license.

DRUG FACTS (STANDARD) TEXT	
	TYPE SIZE/STYLE
DRUG FACTS TITLE	9.0 pt. / Helvetica Condensed Bold Obl.
HEADINGS	8.0 pt. / Helvetica Condensed Bold Obl.
SUBHEADS	6.0 pt. / Helvetica Condensed Bold
BODY TEXT	6.0 pt. / Helvetica Condensed
BULLETS	5.0 pt. / Helvetica Condensed
TRACKING: 0	HORIZONTAL SCALE: 95% LEADING: 6.5 pt.
CHARACTERS/INCH: <39	
BARLINES, HAIRLINES: 1.5 pt., 0.5 pt.	

8¹⁵/₁₆



NDC 0067-6286-29

BONUS 14 FREE capsules!
 (42 capsules for the price of 28)

PREVAACID[®]

Lansoprazole delayed-release capsules 15 mg / acid reducer

- May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours
- Clinically Proven To Treat Frequent Heartburn

42 28 CAPSULES

THREE 14-DAY COURSES OF TREATMENT

Sodium Free

Drug Facts (continued)

Other information

- Read the directions, warnings and package insert before use
- Keep the carton and package insert. They contain important information.
- Store at 20-25°C (68-77°F)
- Keep product out of high heat and humidity
- Protect product from moisture

Inactive ingredients: colloidal silicon dioxide, D&C red No. 28, FD&C blue No. 1, FD&C green No. 3, FD&C red No. 40, polyvinyl hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, magnesium carbonate, methacrylic acid copolymer, polyethylene glycol, polyacrylate 90, starch, sucrose, sugar sphere, talc, titanium dioxide

Questions or comments?
 Visit us at www.druginfo24hr.com
 1-800-432-0061

KEEP THE CARTON AND PACKAGE INSERT. THEY CONTAIN IMPORTANT INFORMATION.

TAMPER-EVIDENT BOTTLE
 DO NOT USE IF THE BOTTLE SEAL IS IMPAIRED OR IF THERE IS A CRACK OR OTHER DAMAGE. A BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BLOWN.

Drug Facts (continued)

- azacinar (medicines for HIV infection)
- methimazole (antithyroid medication)
- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get dizziness

Sup use and ask a doctor if:

- If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children.
- In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults: 18 years of age and older
- This product is to be used once a day (every 24 hours), every day for 14 days
- It may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts

Active ingredient (in each capsule)
 Lansoprazole 15 mg

Purpose
 Acid Reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking:

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)

MADE IN CHINA

EXP: _____

LOT: _____


FPO

0067-6286-29

3 0067-6286-29 7

Produced by Novartis Consumer Health, Inc.
 Product # 0067-6286-29 ©2016
 PREVAACID is a registered trademark of Novartis
 Pharmaceuticals North America, Inc. and is used under license.

DRUG FACT FORMATTING: STANDARD		
DRUG FACTS TEXT	TYPE SIZE	TYPE STYLE
DRUG FACTS TITLE	9 pt.	Helvetica Cond Bold Obl
DRUG FACTS CONT.	8 pt.	Helvetica Cond Bold Obl
HEADINGS	8 pt.	Helvetica Cond Bold Obl
SUBHEADS	6 pt.	Helvetica Cond Bold
BODY TEXT	6 pt.	Helvetica Cond
TRACKING	0	
CONDENSATION	85%	
LEADING	6.5 pt.	
BARLINES, HAIRLINES	1.5, 0.5 pt.	
CHARACTERS/INCH	<30	
BULLETS	5 pt.	



PREVACID[®] 24HR
Lansoprazole delayed-release capsules 15 mg/acid reducer

• May take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

Please read the entire package insert before taking PREVACID[®] 24HR. Save for future reference.

How PREVACID[®] 24HR Treats Your Frequent Heartburn
PREVACID[®] 24HR stops acid production at the source – the pumps that release acid into the stomach. PREVACID[®] 24HR is taken once a day (every 24 hours), every day for 14 days.

What You Can Expect When Taking PREVACID[®] 24HR
Frequent heartburn can occur anytime during the 24-hour period (day or night). Take PREVACID[®] 24HR in the morning before eating. PREVACID[®] 24HR is clinically proven to treat frequent heartburn. Although some people get complete relief of symptoms within 24 hours, it may take 1 to 4 days for full effect. Make sure you take PREVACID[®] 24HR every day for 14 days to treat your frequent heartburn.

Who Should Take PREVACID[®] 24HR
Adults (18 years and older) with frequent heartburn – when you have heartburn 2 or more days a week.

Who Should NOT Take PREVACID[®] 24HR
People who have one episode of heartburn a week or less, or who want immediate relief of heartburn.

How to Take PREVACID[®] 24HR

14-DAY Course of Treatment

- Swallow 1 capsule with a glass of water before eating in the morning.
- Take every day for 14 days.
- Do not take more than 1 capsule a day.
- Swallow whole. Do not crush or chew capsules.
- Do not use for more than 14 days unless directed by your doctor.


When to Take PREVACID[®] 24HR Again
You may repeat a 14-day course of therapy every 4 months.

When to Talk to Your Doctor
Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.

Warnings and When to Ask Your Doctor
Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.



Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin (blood-thinning medicine)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- theophylline (asthma medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- atazanavir (medicine for HIV infection)
- methotrexate (arthritis medication)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Clinical studies prove PREVACID[®] 24HR effectively treats frequent heartburn
In three clinical studies, PREVACID[®] 24HR was shown to be significantly better than placebo in treating frequent heartburn.

How PREVACID[®] 24HR is Sold
PREVACID[®] 24HR is available in 14 capsule, 28 capsule and 42 capsule sizes. These sizes contain one, two and three 14-day courses of treatment, respectively. Do not use for more than 14 days in a row unless directed by your doctor. For the 28 count (two 14-day courses) and the 42 count (three 14-day courses), you may repeat a 14-day course every 4 months.

For Questions or Comments About PREVACID[®] 24HR
Call 1-800-452-0051 or visit us at www.prevacid24hr.com

Distributed by: Novartis Consumer Health, Inc., Parsippany, NJ 07054-0622
©2014 141121-PI
PREVACID[®] is a registered trademark of Takeda Pharmaceuticals North America, Inc., and is used under license.

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
06/30/2015