



Data Matrix Code:  
XX



Data Matrix Code:  
XX





Data Matrix Code:  
XX

37/64  
Datamatrix Code  
0000000  
6mm

1 1/32

FPQ

LOT/EXP AREA  
NO VARNISH

\$5.00  
Coupon Inside

\$5.00  
Coupon Inside

NDC 0000-0000-00

gsk

## Nicorette

nicotine polacrilex lozenge, 2 mg  
stop smoking aid

### mini Lozenge

2

mg

FOR THOSE WHO SMOKE  
THEIR FIRST CIGARETTE  
MORE THAN 30 MINUTES  
AFTER WAKING UP.

If you smoke your first cigarette  
WITHIN 30 MINUTES of waking up,  
use Nicorette 4 mg Lozenge.

20 LOZENGES

2 mg Each

**Drug Facts**

Active ingredient (in each lozenge)	Purpose
Nicotine polacrilex, 2 mg	Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Flip open for Directions and additional information  
Retain this package for complete product information

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

**TAMPER EVIDENT FEATURE:** Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken. Retain outer carton for full product uses, directions and warnings.

6

**Drug Facts (continued)**

**Directions**

- if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge.
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
- it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- store at 20 - 25°C (68 - 77°F)
- keep vial tightly closed and protect from light

**Inactive ingredients** acesulfame potassium, calcium polycarbophil, flavors, magnesium stearate, mannitol, potassium bicarbonate, sodium alginate, sodium carbonate, sucralose, xanthan gum

**Questions or comments?** call toll-free 1-888-569-1743

Distributed by:  
**GSK Consumer Healthcare**  
Warren, NJ 07059

Trademarks are owned by or licensed to the GSK group of companies.  
©2020 GSK group of companies or its licensor.

NPA001552

FPQ

\$5.00  
Coupon Inside

## Nicorette

nicotine polacrilex lozenge, 2 mg  
stop smoking aid

### mini Lozenge

2

mg

FOR THOSE WHO SMOKE  
THEIR FIRST CIGARETTE  
MORE THAN 30 MINUTES  
AFTER WAKING UP.

If you smoke your first cigarette  
WITHIN 30 MINUTES of waking up,  
use Nicorette 4 mg Lozenge.

20 LOZENGES

2 mg Each

**Drug Facts (continued)**

**Directions**

- if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge.
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
- it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- store at 20 - 25°C (68 - 77°F)
- keep vial tightly closed and protect from light

**Inactive ingredients** acesulfame potassium, calcium polycarbophil, flavors, magnesium stearate, mannitol, potassium bicarbonate, sodium alginate, sodium carbonate, sucralose, xanthan gum

**Questions or comments?** call toll-free 1-888-569-1743

Distributed by:  
**GSK Consumer Healthcare**  
Warren, NJ 07059

Trademarks are owned by or licensed to the GSK group of companies.  
©2020 GSK group of companies or its licensor.

NPA001552

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Use at least 9 Nicorette mini Lozenges per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette mini Lozenges for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

**Opening Directions:**  
Push in child resistant tab on the vial with thumb. Flip up the top of vial. Turn upside down and shake to remove lozenge.

**For more information and for a FREE individualized stop smoking program, please visit [www.nicorette.com](http://www.nicorette.com) or see inside for more details.**

SAVE \$5.00

on any one (1) Nicorette Mini 81ct

Redemption of this coupon requires purchase of Nicorette Mini 81ct at participating retailer. Coupon is valid only for one (1) Nicorette Mini 81ct purchase. Limit 1 coupon per purchase of Nicorette Mini 81ct. No more than four (4) identical coupons can be used for the same product in same transaction. Coupons are not authorized to be used when the consumer is purchasing products for resale. You pay any sales tax. RETAILER: Coupons may not be reproduced, photocopied, altered or altered in any way. GSK will reimburse you the face value of this coupon plus 12¢ Handling Fee if submitted in compliance with our Redemption Policy. Copy available upon request. Cash value 1/100 of 1¢. Coupon is void if altered, transferred, exchanged, sold or auctioned, applied or if prohibited by law. Good only in USA and at APOs. FPOs. Send coupons to: GSK 1014, NCH Marketing Services, P.O. Box 880001, El Paso, TX 88888-0001, ©2020 GSK.

030766-078095

FPQ

2 5/8      2 5/8

5 1/4

1 1/4

4 11/32

**Opening Directions:**  
Push in child resistant tab on the vial with thumb. Flip up the top of vial. Turn upside down and shake to remove lozenge.



LOT/EXP AREA  
NO VARNISH

**Drug Facts**  
*(continued)*

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge.
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- do not use more than 5 lozenges in 6 hours.
- Do not use more than 20 lozenges per day.
- It is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- store at 20 - 25°C (68 - 77°F)
- keep vial tightly closed and protect from light

**Inactive ingredients**  
acesulfame potassium, calcium polycarbophil, flavors, magnesium stearate, mannitol, potassium bicarbonate, sodium alginate, sodium carbonate, sucralose, xanthan gum

**Questions or comments?**  
call toll-free 1-888-569-1743

Trademarks are owned by or licensed to the GSK group of companies.

Distributed by:  
**GSK Consumer Healthcare**  
Warren, NJ 07059  
©2020 GSK group of companies or its licensor. NPA001554

NDC 0000-0000-00

gsk

# Nicorette

nicotine polacrilex  
lozenge, 2 mg  
stop smoking aid

## mini Lozenge

Mint

2  
mg

FOR THOSE WHO  
SMOKE THEIR FIRST  
CIGARETTE MORE  
THAN 30 MINUTES  
AFTER WAKING UP.

If you smoke your first  
cigarette **WITHIN 30  
MINUTES** of waking up,  
use Nicorette 4 mg Lozenge.

81 LOZENGES, 2 mg Each  
(3 Vials of 27)

For more information and for a FREE  
individualized stop smoking program, please visit  
[www.Nicorette.com](http://www.Nicorette.com) or see inside for more details.

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Use at least 9 Nicorette mini Lozenges per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette mini Lozenges for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

■ not for sale to those under 18 years of age

■ proof of age required

■ not for sale in vending machines or from any source where proof of age cannot be verified

**TAMPER EVIDENT FEATURE:**  
Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken. Retain outer carton for full product uses, directions and warnings.

**Drug Facts**

Active ingredient (in each lozenge)	Purpose
Nicotine polacrilex, 2 mg	Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- If you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

7 21/32



0 0000-0000-00 0

12 7/64



0000000000

EV CODE  
0000000000



Data Matrix Code:  
XX

37/64  
Datamatrix Code  
0000000  
6mm

1 1/32

**FPO**

LOT/EXP AREA  
NO VARNISH

**\$5.00**  
Coupon Inside

**\$5.00**  
Coupon Inside

NDC 0000-0000-00 **gsk**

# Nicorette

nicotine polacriflex lozenge, 4 mg  
stop smoking aid

## mini Lozenge

**4**  
mg

FOR THOSE WHO SMOKE  
THEIR FIRST CIGARETTE  
WITHIN 30 MINUTES OF  
WAKING UP.

If you smoke your first cigarette  
MORE THAN 30 MINUTES after  
waking up, use Nicorette 2 mg Lozenge.

20 LOZENGES      4 mg Each

**Drug Facts**

Active ingredient (in each lozenge)	Purpose
Nicotine polacriflex, 4 mg	Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

*Flip open for Directions and additional information. Retain this package for complete product information.*

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

**TAMPER EVIDENT FEATURE:** Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken. Retain outer carton for full product uses, directions and warnings.

FPO

6

**Drug Facts (continued)**

**Directions**

- if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge.
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
- it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- store at 20 - 25°C (68 - 77°F)
- keep vial tightly closed and protect from light

**Inactive ingredients** acesulfame potassium, calcium polycarbophil, flavors, magnesium stearate, mannitol, potassium bicarbonate, sodium alginate, sodium carbonate, sucralose, xanthan gum

**Questions or comments?** call toll-free 1-888-569-1743

Distributed by:  
**GSK Consumer Healthcare**  
Warren, NJ 07059

Trademarks are owned by or licensed to the GSK group of companies.  
©2020 GSK group of companies or its licensor.

NPA001553

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Use at least 9 Nicorette mini Lozenges per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette mini Lozenges for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

**Opening Directions:**  
Push in child resistant tab on the vial with thumb. Flip up the top of vial. Turn upside down and shake to remove lozenge.

**For more information and for a FREE individualized stop smoking program, please visit [www.nicorette.com](http://www.nicorette.com) or see inside for more details.**

MANUFACTURER'S COUPON | EXPIRES 12/31/2020

SAVE \$5.00

on any one (1) Nicorette Mini 81ct

Redeemable at participating drug and mass merchandise stores. CONSUMER: Coupons are redeemable only by a consumer purchasing the GSK product(s) indicated and as specified, purchasing the quantity and/or product size specified. Limit 1 coupon per purchase of indicated product(s). No more than four (4) identical coupons can be used for the same product in same transaction. Coupons are not authorized to be used when the consumer is purchasing products for resale. You pay any sales tax. RETAILER: Coupons may not be reproduced, photocopied, altered or altered in any way. GSK will reimburse you the face value of this coupon plus 12¢ Handling Fee if submitted in compliance with our Redemption Policy. Copy available upon request. Cash value 1/100 of 1¢. Coupon is void if altered, transferred, exchanged, sold or auctioned, applied or if prohibited by law. Good only in USA and at APOs. FPOs. Send coupons to: GSK 1014, NCH Marketing Services, P.O. Box 880001, El Paso, TX 88888-0001, ©2020 GSK.

030766-078095

FPO

2 5/8      2 5/8

5 1/4

1 1/4
4 11/32
7 21/32

12 7/64
EV CODE  
XXXXXXXXXX

**Opening Directions:**  
Push in child resistant tab on the vial with thumb. Flip up the top of vial. Turn upside down and shake to remove lozenge.



LOT/EXP AREA  
NO VARNISH

NDC 0000-0000-00 gsk

# Nicorette

**nicotine polacrilex  
lozenge, 4 mg  
stop smoking aid**

## mini Lozenge

**Mint**

4  
mg

**FOR THOSE WHO  
SMOKE THEIR FIRST  
CIGARETTE WITHIN  
30 MINUTES OF  
WAKING UP.**

If you smoke your first  
cigarette **MORE THAN 30  
MINUTES** after waking up,  
use Nicorette 2 mg Lozenge.

**81 LOZENGES, 4 mg Each  
(3 Vials of 27)**

For more information and for a FREE individualized stop smoking program, please visit [www.Nicorette.com](http://www.Nicorette.com) or see inside for more details.

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** — Use at least 9 Nicorette mini Lozenges per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette mini Lozenges for the full 12 weeks.
4. Use with a support program as directed in the enclosed User's Guide.

■ not for sale to those under 18 years of age

■ proof of age required

■ not for sale in vending machines or from any source where proof of age cannot be verified

**TAMPER EVIDENT FEATURE:**  
Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken. Retain outer carton for full product uses, directions and warnings.

**Drug Facts**  
*(continued)*

**nicotine lozenge is a medicine and must be used a certain way to get the best results**

- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge.
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- do not use more than 5 lozenges in 6 hours.
- Do not use more than 20 lozenges per day.
- It is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- store at 20 - 25°C (68 - 77°F)
- keep vial tightly closed and protect from light

**Inactive ingredients**  
acesulfame potassium, calcium polycarbophil, flavors, magnesium stearate, mannitol, potassium bicarbonate, sodium alginate, sodium carbonate, sucralose, xanthan gum

**Questions or comments?**  
call toll-free 1-888-569-1743

Trademarks are owned by or licensed to the GSK group of companies.

Distributed by:  
**GSK Consumer Healthcare**  
Warren, NJ 07059  
©2020 GSK group of companies or its licensor. NPA001555

**Drug Facts**

Active ingredient (in each lozenge)	Purpose
Nicotine polacrilex, 4 mg	Stop smoking aid

**Use** - reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**

**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours



0 0000-0000-00 0



0000000000

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

NUSHIN F TODD  
03/05/2021 01:36:34 PM