

214534

**REGULATORY LEGEND**

	<b>Actual Size</b>
DRUG FACTS TITLE	N/A
DRUG FACTS CONTINUED	N/A
HEADINGS	N/A
SUBHEADINGS/BODY TEXT	N/A
LEADING	N/A
BULLETS	N/A
# OF CHARACTERS PER INCH	N/A
SPACE BEFORE BULLET	N/A
BARLINES, HAIRLINES	N/A
SPACE BETWEEN HAIRLINES AND BOX END	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	7 pt

**\$5.00**  
Coupon Inside

**Drug Facts**

**Active ingredient (in each lozenge)** Nicotine polacrilex, 2 mg. **Purpose** Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Flip open for Directions and additional information**  
Retain this package for complete product information

not for sale to those under 18 years of age  
proof of age required  
not for sale in vending machines or from any source where proof of age cannot be verified

**TAMPER EVIDENT FEATURE:** Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken. Retain outer carton for full product uses, directions and warnings.

3 07663 07577 3

NEW CHERRY PEPPERMINT FLAVOR

NDC 0135-8070-20 HALEON

**\$5.00**  
Coupon Inside

**NICORETTE**  
nicotine polacrilex lozenge, 2 mg  
stop smoking aid  
mini  
Lozenge

**2 mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Lozenge.

**20 LOZENGES**  
2 mg Each

ACTUAL SIZE

CHERRY PEPPERMINT

**Drug Facts (continued)**

**Directions**

- If you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge
- If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge.
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- To improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- Do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
- It is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

**Other information** • store between 20°-25°C (68°-77°F)  
• keep the vial tightly closed and protect from light

**Inactive ingredients**  
acesulfame potassium, anhydrous dibasic calcium phosphate, carmine (color), croscopolidone, flavors, hypromellose, magnesium stearate, mannitol, microcrystalline cellulose, sodium bicarbonate, sodium carbonate anhydrous, sucralose, xanthan gum, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Switzerland

Trademarks are owned by or licensed to the Haleon group of companies.  
©2024 Haleon group of companies or its licensor.  
Patent Pending  
214535

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Use at least 9 Nicorette mini Lozenges per day during the first six weeks.
- Use Long Enough** — Use Nicorette mini Lozenges for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.

**Opening Directions:**  
Squeeze child resistant tabs on each side. Flip up the top of vial, turn upside down and shake to remove lozenge.

For more information and for a FREE individualized stop smoking program, please visit [www.Nicorette.com](http://www.Nicorette.com) or see inside for more details.

MANUFACTURER'S COUPON | EXPIRES 12/31/2026

**SAVE \$5.00**

on any one (1) Nicorette (nicotine polacrilex) lozenge 2 mg or 4 mg, mini 81 ct

0307661035429

Redeemable at food, drug and mass merchandise stores.  
CONSUMER: Limit 1 coupon per purchase of indicated product(s). You pay any sales tax.  
RETAILER: Haleon Group of Companies will reimburse you the face value of this coupon plus 12¢ Handling Fee if submitted in compliance with our Redemption Policy. Copy available upon request. Cash value 1/100 of 1¢. Coupon is void if altered, transferred, exchanged, sold or auctioned, copied or if prohibited by law. Good only in USA and at APOs, FPOs. Send coupons to: Haleon Group of Companies, P.O. Box 880001, El Paso, TX 88588-0001.  
©2024 Haleon Group of Companies. Read and follow label directions.

214535

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	12 pt
DRUG FACTS CONTINUED	8 pt
HEADINGS	8 pt
SUBHEADINGS/BODY TEXT	6 pt, 6 pt
LEADING	6.5 pt
BULLETS	5 pt
# OF CHARACTERS PER INCH	29
SPACE BEFORE BULLET	2 ems
BARLINES, HAIRLINES	1.5 pt, 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	18.5 pt

9 5 5 7 1 2

**Opening Directions:**  
Squeeze child resistant tabs on each side. Flip up the top of vial. Turn upside down and shake to remove lozenge.



NEW CHERRY PEPPERMINT FLAVOR    NDC 005-9076-01    HALEON

# NICORETTE

nicotine polacrilex lozenge, 2 mg  
stop smoking aid  
mini  
Lozenge

2 mg

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicorette 4 mg Lozenge.



ACTUAL SIZE

81 LOZENGES,  
2 mg EACH (3 Vials of 27)

CHERRY PEPPERMINT

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Use at least 9 Nicorette mini Lozenges per day during the first six weeks.
- Use Long Enough** — Use Nicorette mini Lozenges for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.

3 07663 07584 1

**Drug Facts (continued)**

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge.
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
- it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- store between 20°-25°C (68°-77°F)
- keep the vial tightly closed and protect from light

**Inactive ingredients**  
ascorbic acid, calcium phosphate, croscopolvidone, flavors, hypromellose, magnesium stearate, mannitol, microcrystalline cellulose, sodium bicarbonate, sodium carbonate anhydrous, sucralose, xanthan gum, xylitol

**Questions or comments?**  
call toll-free 1-888-569-1743

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Switzerland  
Trademarks are owned by or licensed to the Haleon group of companies.  
©2024 Haleon group of companies or its licensor.  
Patent Pending  
214536

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Use at least 9 Nicorette mini Lozenges per day during the first six weeks.
- Use Long Enough** — Use Nicorette mini Lozenges for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.

**not for sale to those under 18 years of age**

**proof of age required**

**not for sale in vending machines or from any source where proof of age cannot be verified**

**TAMPER EVIDENT FEATURE:**  
Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken. Retain outer carton for full product uses, directions and warnings.

For more information and for a FREE individualized stop smoking program, please visit [www.Nicorette.com](http://www.Nicorette.com) or see inside for more details.

**Drug Facts**

Active ingredient (in each lozenge)	Purpose
Nicotine polacrilex, 2 mg	Stop smoking aid

**Use**  
reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**


- if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

214536

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	10 pt
DRUG FACTS CONTINUED	8 pt
HEADINGS	8 pt
SUBHEADINGS/BODY TEXT	6 pt, 6 pt
LEADING	6.5 pt
BULLETS	5 pt
# OF CHARACTERS PER INCH	29
SPACE BEFORE BULLET	N/A
BARLINES, HAIRLINES	1.5 pt, 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	17.5 pt



214537

**REGULATORY LEGEND**

	<b>Actual Size</b>
DRUG FACTS TITLE	N/A
DRUG FACTS CONTINUED	N/A
HEADINGS	N/A
SUBHEADINGS/BODY TEXT	N/A
LEADING	N/A
BULLETS	N/A
# OF CHARACTERS PER INCH	N/A
SPACE BEFORE BULLET	N/A
BARLINES, HAIRLINES	N/A
SPACE BETWEEN HAIRLINES AND BOX END	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	7 pt

**\$5.00**  
Coupon Inside

**Drug Facts**

**Active ingredient (in each lozenge)** Nicotine polacrilex, 4 mg. **Purpose** Stop smoking aid

**Use** • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Flip open for Directions and additional information  
Retain this package for complete product information

not for sale to those under 18 years of age  
proof of age required  
not for sale in vending machines or from any source where proof of age cannot be verified

**TAMPER EVIDENT FEATURE:** Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken. Retain outer carton for full product uses, directions and warnings.

3 07663 07574 2

NEW CHERRY PEPPERMINT FLAVOR

NDC 0135-0071-20 HALEON

**\$5.00**  
Coupon Inside

**NICORETTE**  
nicotine polacrilex lozenge, 4 mg  
stop smoking aid  
mini  
Lozenge

**4mg**

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicorette 2 mg Lozenge.

20 LOZENGES  
4 mg Each

CHERRY PEPPERMINT

ACTUAL SIZE

**Drug Facts (continued)**

**Directions**

- if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 6 hours

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge.
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
- it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

**Other information** • store between 20°-25°C (68°-77°F)  
• keep the vial tightly closed and protect from light

**Inactive ingredients**  
acesulfame potassium, anhydrous dibasic calcium phosphate, carmine (color), crospovidone, flavors, hypromellose, magnesium stearate, mannitol, microcrystalline cellulose, sodium bicarbonate, sodium carbonate anhydrous, sucralose, xanthan gum, xylitol

**Questions or comments?** call toll-free 1-888-569-1743

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Switzerland

Trademarks are owned by or licensed to the Haleon group of companies.  
©2024 Haleon group of companies or its licensor.  
Patent Pending  
214538

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough — Use at least 9 Nicorette mini Lozenges per day during the first six weeks.
- Use Long Enough — Use Nicorette mini Lozenges for the full 12 weeks.
- Use with a support program as directed in the enclosed User's Guide.

**Opening Directions:**  
Squeeze child resistant tabs on each side. Flip up the top of vial. Turn upside down and shake to remove lozenge.

For more information and for a FREE individualized stop smoking program, please visit [www.Nicorette.com](http://www.Nicorette.com) or see inside for more details.

MANUFACTURER'S COUPON | EXPIRES 12/31/2026

**SAVE \$5.00**

on any one (1) Nicorette (nicotine polacrilex) lozenge 2 mg or 4 mg, mini 81 ct

Redeemable at food, drug and mass merchandise stores.  
CONSUMER: Limit 1 coupon per purchase of indicated product(s). You pay any sales tax.  
RETAILER: Haleon Group of Companies will reimburse you the face value of this coupon plus 12¢ Handling Fee if submitted in compliance with our Redemption Policy. Copy available upon request. Cash value 1/100 of 1¢. Coupon is void if altered, transferred, exchanged, sold or auctioned, copied or if prohibited by law. Good only in USA and at APOs, FPOs. Send coupons to: Haleon Group of Companies, P.O. Box 880001, El Paso, TX 88589-0001.  
©2024 Haleon Group of Companies. Read and follow label directions.

030766-035429

214538

**REGULATORY LEGEND**

	Actual Size
DRUG FACTS TITLE	12 pt
DRUG FACTS CONTINUED	8 pt
HEADINGS	8 pt
SUBHEADINGS/BODY TEXT	6 pt, 6 pt
LEADING	6.5 pt
BULLETS	5 pt
# OF CHARACTERS PER INCH	29
SPACE BEFORE BULLET	2 ems
BARLINES, HAIRLINES	1.5 pt, 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	18.5 pt

655712

**Opening Directions:**  
Squeeze child resistant tabs on each side. Flip up the top of vial. Turn upside down and shake to remove lozenge.



NEW CHERRY PEPPERMINT FLAVOR    NDC 035-9074-01    HALEON

# NICORETTE

nicotine polacrilex lozenge, 4 mg  
stop smoking aid  
mini  
Lozenge

4mg

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Nicorette 2 mg Lozenge.



**81 LOZENGES, 4 mg EACH (3 Vials of 27)**    CHERRY PEPPERMINT

*For more information and for a FREE individualized stop smoking program, please visit [www.Nicorette.com](http://www.Nicorette.com) or see inside for more details.*

**TO INCREASE YOUR SUCCESS IN QUITTING:**

- You must be motivated to quit.
- Use Enough** — Use at least 9 Nicorette mini Lozenges per day during the first six weeks.
- Use Long Enough** — Use Nicorette mini Lozenges for the full 12 weeks.
- Use with a support program** as directed in the enclosed User's Guide.



3 07663 07581 0

**Drug Facts (continued)**

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge.
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
- it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

**Other information**

- store between 20°-25°C (68°-77°F)
- keep the vial tightly closed and protect from light

**Inactive ingredients**  
asculfame potassium, anhydrous dibasic calcium phosphate, carmine (color), croscopolidone, flavors, hydroxypropylcellulose, magnesium stearate, mannitol, microcrystalline cellulose, sodium bicarbonate, sodium carbonate anhydrous, sucralose, xanthan gum, xylitol

**Questions or comments?**  
call toll-free 1-888-569-1743

Distributed by:  
Haleon  
Warren, NJ 07059  
Made in Switzerland  
Trademarks are owned by or licensed to the Haleon group of companies.  
©2024 Haleon group of companies or its licensor.  
Patent Pending  
214539

**Drug Facts**

Active ingredient (in each lozenge)	Purpose
Nicotine polacrilex, 4 mg	Stop smoking aid

**Use**  
reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

214539

REGULATORY LEGEND	
	Actual Size
DRUG FACTS TITLE	10 pt
DRUG FACTS CONTINUED	8 pt
HEADINGS	8 pt
SUBHEADINGS/BODY TEXT	6 pt, 6 pt
LEADING	6.5 pt
BULLETS	5 pt
# OF CHARACTERS PER INCH	29
SPACE BEFORE BULLET	N/A
BARLINES, HAIRLINES	1.5 pt, 0.5 pt
SPACE BETWEEN HAIRLINES AND BOX END	2 spaces
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	N/A
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	17.5 pt