


Expiration Date

Lot #



Drug Facts (continued)

Directions

adults and children 6 years and over	one 10 mg capsule once daily; do not take more than one 10 mg capsule in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
adults 65 years and over	ask a doctor
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- protect from light
- do not use if Tamper-Evident seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing

Inactive ingredients FD&C Blue #1, FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, mannitol, pharmaceutical ink, polyethylene glycol 400, purified water, sodium hydroxide, sorbitan, sorbitol

Questions or comments? call 1-800-447-1140

* This product is not manufactured or distributed by McNeil Consumer Healthcare Division of McNeil-PPC, Inc. owner of the registered trademark of Zyrtec®
 Manufactured by: Banner Pharmacaps Inc.
 4125 Premier Drive
 High Point, NC 27265

NDC 010888-7127-12

Compare to the Active Ingredient of Zyrtec®*

Cetirizine HCl Capsules, 10 mg

Antihistamine

ALLERGY

Indoor & Outdoor Allergies

24 hour Relief of ■ Sneezing ■ Running Nose
 ■ Itchy, Watery eyes ■ Itchy Throat or Nose

200 Softgels**
 Liquid-filled Capsules**

NDC 010888-7127-12

Compare to the Active Ingredient of Zyrtec®*

Cetirizine HCl Capsules, 10 mg

Antihistamine

ALLERGY

Indoor & Outdoor Allergies

24 hour Relief of ■ Sneezing ■ Running Nose
 ■ Itchy, Watery eyes ■ Itchy Throat or Nose

200 Softgels**
 Liquid-filled Capsules**

Drug Facts

Active ingredient (in each capsule)	Purpose
Cetirizine HCl 10 mg	Antihistamine

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
 Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
 Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.


DIE 0163C

11.625 x 12.5 flat
 2.75 x 2.75 x 5.5

Labeling Format Information:

Fonts: Helvetica Condensed, Bold, Bold Italic	
Drug Facts: 10 pt	Leading: 6.5 pt
Header: 8 pt	Bullets: 6.5 pt
Subheader: 6.2 pt	Barlines: 2.5 pt
Body Text: 6.2 pt	Hairlines: .75 pt
Drug Facts (continued): 8 pt	
Horizontal Scale: 100% Average Kerning: 0	

(b) (4)



FOLD PROOF
ALONG
DOTTED LINE
to simulate an
actual printed
label layout



(b) (4)

Labeling Format Information:			
Fonts: Helvetica Condensed, Bold, Bold Italic			
Drug Facts:	NA pt	Leading:	5.4 pt
Header:	5.4 pt	Bullets:	3.6 pt
Subheader:	5.4 pt	Barlines:	NA pt
Body Text:	5.4 pt	Hairlines:	NA pt
Drug Facts (continued):	NA pt		
Horizontal Scale: 100%		Average Kerning: 0	

NDC 010888-7127-12 Compare to the Active Ingredient of Zyrtec®*

Cetirizine HCl Capsules, 10 mg
Antihistamine
Allergy

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

200 Liquid-filled capsules **

Active ingredient (in each capsule)
Cetirizine HCl 10 mg Antihistamine

Purpose
Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:
■ if breast-feeding: not recommended
■ if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions Adults and children 6 years and over: one 10 mg capsule once daily, do not take more than one 10 mg capsule in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
Adults 65 years and over: ask a doctor. Children under 6 years of age: ask a doctor. Consumers with liver or kidney disease: ask a doctor.

Other information ■ store at 20-25°C (68-77°F) ■ avoid high humidity and excessive heat above 40°C (104°F) ■ protect from light ■ do not use if Tamper-Evident seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing

Inactive ingredients FD&C Blue #1, FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, mannitol, pharmaceutical ink, polyethylene glycol 400, purified water, sodium hydroxide, sorbitan, sorbitol

* This product is not manufactured or distributed by McNeil Consumer Healthcare Division of McNeil-PPC, Inc. owner of the registered trademark of Zyrtec®

Manufactured by:
Banner Pharmaceuticals Inc.
4125 Premier Drive
High Point, NC 27265

Lot #
Expiration Date

(b) (4)

Labeling Format Information:

Fonts: Helvetica Condensed, Bold, Bold Italic

Drug Facts:	NA pt	Leading:	6.5 pt
Header:	8 pt	Bullets:	5 pt
Subheader:	6 pt	Barlines:	NA pt
Body Text:	6 pt	Hairlines:	NA pt
Drug Facts (continued):	NA pt		

Horizontal Scale: 88%

Average Kerning: 0

DIE 0625C
 5.432 x 6.625 Flat
 1.5 x 1.5 x 2.655

Drug Facts (continued)

Directions	
adults and children 6 years and over	one 10 mg capsule once daily; do not take more than one 10 mg capsule in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
adults 65 years and over	ask a doctor
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information ■ store at 20-25°C (68-77°F) ■ avoid high humidity and excessive heat above 40°C (104°F) ■ protect from light
 ■ do not use if Tamper-Evident seal under bottle cap imprinted with "Sealed for Your Protection" is Broken or missing

NDC 010888-7127-11 Compare to the Active Ingredient of Zyrtec®*

Cetirizine HCl Capsules, 10 mg
Antihistamine
ALLERGY
Indoor & Outdoor Allergies

24 hour Relief of ■ Sneezing ■ Running Nose
 ■ Itchy, Watery eyes ■ Itchy Throat or Nose

10 Softgels**
 Liquid-filled capsules**

Drug Facts

Active ingredient (in each capsule)	Purpose
Cetirizine HCl 10 mg	Antihistamine

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes
 ■ itching of the nose or throat

Warnings
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
 Ask a doctor before use if you have liver or kidney disease.
 Your doctor should determine if you need a different dose.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product ■ drowsiness may occur ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if an allergic reaction to this product occurs.
 Seek medical help right away.

If pregnant or breast-feeding: ■ if breast-feeding: not recommended
 ■ if pregnant: ask a health professional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Inactive ingredients FD&C Blue #1, FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, mannitol, pharmaceutical ink, polyethylene glycol 400, purified water, sodium hydroxide, sorbitan, sorbitol


Questions or comments?
 call 1-800-447-1140

*This product is not manufactured or distributed by McNeil Consumer Healthcare Division of McNeil-PPC, Inc., owner of the registered trademark of Zyrtec®.


Manufactured by:
 Banner Pharmaceuticals Inc.
 4125 Premier Drive
 High Point, NC 27265

Expiration Date

Lot #



000000-000000



#dd
pp
12of5

(b) (4)

Labeling Format Information:

Fonts: Helvetica Condensed, Bold, Bold Italic

Drug Facts:	10 pt	Leading:	6.5 pt
Header:	8 pt	Bullets:	5 pt
Subheader:	6 pt	Barlines:	2.5 pt
Body Text:	6 pt	Hairlines:	.5 pt
Drug Facts (continued):	8 pt		

Horizontal Scale: 93%

Average Kerning: 0

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER
02/06/2013