


BACK

FOR BACK ONLY



090177e1861a8eb9\Final\Final On: 10-Feb-2015 14:32

(b) (4)



FRONT

090177e1861a8eb5\Final\Final On: 09-Feb-2015 17:35

(b) (4)


090177e1861a848fFinalOn: 09-Feb-2015 17:30



BACK

FRONT

ACTUAL SIZE



090177e185c4fcfb\Final\Final On: 02-Oct-2014 18:37

(b) (4)

3/16" x 3/16" DATA MATRIX CODE (ECC200) **012221**,
 PRINTED IN BLACK OR DARK BLUE ON
 UNPRINTED WHITE BACKGROUND

LIFT HERE
For More Drug Facts

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

<p>Drug Facts</p> <p>Active ingredients (in each tablet)</p> <p>Ibuprofen 200 mg (NSAID) * ..Pain reliever/fever reducer Phenylephrine HCl 10 mg Nasal decongestant *nonsteroidal anti-inflammatory drug</p> <p>Purposes</p> <p>temporarily relieves these symptoms associated with the common cold or flu:</p> <ul style="list-style-type: none"> ■ headache ■ fever ■ sinus pressure ■ nasal congestion ■ minor body aches and pains ■ reduces swelling of the nasal passages ■ temporarily restores freer breathing through the nose 	<p>Drug Facts (continued)</p> <p>Warnings</p> <p>Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> ■ are age 60 or older
---	---

PAA012221

PANEL 1

<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed <p>Do not use</p> <ul style="list-style-type: none"> ■ in children under 12 years of age because this product contains too much medication for children under this age ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> ■ stomach bleeding warning applies to you 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes or have trouble urinating due to an enlarged prostate gland ■ you are taking a diuretic <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ under a doctor's care for any serious condition ■ taking any other product that contains phenylephrine or any other nasal decongestant ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug <p>When using this product</p> <ul style="list-style-type: none"> ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you experience any of the following signs of stomach bleeding: ■ feel faint
--	---

PANEL 2

<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ nasal congestion lasts for more than 7 days ■ symptoms continue or get worse ■ redness or swelling is present in the painful area ■ you get nervous, dizzy, or sleepless ■ any new symptoms appear <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed ■ adults and children 12 years of age and over: <ul style="list-style-type: none"> ■ take 1 tablet every 4 hours while symptoms persist. ■ do not use more than 6 tablets in any 24-hour period unless directed by a doctor 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ children under 12 years of age: do not use because this product contains too much medication for children under this age <p>Other information</p> <ul style="list-style-type: none"> ■ store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). ■ read all warnings and directions before use. Keep carton. <p>Inactive ingredients</p> <p>acesulfame potassium, artificial flavor, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, sodium lauryl sulfate, stearic acid, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum</p> <p>Questions or comments?</p> <p>Call weekdays from 9 AM to 5 PM EST toll free at 1-800-88-ADVIL</p>
---	---

BASE PANEL

090177e1857b6ddc\Final\Final On: 01-Jul-2014 04:52



RELEASED

Name _____ Date _____

DEPARTMENT	REVIEWED BY		DATE	✓ with Changes
	PRINT NAME	SIGNATURE		
Graphic Artist				<input type="checkbox"/>
Labeling Compliance				<input type="checkbox"/>
Medical				<input type="checkbox"/>
Regulatory Affairs				<input type="checkbox"/>
Marketing				<input type="checkbox"/>
Legal				<input type="checkbox"/>
Trademark				<input type="checkbox"/>
Display Services				<input type="checkbox"/>
Promotions				<input type="checkbox"/>
Graphic Artist Closed				<input type="checkbox"/>