

NDA 22572/S-009

Page 3

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use MITOSOL® safely and effectively. See full prescribing information for MITOSOL®.

MITOSOL® (mitomycin for solution) for ophthalmic use
Initial U.S. Approval: 1974

INDICATIONS AND USAGE

Mitosol® is an antimetabolite indicated as an adjunct to ab externo glaucoma surgery. (1)

DOSAGE AND ADMINISTRATION

Mitosol® is intended for topical application to the surgical site of glaucoma filtration surgery. It is not intended for intraocular administration. (2)

- Each vial of Mitosol® contains 0.2 mg of mitomycin and mannitol in a 1:2 concentration ratio. To reconstitute, add 1 mL of Sterile Water for Injection, then shake to dissolve. If product does not dissolve immediately, allow to stand at room temperature until the product has dissolved into solution. (2.1)
- Fully saturate sponges provided within the Mitosol® Kit utilizing the entire reconstituted contents of the vial in the manner prescribed in the Instructions for Use. (2.2)
- Apply fully saturated sponges equally to the treatment area, in a single layer, with the use of a surgical forceps. Keep the sponges on the treatment area for two (2) minutes, then remove and return to the Mitosol® Tray for defined disposal. (2.2)

DOSAGE FORMS AND STRENGTHS

Each vial contains a sterile lyophilized mixture of 0.2 mg mitomycin and 0.4 mg mannitol; when reconstituted with Sterile Water for Injection, the solution contains 0.2 mg/mL mitomycin. (3)

CONTRAINDICATIONS

- Hypersensitivity to mitomycin. (4.1)

WARNINGS AND PRECAUTIONS

- **Cell Death:** Mitomycin is cytotoxic. Use of mitomycin in concentrations higher than 0.2 mg/mL or use for longer than 2 minutes may lead to unintended corneal and/or scleral damage including thinning or perforation. Direct contact with the corneal endothelium will result in cell death. (5.1)
- **Hypotony:** The use of mitomycin has been associated with an increased incidence of post-operative hypotony. (5.2)
- **Cataract Development:** Use in phakic patients has been correlated to a higher incidence of lenticular change and cataract formation. (5.3)
- **Embryo-Fetal Toxicity:** Can cause fetal harm. Advise of potential risk to a fetus. Verify pregnancy status in females of reproductive potential prior to use. (5.4, 8.1, 8.3)

ADVERSE REACTIONS

The most frequent adverse reactions to Mitosol® occur locally and include hypotony, hypotony maculopathy, blebitis, endophthalmitis, vascular reactions, corneal reactions, and cataract. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Mobius Therapeutics LLC 1-877-393-6486 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION

Revised: 4/2021

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Administration Instructions
- 2.2 Method of Reconstitution
- 2.3 Method of Use
- 2.4 Stability

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

- 4.1 Hypersensitivity

5 WARNINGS AND PRECAUTIONS

- 5.1 Cell Death
- 5.2 Hypotony
- 5.3 Cataract Formation
- 5.4 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS

- 6.1 Ophthalmic Adverse Reactions

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

NDA 22572/S-009

Page 4

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Mitosol[®] is an antimetabolite indicated for use as an adjunct to ab externo glaucoma surgery.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions

Mitosol[®] is intended for topical application to the surgical site of glaucoma filtration surgery. Mitosol[®] is a cytotoxic drug. It is not intended for intraocular administration. If intraocular administration occurs, cell death leading to corneal infarction, retinal infarction, and ciliary body atrophy may result. Verify pregnancy status in females of reproductive potential prior to using Mitosol[®].

2.2 Method of Reconstitution

Each vial of Mitosol[®] contains 0.2 mg of mitomycin and mannitol in a 1:2 concentration ratio. To reconstitute, add 1 mL of Sterile Water for Injection, then shake to dissolve. If product does not dissolve immediately, allow to stand at room temperature until the product dissolves into solution.

2.3 Method of Use

Sponges provided within the Mitosol[®] Kit should be fully saturated with the entire reconstituted contents in the manner prescribed in the Instructions for Use. A treatment area approximating 10mm x 6mm +/- 2mm should be treated with the Mitosol[®]. Apply fully saturated sponges equally to the treatment area, in a single layer, with the use of a surgical forceps. Keep the sponges on the treatment area for two (2) minutes, then remove and return to the Mitosol[®] Tray for defined disposal in the Chemotherapy Waste Bag provided.

2.4 Stability

Lyophilized Mitosol[®] stored at 20°C to 25°C (68°F to 77°F) is stable for the shelf life indicated on the package. Avoid excessive heat. Protect from light.

Reconstituted with 1 mL of Sterile Water for Injection at a concentration of 0.2 mg/mL, mitomycin is stable for one (1) hour at room temperature.

3. DOSAGE FORMS AND STRENGTHS

Mitosol[®] is a sterile lyophilized mixture of mitomycin and mannitol, which, when reconstituted with Sterile Water for Injection, provides a solution for application in glaucoma filtration surgery. Mitosol[®] is supplied in vials containing 0.2 mg of mitomycin. Each vial also contains mannitol 0.4 mg, at a 1:2 ratio of mitomycin to mannitol. Each mL of reconstituted solution contains 0.2 mg mitomycin and has a pH between 5.0 and 8.0.

4. CONTRAINDICATIONS

4.1 Hypersensitivity

Mitosol[®] is contraindicated in patients that have demonstrated a hypersensitivity to mitomycin in the past.

5. WARNINGS AND PRECAUTIONS

5.1 Cell Death

Mitomycin is cytotoxic. Use of mitomycin in concentrations higher than 0.2 mg/mL or use for longer than 2 minutes may lead to unintended corneal and/or scleral damage including thinning or perforation. Direct contact with the corneal endothelium will result in cell death.

5.2 Hypotony

The use of mitomycin has been associated with an increased incidence of post-operative hypotony.

NDA 22572/S-009

Page 5

5.3 Cataract Formation

Use in phakic patients has been correlated to a higher incidence of lenticular change and cataract formation.

5.4 Embryo-Fetal Toxicity

Based on findings in animals and mechanism of action, Mitosol® can cause fetal harm when administered to a pregnant woman. In animal reproduction studies, parenteral administration of mitomycin resulted in teratogenicity [see *Use in Specific Populations (8.1, 8.3) and Clinical Pharmacology (12.1)*].

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Cell Death [see *Warnings and Precautions (5.1)*]
- Hypotony [see *Warnings and Precautions (5.2)*]
- Cataract Formation [see *Warnings and Precautions (5.3)*]

6.1 Ophthalmic Adverse Reactions

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice. The most frequent adverse reactions to Mitosol® occur locally, as an extension of the pharmacological activity of the drug. These reactions include:

Blebitis: bleb ulceration, chronic bleb leak, encapsulated/cystic bleb, bleb-related infection, wound dehiscence, conjunctival necrosis, thin-walled bleb

Cornea: corneal endothelial damage, epithelial defect, anterior synechiae, superficial punctuate keratitis, Descemet's detachment, induced astigmatism

Endophthalmitis

Hypotony: choroidal reactions (choroidal detachment, choroidal effusion, serous choroidal detachment, suprachoroidal hemorrhage, hypotony maculopathy, presence of supraciliochoroidal fluid, hypoechogenic suprachoroidal effusion)

Inflammation: iritis, fibrin reaction

Lens: cataract development, cataract progression, capsule opacification, capsular constriction and/or capsulotomy rupture, posterior synechiae

Retina: retinal pigment epithelial tear, retinal detachment (serous and rhegmatogenous)

Scleritis: wound dehiscence

Vascular: hyphema, central retinal vein occlusion, hemiretinal vein occlusion, retinal hemorrhage, vitreal hemorrhage and blood clot, subconjunctival hemorrhage, disk hemorrhage

Additional Reactions: macular edema, sclera thinning or ulceration, intraocular lens capture, disk swelling, malignant glaucoma, lacrimal drainage system obstruction, ciliary block, corneal vascularization, visual acuity decrease, cystic conjunctival degeneration, upper eyelid retraction, dislocated implants, severe loss of vision.

NDA 22572/S-009

Page 6

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on findings in animals and mechanism of action [see *Clinical Pharmacology (12.1)*], Mitosol® can cause fetal harm when administered to a pregnant woman. There are no available data on Mitosol® use in pregnant women to inform the drug-associated risk. In animal reproduction studies, parenteral administration of mitomycin resulted in teratogenicity (see *Data*). Advise pregnant women of the potential risk to a fetus.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% - 4% and 15% - 20%, respectively.

Data

Animal Data

Parenteral administration of mitomycin in animal reproduction studies produced fetal malformations and embryofetal lethality.

8.2 Lactation

Risk Summary

There are no data on the presence of mitomycin in human milk, the effects on the breastfed child, or the effects on milk production. Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during and for 1 week following administration of Mitosol®.

8.3 Females and Males of Reproductive Potential

Mitosol® can cause fetal harm when administered to pregnant women [see *Use in Specific Populations (8.1)*].

Pregnancy Testing

Verify pregnancy status in females of reproductive potential prior to using Mitosol®.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

8.5 Geriatric Use

No overall differences in safety and effectiveness have been observed between elderly and younger patients.


11 DESCRIPTION

Mitomycin is an antibiotic isolated from the broth of *Streptomyces verticillus Yingtanensis* which has been shown to have antimetabolic activity.

Mitomycin is a blue-violet crystalline powder with the molecular formula of C₁₅H₁₈N₄O₅ and a molecular weight of 334.33. Its chemical name is 7-amino-9 α -methoxymitosane and it has the following structural formula:

NDA 22572/S-009

Page 7



Mitosol[®] is a sterile lyophilized mixture of mitomycin and mannitol, which, when reconstituted with Sterile Water for Injection, provides a solution for application in glaucoma filtration surgery. Mitosol[®] is supplied in vials containing 0.2 mg of mitomycin. Each vial also contains mannitol 0.4 mg, at a 1:2 ratio of mitomycin to mannitol. Each mL of reconstituted solution contains 0.2 mg mitomycin and has a pH between 5.0 and 8.0.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Mitosol[®] inhibits the synthesis of deoxyribonucleic acid (DNA). The guanine and cytosine content correlates with the degree of mitomycin-induced cross-linking. Cellular RNA and protein synthesis may also be suppressed.

12.3 Pharmacokinetics

Absorption

The systemic exposure of mitomycin following ocular administration of Mitosol[®] in humans is unknown. Based on a comparison of the proposed dose of up to 0.2 mg to intravenous (IV) doses of mitomycin used clinically for treatment of oncologic indications (up to 20 mg/m²), systemic concentrations in humans upon ocular administration are expected to be multiple orders of magnitude lower than those achieved by IV administration.

Elimination

Metabolism

In humans, mitomycin is cleared from ophthalmic tissue after intraoperative topical application and irrigation, as metabolism occurs in other affected tissues. Systemic clearance is affected primarily by metabolism in the liver. The rate of clearance is inversely proportional to the maximal serum concentration because of saturation of the degradative pathways.

Excretion

Approximately 10% of an injectable dose of mitomycin is excreted unchanged in the urine. Since metabolic pathways are saturated at relatively low doses, the percent of a dose excreted in urine increases.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Adequate long-term studies in animals to evaluate carcinogenic potential have not been conducted with Mitosol[®]. Intravenous administration of mitomycin has been found to be carcinogenic in rats and mice. At doses approximating the recommended clinical injectable dose in humans, mitomycin produces a greater than 100 percent increase in tumor incidence in male Sprague-Dawley rats, and a greater than 50 percent increase in tumor incidence in female Swiss mice.

The effect of Mitosol[®] on fertility is unknown.

NDA 22572/S-009

Page 8

14 CLINICAL STUDIES

In placebo-controlled studies reported in the medical literature, mitomycin reduced intraocular pressure (IOP) by 3 mmHg in patients with open-angle glaucoma when used as an adjunct to ab externo glaucoma surgery by Month 12.

In studies with a historical control reported in the medical literature, mitomycin reduced intraocular pressure (IOP) by 5 mmHg in patients with open-angle glaucoma when used as an adjunct to ab externo glaucoma surgery by Month 12.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Mitosol[®] (mitomycin for solution) is available in a kit containing:

One	Vial containing 0.2 mg mitomycin
One	1 mL syringe (Sterile Water For Injection) with Safety Connector
One	Plunger Rod
One	Vial Adapter with Spike
One	1 mL TB Syringe, Luer Lock
One	Sponge Container
Six	3 mm Absorbent Sponges
Six	6 mm Absorbent Sponges
Six	Half Moon Sponges
One	Instrument Wedge Sponge
One	Protective Foam Pouch
One	Chemotherapy Waste Bag
One	Label, MMC (mitomycin)

Three kits are supplied in each carton (NDC 49771-002-03).

16.2 Storage and Handling

Storage

Store kits at 20°C to 25°C (68°F to 77°F). Avoid excessive heat. Protect from light.

Handling Procedures

Mitosol[®] is a cytotoxic drug. Procedures for Proper Handling and Disposal of anti-cancer drugs should be followed. Appropriate containment and disposal devices are included within the Mitosol[®] (mitomycin for solution) Kit for Ophthalmic Use.

17 PATIENT COUNSELING INFORMATION

- Instruct patients to discuss with their physician if they are pregnant or if they might become pregnant [see *Use in Specific Populations (8.1)*].
- Instruct patients to discuss with their physician if they have demonstrated a hypersensitivity to mitomycin in the past [see *Contraindications (4.1)*].
- Nursing mothers should be advised that it is not known if Mitosol[®] is excreted in human milk. Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during and for 1 week following administration of Mitosol[®] [see *Use in Specific Populations (8.2)*].
- Patients should be advised of the toxicity of Mitosol[®] and potential complications.

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

NDA 22572/S-009

Page 9

Manufactured for:
Mobius Therapeutics, LLC
1000 Executive Parkway
Suite 224
St. Louis, MO 63141