


**CENTER FOR DRUG EVALUATION AND RESEARCH**

**APPLICATION NUMBER: 40-323**

**PRINTED LABELING**



BOTTLE LABEL  
480 mL


Margo

3 51079-888-44 1

1.5 mg/5 mL

EACH 5 mL (ONE TEASPOONFUL) CONTAINS 15 mg of PREDNISOLONE benzoate and 0.1 % sodium as a preservative. Also alcohol 5%.

**NDC 51079-888-44**



**PREDNISOLONE SYRUP, USP**

**15 mg/5 mL**

**240 mL**

**R only**

**USUAL DOSAGE:** See package insert for complete prescribing information.

Keep this and all drugs out of the reach of children.

**STORE AT CONTROLLED ROOM TEMPERATURE 15°-30°C (59°-86°F). DO NOT REFRIGERATE.**

Dispense in tight, light-resistant and child-resistant containers as defined in USP.

**PHARMACIST:** Dispense with a suitable Calibrated Measuring Device.

**LJL Laboratories, Inc.**  
Rockford, IL 61103

FP1021

BOTTLE LABEL  
240 mL

FF1023




**PREDNISOLONE  
SYRUP, USP**  
15 mg per 5 mL

**R**  
only

**DESCRIPTION:** Prednisolone syrup contains prednisolone which is a glucocorticoid. Glucocorticoids are adrenocortical steroids, both naturally occurring and synthetic, which are readily absorbed from the gastrointestinal tract. Prednisolone is a white to practically white, odorless, crystalline powder. It is very slightly soluble in water, soluble in methanol and in dioxane; sparingly soluble in acetone and alcohol, slightly soluble in chloroform.

The chemical name for Prednisolone is 11 $\beta$ ,17,21-Trihydroxypregna-1,4-diene-3,20-dione (anhydrous). Its molecular weight is 360.45. The molecular formula is C<sub>21</sub>H<sub>28</sub>O<sub>5</sub>, and the structural formula is:



Each 5 mL (teaspoonful) contains 15 mg of Prednisolone Syrup, USP. In addition, each 5 mL (teaspoonful) contains the following inactive ingredients: Benzoic acid 0.1% added as a preservative. It also contains alcohol 5%, citric acid, edetate disodium, FD&C red #40, flavor wild cherry, glycerin, propylene glycol, purified water, sodium saccharin and sucrose.

**CLINICAL PHARMACOLOGY:** Naturally occurring glucocorticoids (hydrocortisone and cortisone), which also have salt-retaining properties, are used as replacement therapy in adrenal deficiency states. synthetic analogs such as prednisolone are primarily used for their potent anti-inflammatory effects in disorders of many organ systems.

zoic acid 0.1% added as a preservative. It also contains alcohol 5%, citric acid, edetate disodium, FD&C red #40, flavor wild cherry, glycerin, propylene glycol, purified water, sodium saccharin and sucrose.

**CLINICAL PHARMACOLOGY:**

Naturally occurring glucocorticoids (hydrocortisone and cortisone), which also have salt-retaining properties, are used as replacement therapy in adrenal deficiency states. Synthetic analogs such as prednisolone are primarily used for their potent anti-inflammatory effects in disorders of many organ systems.

Glucocorticoids such as prednisolone cause profound and varied metabolic effects. In addition, they modify the body's immune responses to diverse stimuli.

**INDICATIONS AND USAGE:**

Prednisolone syrup is indicated in the following conditions:

**1. Endocrine Disorders**

Primary or secondary adrenocortical insufficiency (hydrocortisone or cortisone is the first choice; synthetic analogs may be used in conjunction with mineralocorticoids where applicable; in infancy mineralocorticoid supplementation is of particular importance).

Congenital adrenal hyperplasia  
Nonsuppurative thyroiditis  
Hypercalcemia associated with cancer

**2. Rheumatic Disorders**

As adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in:

Psoriatic arthritis  
Rheumatoid arthritis, including juvenile rheumatoid arthritis (selected cases may require low-dose maintenance therapy)  
Ankylosing spondylitis  
Acute and subacute bursitis  
Acute nonspecific tenosynovitis  
Acute gouty arthritis  
Post-traumatic osteoarthritis  
Synovitis of osteoarthritis  
Epicondylitis

**3. Collagen Diseases**

During an exacerbation or as maintenance therapy in selected cases of:

Systemic Lupus erythematosus  
Acute rheumatic carditis

**4. Dermatologic Diseases**

Pemphigus  
Bullous dermatitis herpetiformis  
Severe erythema multiforme (Stevens-Johnson syndrome)  
Exfoliative dermatitis  
Mycosis fungoides  
Severe psoriasis  
Severe seborrheic dermatitis

**5. Allergic States**

Control of severe or incapacitating allergic conditions intractable to adequate trials of conventional treatment:

5  
Cutaneous eruptions  
herpetiformis  
Severe erythema  
multiforme (Stevens-  
Johnson syndrome)  
Exfoliative dermatitis  
Mycosis fungoides  
Severe psoriasis  
Severe seborrheic  
dermatitis

**5. Allergic States**

Control of severe or  
incapacitating allergic  
conditions intractable  
to adequate trials of  
conventional treatment:

Seasonal or  
perennial  
allergic rhinitis  
Bronchial asthma  
Contact dermatitis  
Atopic dermatitis  
Serum sickness  
Drug hypersensitivity  
reactions

**6. Ophthalmic Diseases**

Severe acute and  
chronic allergic and in-  
flammatory processes  
involving the eye and  
its adnexa such as:

Allergic corneal  
marginal ulcers  
Herpes zoster  
ophthalmicus  
Anterior segment  
inflammation  
Diffuse posterior  
uveitis and  
choroiditis  
Sympathetic  
ophthalmia  
Allergic conjunctivitis  
Keratitis  
Chorioretinitis  
Optic neuritis  
Iritis and Iridocyclitis

**7. Respiratory Diseases**

Symptomatic  
sarcoidosis  
Loeffler's syndrome  
not manageable by  
other means  
Berylliosis  
Fulminating or  
disseminated  
pulmonary  
tuberculosis when  
used concurrently  
with appropriate  
chemotherapy  
Aspiration  
pneumonitis

**8. Hematologic Disorders**

Idiopathic  
thrombocytopenic  
purpura in adults  
Secondary  
thrombocytopenia  
in adults  
Acquired  
(autoimmune)  
hemolytic anemia  
Erythroblastopenia  
(RBC anemia)  
Congenital  
(erythroid)  
hypoplastic anemia

**9. Neoplastic Diseases**

For palliative manage-  
ment of:

Leukemias and  
lymphomas in  
adults  
Acute leukemia of  
childhood

**10. Edematous States**

To induce a diuresis or  
remission of proteinuria  
in the nephrotic syn-  
drome, without uremia,  
of the idiopathic type  
or that due to lupus  
erythematosus.

**11. Gastrointestinal  
Diseases**

To tide the patient over  
a critical period of the  
disease in:

Ulcerative colitis  
Regional enteritis

**12. Miscellaneous**

Tuberculosis meningitis

...hematoma of  
childhood

6

**10. Edematous States**  
To induce a diuresis or remission of proteinuria in the nephrotic syndrome, without uremia, of the idiopathic type or that due to lupus erythematosus.

**11. Gastrointestinal Diseases**  
To tide the patient over a critical period of the disease in:  
Ulcerative colitis  
Regional enteritis

**12. Miscellaneous**  
Tuberculous meningitis with subarachnoid block or impending block used concurrently with appropriate anti-tuberculous chemotherapy. Trichinosis with neurologic or myocardial involvement.

In addition to the above indications prednisolone syrup is indicated for systemic dermatomyositis (polymyositis).

**CONTRAINDICATIONS.**  
Systemic fungal infections.

**WARNINGS:** In patients on corticosteroid therapy subjected to unusual stress, increased dosage of rapidly acting corticosteroids before, during, and after the stressful situation is indicated.

Corticosteroids may mask some signs of infection, and new infections may appear during their use. There may be decreased resistance and inability to localize infection when corticosteroids are used.

Prolonged use of corticosteroids may produce posterior subcapsular cataracts, glaucoma with possible damage to the optic nerves, and may enhance the establishment of secondary ocular infections due to fungi or viruses.

Average and large doses of hydrocortisone or cortisone can cause elevation of blood pressure, salt and water retention, and increased excretion of potassium. These effects are less likely to occur with the synthetic derivatives except when used in large doses. Dietary salt restriction and potassium supplementation may be necessary. All corticosteroids increase calcium excretion.

**While on corticosteroid therapy, patients should not be vaccinated against smallpox. Other immunization procedures should not be undertaken in patients who are on corticosteroids, especially on high dose, because of possible hazards of neurological complications and a lack of antibody response.**

Persons who are on drugs which suppress the immune system are more susceptible to infections than healthy individuals. Chickenpox and measles, for example, can have a more serious or even fatal course in non-immune children or adults on corticosteroids. In such children or adults who have not had these diseases, particular care should be taken to avoid exposure. How the dose, route and duration of corticosteroid administration affects the

7

Corticosteroids may mask some signs of infection, and new infections may appear during their use. There may be decreased resistance and inability to localize infection when corticosteroids are used.

Prolonged use of corticosteroids may produce posterior subcapsular cataracts, glaucoma with possible damage to the optic nerves, and may enhance the establishment of secondary ocular infections due to fungi or viruses.

Average and large doses of hydrocortisone or cortisone can cause elevation of blood pressure, salt and water retention, and increased excretion of potassium. These effects are less likely to occur with the synthetic derivatives except when used in large doses. Dietary salt restriction and potassium supplementation may be necessary. All corticosteroids increase calcium excretion.

While on corticosteroid therapy, patients should not be vaccinated against smallpox. Other immunization procedures should not be undertaken in patients who are on corticosteroids, especially on high dose, because of possible hazards of neurological complications and a lack of antibody response.

Persons who are on drugs which suppress the immune system are more susceptible to infections than healthy individuals. Chickenpox and measles, for example, can have a more serious or even fatal course in non-immune children or adults on corticosteroids. In such children or adults who have not had these diseases, particular care should be taken to avoid exposure. How the dose, route and duration of corticosteroid administration affects the risk of developing a disseminated infection is not known. The contribution of the underlying disease and/or prior corticosteroid treatment to the risk is also not known. If exposed to chickenpox, prophylaxis with varicella zoster immune globulin (VZIG) may be indicated. If exposed to measles, prophylaxis with pooled intravenous immunoglobulin (IVIG) may be indicated. (See the respective package inserts for complete VZIG and IVIG prescribing information). If chickenpox develops, treatment with antiviral agents may be considered.

The use of prednisolone syrup in active tuberculosis

should be restricted to those cases of fulminating or disseminated tuberculosis in which the corticosteroid is used for the management of the disease in conjunction with an appropriate antituberculous regimen.

If corticosteroids are indicated in patients with latent tuberculosis or tuberculin reactivity, dose observation is necessary as reactivation of the disease may occur. During prolonged corticosteroid therapy, these patients should receive chemoprophylaxis.

*Use in pregnancy:* Since adequate human reproduction studies have not been done with corticosteroids, the use of these drugs in pregnancies, nursing mothers or women of childbearing potential requires that the possible benefits of the drug be weighed against the potential hazards to the mother and embryo or fetus. Infants born of mothers who have received substantial doses of corticosteroids during pregnancy should be carefully observed for signs of hypoadrenalism.

**PRECAUTIONS:**

**General:** Drug-induced secondary adrenocortical insufficiency may be minimized by gradual reduction of dosage. This type of relative insufficiency may persist for months after discontinuation of therapy; therefore, in any situation of stress occurring during that period, hormone therapy should be reinstated. Since mineralocorticoid secretion may be impaired, salt and/or a mineralocorticoid should be administered concurrently.

There is an enhanced effect of corticosteroids on patients with hypothyroidism and in those with cirrhosis.

Corticosteroids should be used cautiously in patients with ocular herpes simplex because of possible corneal perforation.

The lowest possible dose of corticosteroid should be used to control the condition under treatment, and when reduction in dosage is possible, the reduction should be gradual.

Psychic derangements may appear when corticosteroids are used, ranging from euphoria, insomnia, mood swings, personality changes, and severe depression, to frank psychotic manifestations. Also, existing emotional instability or psychotic tendencies may be aggravated by corticosteroids.

Aspirin should be used cautiously in conjunction with corticosteroids in hypoprothrombinemia.

Steroids should be used with caution in nonspecific Ulcerative Colitis if there is a probability of impending perforation, abscess or other pyogenic infections, diverticulitis, fresh intestinal anastomoses, active or latent peptic ulcer, renal insufficiency, hypertension, osteoporosis, and myasthe-

9

Aspirin should be used cautiously in conjunction with corticosteroids in hypoprothrombinemia.

Steroids should be used with caution in nonspecific Ulcerative Colitis if there is a probability of impending perforation, abscess or other pyogenic infections, diverticulitis, fresh intestinal anastomoses, active or latent peptic ulcer, renal insufficiency, hypertension, osteoporosis, and myasthenia gravis.

Growth and development of infants and children on prolonged corticosteroid therapy should be carefully observed.

Information for patients: Patients who are on immunosuppressant doses of corticosteroids should be warned to avoid exposure to chickenpox or measles. Patients should also be advised that if they are exposed, medical advice should be sought without delay.

**ADVERSE REACTIONS:**

**Fluid and Electrolyte**

**Disturbances**

Sodium retention  
Fluid retention  
Congestive heart failure in susceptible patients  
Potassium loss  
Hypokalemic alkalosis  
Hypertension

**Musculoskeletal**

Muscle weakness  
Steroid myopathy  
Loss of muscle mass  
Osteoporosis  
Vertebral compression fractures  
Aseptic necrosis of femoral and humeral heads  
Pathologic fracture of long bones

**Gastrointestinal**

Peptic ulcer with possible perforation and hemorrhage  
Pancreatitis  
Abdominal distention  
Ulcerative esophagitis

**Dermatologic**

Impaired wound healing  
Thin fragile skin  
Petechiae and ecchymoses  
Facial erythema  
Increased sweating  
May suppress reactions to skin tests

**Neurological**

Convulsions  
Increased intracranial pressure with papilledema (pseudo-tumor cerebri) usually after treatment  
Vertigo  
Headache

**Endocrine**

Menstrual irregularities  
Development of Cushingoid state  
Suppression of growth in pediatric patients  
Secondary adrenocortical and pituitary unresponsiveness, particularly in times of stress, as in trauma, surgery or illness  
Decreased carbohydrate tolerance  
Manifestations of latent diabetes mellitus  
Increased requirements for insulin or oral hypoglycemic agents in diabetics

**Ophthalmic**

Posterior subcapsular cataracts  
Increased intraocular pressure

10

**Endocrine**

Menstrual irregularities  
Development of  
Cushingoid state  
Suppression of growth  
in pediatric patients  
Secondary  
adrenocortical and  
pituitary unresponsive-  
ness, particularly  
in times of stress, as in  
trauma, surgery or  
illness  
Decreased carbohy-  
drate tolerance  
Manifestations of latent  
diabetes mellitus  
Increased requirements  
for insulin or oral  
hypoglycemic agents  
in diabetics

**Ophthalmic**

Posterior subcapsular  
cataracts  
Increased intraocular  
pressure  
Glaucoma  
Exophthalmos

**Metabolic**

Negative nitrogen  
balance due to protein  
catabolism

**DOSAGE AND ADMINISTRATION:**

Dosage of prednisolone syrup should be individualized according to the severity of the disease and the response of the patient. For pediatric patients, the recommended dosage should be governed by the same considerations rather than strict adherence to the ratio indicated by age or body weight.

Hormone therapy is an adjunct to and not a replacement for conventional therapy.

Dosage should be decreased or discontinued gradually when the drug has been administered for more than a few days.

The severity, prognosis, expected duration of the disease, and the reaction of the patient to medication are primary factors in determining dosage.

If a period of spontaneous remission occurs in a chronic condition, treatment should be discontinued.

Blood pressure, body weight, routine laboratory studies, including two-hour postprandial blood glucose and serum potassium, and a chest X-ray should be obtained at regular intervals during prolonged therapy. Upper GI X-rays are desirable in patients with known or suspected peptic ulcer disease.

The initial dosage of prednisolone syrup may vary from 5 mg to 60 mg per day depending on the specific disease entity being treated. In situations of less severity lower doses will generally suffice while in selected patients higher initial doses may be required. The initial dosage should be maintained or adjusted until a satisfactory response is noted. If after a reasonable period of time there is a lack of satisfactory clinical response, prednisolone syrup should be discontinued and the patient transferred to other appropriate therapy. **IT SHOULD BE EMPHASIZED THAT DOSAGE REQUIREMENTS ARE VARIABLE AND MUST BE INDIVIDUALIZED ON THE BASIS OF**

The initial dosage should be maintained or adjusted until a satisfactory response is noted. If after a reasonable period of time there is a lack of satisfactory clinical response, prednisolone syrup should be discontinued and the patient transferred to other appropriate therapy. **IT SHOULD BE EMPHASIZED THAT DOSAGE REQUIREMENTS ARE VARIABLE AND MUST BE INDIVIDUALIZED ON THE BASIS OF THE DISEASE UNDER TREATMENT AND THE RESPONSE OF THE PATIENT.**

After a favorable response is noted, the proper maintenance dosage should be determined by decreasing the initial drug dosage in small decrements at appropriate time intervals until the lowest dosage which will maintain an adequate clinical response is reached. It should be kept in mind that constant monitoring is needed in regard to drug dosage. Included in the situations which may make dosage adjustments necessary are changes in clinical status secondary to remissions or exacerbations in the disease process, the patient's individual drug responsiveness, and the effect of patient exposure to stressful situations not directly related to the disease entity under treatment. In this latter situation it may be necessary to increase the dosage of prednisolone syrup for a period of time consistent with the patient's condition. If after long-term therapy the drug is to be stopped, it is recommended that it be withdrawn gradually rather than abruptly.

**HOW SUPPLIED:** Prednisolone Syrup, USP is a cherry flavored red liquid containing 15 mg of Prednisolone in each 5 mL (teaspoonful). It is supplied as follows:

NDC 51079-888-44

Bottles of 240 mL

NDC 51079-888-38

Bottles of 480 mL

**Pharmacist:** Dispense with a suitable calibrated measuring device to assure proper measuring of dose.

**Dose/Volume Chart**

15 mg prednisolone =  
1 teaspoon

10 mg prednisolone =  
2/3 teaspoon


7.5 mg prednisolone =  
1/2 teaspoon

5 mg prednisolone =  
1/3 teaspoon

Dispense in tight, light-resistant and child-resistant containers as defined in USP/NF.

Store at controlled room temperature 15°-30°C (59°-86°F). Do Not Refrigerate.



 UDL Laboratories, Inc.  
Rockford, IL 61103

FP1023  
11/98