

061813EN

PMS Black

Erythrocin™ Lactobionate - IV

Erythromycin Lactobionate for Injection, USP

INTRAVENOUS USE ONLY
Vials

Rx only



Erythrocin™ Lactobionate - IV

Erythromycin Lactobionate for Injection, USP

INTRAVENOUS USE ONLY
Vials

Rx only



Hospira, Inc., Lake Forest, IL 60045 USA

Revised: 01/2013

1. Clinical and Laboratory Standards Institute. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically*. Approved Standard—9th Edition. Wayne, Pennsylvania: Clinical and Laboratory Standards Institute, 2012.
2. Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Disk Susceptibility Tests*. Approved Standard—11th Edition. Wayne, Pennsylvania: Clinical and Laboratory Standards Institute, 2012.
3. Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing*. 22nd International Supplement. CLSI document M100-S22. Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087-1898 USA, 2012.
4. Committee on Rheumatic Fever and Infective Endocarditis of the Council on Cardiovascular Disease of the Young. Prevention of Rheumatic Fever, *Circulation* 70(6):1184-1122A, December 1984.
5. Committee on Rheumatic Fever and Infective Endocarditis of the Council on Cardiovascular Disease of the Young. Prevention of Bacterial Endocarditis. *Circulation* 70(6):1123A-1127A, December 1984.
6. Gitter, B. et al. *Torsades de Pointes Induced by Erythromycin*. *Chest*, Volume 105: 368-72, February 1994.

REFERENCES
 Clinical and Laboratory Standards Institute. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically*. Approved Standard—9th Edition. Wayne, Pennsylvania: Clinical and Laboratory Standards Institute, 2012.

HOW SUPPLIED
 Erythrocin Lactobionate IV (erythromycin lactobionate for injection, USP) is supplied as a sterile, lyophilized powder in packages of ten vials (NDC 0409-6482-01), each vial containing the equivalent of 500 mg of erythromycin.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution storage.

The final diluted solution of erythromycin lactobionate should be completely administered within 8 hours, since it is not suitable for storage.

Physical stability of the solution has first been determined.

No drug or chemical agent should be added to an erythromycin lactobionate-IV fluid admixture unless its effect on the chemical and physical stability of the final diluted solution of erythromycin lactobionate.

Desirable for the final diluted solution of erythromycin lactobionate are unstable and lose their potency rapidly. A pH of at least 5.5 is acceptable for the final diluted solution of erythromycin lactobionate.

Neutral (pH) sodium lactobionate, USP, must be added to these solutions so that their pH is in the optimum range for erythromycin lactobionate stability. Each vial contains the equivalent of 500 mg of erythromycin lactobionate and 0.9% sodium chloride injection, USP.

5% DEXTROSE AND LACTATED RINGERS INJECTION, USP
 5% DEXTROSE INJECTION, USP
 HOSPIRA) by adding 1 mL of Neut™ per 100 mL of solution.

3. THE FOLLOWING SOLUTIONS MAY ALSO BE USED PROVIDING THEY ARE FIRST BUFFERED WITH NEUT™ (4% SODIUM BICARBONATE, 0.9% SODIUM CHLORIDE INJECTION, USP; LACTATED RINGERS INJECTION, USP; NORMOSOL™-R).

erythromycin activity per liter (1 mg/mL) for continuous infusion or 1 to 2 mg/mL for intermittent infusion:
 2. ADD THE INITIAL DILUTION TO ONE OF THE FOLLOWING DILUENTS BEFORE ADMINISTRATION TO GIVE A CONCENTRATION OF 1 g OR 2 g PER 24 HOURS AT ROOM TEMPERATURE.

After reconstitution, each mL contains 50 mg of erythromycin base. The initial solution is stable at refrigerator temperature for two weeks, or for 24 hours at room temperature.

1. PREPARE THE INITIAL SOLUTION OF ERYTHROCIIN LACTOBIONATE IV BY ADDING 10 mL OF STERILE WATER FOR INJECTION, USP, TO THE 500 MG VIAL. Use only Sterile Water for Injection, USP, as other diluents may cause precipitation during reconstitution. Do not use diluents containing preservatives or inorganic salts.

Preparation of Solution
 Erythrocin Lactobionate IV (erythromycin lactobionate for injection, USP) is supplied as a sterile, lyophilized powder in packages of ten vials (NDC 0409-6482-01), each vial containing the equivalent of 500 mg of erythromycin.

In prophylaxis against bacterial endocarditis (see **INDICATIONS AND USAGE**), the oral regimen for penicillin allergic patients is erythromycin 1 gram, 1 hour before the procedure followed by 500 mg six hours later.

Administration of doses of ≥ 4 g/day may increase the risk for the development of erythromycin-induced hearing loss in elderly patients, particularly those with reduced renal or hepatic function.

For treatment of severe infections in adults and pediatric patients, the recommended intravenous dose of erythromycin lactobionate is 1 to 2 g q4h/day, higher doses, up to 4 g q4h/day, may be given for severe infections.

DOSE AND ADMINISTRATION
 Erythromycin is not removed by peritoneal dialysis or hemodialysis.

OVERDOSE
 In the case of overdose, erythromycin infusion should be discontinued and all other appropriate measures should be instituted.

Continuous infusion of erythromycin lactobionate is preferred due to the slower infusion rate and lower concentration of erythromycin base. However, intermittent infusion at six hour intervals is also effective.

High doses of erythromycin. The final diluted solution of erythromycin lactobionate is prepared to give a concentration of 1 g per liter (1 mg/mL).

For slow continuous infusion: The final diluted solution of erythromycin lactobionate is prepared to give a concentration of 1 g per liter (1 mg/mL).

For intermittent infusion: administer one-fourth the total daily dose of erythromycin lactobionate by intravenous infusion in 20 to 60 minutes at intervals not greater than every six hours. The final diluted solution of erythromycin lactobionate is prepared to give a 60 mg/mL concentration of 1 to 5 mg/mL. (No less than 100 mL of IV fluid should be used. Infusion should be subsided slowly to minimize fluid along the vein.)

For treatment of acute pelvic inflammatory disease caused by *N. gonorrhoeae*, in female patients hypersensitive to penicillins, administer 500 mg erythromycin lactobionate every six hours for three days, followed by oral administration of 250 mg erythromycin stearate or base every six hours for seven days.

For treatment of Legionnaires' Disease: Although optimal doses have not been established, doses utilized in reported clinical case were bases every six hours for seven days.

Administration of doses of ≥ 4 g/day may increase the risk for the development of erythromycin-induced hearing loss in elderly patients, particularly those with reduced renal or hepatic function.

For treatment of severe infections in adults and pediatric patients, the recommended intravenous dose of erythromycin lactobionate is 1 to 2 g q4h/day, higher doses, up to 4 g q4h/day, may be given for severe infections.

ADVERSE REACTIONS
 Erythromycin has been associated with QT prolongation and ventricular arrhythmias, including ventricular tachycardia and torsades de pointes. (See **WARNINGS**.)

Side effects following the use of intravenous erythromycin are rare. Occasional inious irritation has been encountered, but if the infusion is given slowly, dilute solution, preferably by continuous intravenous infusion or intermittent infusion in no less than 20 to 60 minutes, allergic reactions ranging from urticaria to anaphylaxis have occurred. Skin reactions ranging from mild eruptions to erythema multiforme, Stevens-Johnson syndrome, and toxic epidermal necrolysis have been reported rarely.

There have been isolated reports of reversible hearing loss occurring chiefly in patients with renal insufficiency and in patients receiving high doses of erythromycin.

Elderly patients, particularly those with reduced renal or hepatic function, may also be at increased risk for developing this effect when receiving doses of 4 grams/day or higher are given. (See **DOSE AND ADMINISTRATION**.)

Diarrhea is a common problem caused by antibiotics which usually ends when the antibiotic is discontinued. Sometimes as late as two or more months after having taken the last dose of the antibiotic, if this occurs, patients should contact their physician as soon as possible.

ADVERSE REACTIONS AND DOSAGE AND ADMINISTRATION
 Elderly patients may experience increased effects of oral anticoagulant therapy while undergoing treatment with Erythrocin™. (See **PRECAUTIONS, Drug Interactions**.)

Information for Patients
 Patients should be counseled that antibacterial drugs including erythromycin should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When erythromycin is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not taking all of the medication may decrease the effectiveness of the immediate and long-term treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by erythromycin or other antibacterial drugs in the future.

Information for Patients
 Patients should be counseled that antibacterial drugs including erythromycin should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When erythromycin is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not taking all of the medication may decrease the effectiveness of the immediate and long-term treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by erythromycin or other antibacterial drugs in the future.

PRECAUTIONS, Drug Interactions
 Erythrocin Lactobionate does not contain sodium.

ADVERSE REACTIONS AND DOSAGE AND ADMINISTRATION
 Elderly patients may experience increased effects of oral anticoagulant therapy while undergoing treatment with Erythrocin™. (See **ADVERSE REACTIONS**.)

Geiatric Use
 Administration of doses of ≥ 4 g/day may increase the risk for the development of erythromycin-induced hearing loss in elderly patients, particularly those with reduced renal or hepatic function.

Pediatric Use
 Erythromycin is excreted in breast milk. Caution should be exercised when erythromycin is administered to a nursing woman.

Nursing Mothers
 The effect of erythromycin on labor and delivery is unknown.

Labor and Delivery
 fetal plasma levels are generally low.

It should be used during pregnancy only if clearly needed. Erythromycin has been reported to cross the placental barrier in humans, but well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if the potential benefits outweigh the risks.

Pregnancy Category B
 There was no evidence of teratogenicity or any other adverse effects through gestation, and during gestation, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if the potential benefits outweigh the risks.

Fertility
 Long-term animal data studies in rats with erythromycin ethylsuccinate and erythromycin base did not provide evidence of reproductive toxicity. Mutagenicity studies in rats with erythromycin base have not been conducted. There was no apparent effect on male or female fertility in rats fed erythromycin (base) at levels up to 0.25% of diet.

Carcinogenesis, Mutagenesis, Impairment of Fertility
 Long-term animal data studies in rats with erythromycin ethylsuccinate and erythromycin base did not provide evidence of reproductive toxicity. Mutagenicity studies in rats with erythromycin base have not been conducted. There was no apparent effect on male or female fertility in rats fed erythromycin (base) at levels up to 0.25% of diet.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.


Concomitant Administration
 Concomitant administration of erythromycin and digoxin has been reported to result in elevated serum digoxin levels.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of erythromycin and other antibacterial drugs, erythromycin should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

DESCRIPTION
 Erythromycin is produced by a strain of *Streptomyces erythraeus* and belongs to the macrolide group of antibiotics. It is basic and readily forms salts with acids.

Erythrocin Lactobionate (erythromycin lactobionate for injection, USP), is a soluble salt of erythromycin suitable for intravenous administration. It is available as a sterile, lyophilized powder in vials containing the equivalent of 500 mg of erythromycin activity. It is prepared as a solution and lyophilized in its final container.

Erythromycin lactobionate is chemically known as erythromycin mono (4-0-β-D-galactopyranosyl-D-gluconate) (salt). The structural formula is:



CLINICAL PHARMACOLOGY
 Erythromycin diffuses readily into most body fluids. In the absence of meningeal inflammation, low concentrations are normally achieved in the spinal fluid but the passage of the drug across the blood-brain barrier increases in meningitis. Erythromycin crosses the placental barrier and is excreted in breast milk. Erythromycin is not removed by peritoneal dialysis or hemodialysis.

In the presence of normal hepatic function, erythromycin is concentrated in the liver and is excreted in the bile; the effect of hepatic dysfunction on biliary excretion of erythromycin is not known. From 12 to 15 percent of intravenously administered erythromycin is excreted in active form in the urine.

Intravenous infusion of 500 mg of erythromycin lactobionate at a constant rate over 1 hour in fasting adults produced a mean serum erythromycin level of approximately 7 mcg/mL at 20 minutes, 10 mcg/mL at 1 hour, 2.6 mcg/mL at 2.5 hours, and 1 mcg/mL at 6 hours.

Microbiology
 Erythromycin is a macrolide antibiotic with activity against Gram-positive and Gram-negative bacteria.

Mechanism of Action
 Erythromycin acts by inhibition of protein synthesis by binding 50 S ribosomal subunits of susceptible organisms. It does not affect nucleic acid synthesis.

Interactions with Other Antibiotics
 Antagonism has been demonstrated *in vitro* between erythromycin and clindamycin, lincomycin and chloramphenicol. Many strains of *Haemophilus influenzae* are resistant to erythromycin, but are susceptible to erythromycin and sulfonamides used concomitantly.

Development of Resistance
 Resistance to erythromycin in *S. aureus* may emerge during therapy.

Erythromycin has been shown to be active against most strains of the following organisms both *in vitro* and in clinical infections (see **INDICATIONS AND USAGE**):

- Gram-positive bacteria Aerobic**
 - Corynebacterium diphtheriae*
 - Corynebacterium minutissimum*
 - Staphylococcus aureus* (methicillin-susceptible strains only)
 - Streptococcus pneumoniae*
 - Streptococcus pyogenes*

- Gram-negative bacteria**
 - Legionella pneumophila*
 - Neisseria gonorrhoeae*

- Other Microorganisms**
 - Mycoplasma pneumoniae*

At least 90 percent of the following microorganisms exhibit an *in vitro* minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoint for erythromycin. However, the efficacy of erythromycin in treating clinical infections due to these microorganisms has not been established in adequate and well-controlled trials.

Gram-negative bacteria
Moraxella catarrhalis

Susceptibility Testing
 When available, the clinical microbiology laboratory should provide cumulative results of the *in vitro* susceptibility test results for antimicrobial drugs used in resident hospitals to the physician as periodic reports that describe the susceptibility profile of nosocomial and community-acquired pathogens. These reports should aid the physician in selecting the most effective antimicrobial.

Dilution techniques
 Quantitative methods are used to determine antimicrobial minimum inhibitory concentrations (MICs). These MICs provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MICs should be determined using a standardized procedure. Standardized procedures are based on a dilution method (broth or agar)¹ or equivalent with standardized inoculum concentrations and standardized concentrations of erythromycin powder. The MIC values should be interpreted according to the criteria provided in Table 1.

Diffusion technique
 Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such standardized procedure² requires the use of standardized inoculum concentrations. This procedure uses paper disks impregnated with 15 mcg of erythromycin to test the susceptibility of microorganisms to erythromycin. The disk diffusion interpretive criteria are provided in Table 1.

Pathogen	Susceptibility Interpretive Criteria					
	Minimum Inhibitory Concentration (mcg/mL)			Disk Diffusion (zone diameter in mm)		
	S	I	R	S	I	R
<i>Staphylococcus spp.</i>	≤0.5	1 to 4	≥8	≥23	14 to 22	≤13
<i>Enterococcus spp.</i>	≤0.5	1 to 4	≥8	≥23	14 to 22	≤13
<i>Streptococcus pneumoniae</i> ^{a,b}	≤0.25	0.5	≥1	≥21	16 to 20	≤15
<i>Streptococcus spp. (β-hemolytic group)</i> ^{a,b}	≤0.25	0.5	≥1	≥21	16 to 20	≤15
<i>Streptococcus spp. (Viridans group)</i> ^{a,b}	≤0.25	0.5	≥1	≥21	16 to 20	≤15

^a The MIC interpretive criteria for *Streptococcus pneumoniae*, *Streptococcus spp. (β-hemolytic group)*, and *Streptococcus spp. (Viridans group)* are applicable only to tests performed by broth microdilution using cation-adjusted Mueller-Hinton broth supplemented with 2 to 5% lysed horse blood inoculated with a direct colony suspension and incubated in ambient air at 35 ± 2°C for 20 to 24 hours.

^b The zone diameter interpretive criteria for *Streptococcus pneumoniae*, *Streptococcus spp. (β-hemolytic group)*, and *Streptococcus spp. (Viridans group)* are applicable only to tests performed using Mueller-Hinton agar supplemented with 5% defibrinated sheep blood inoculated with a direct colony suspension and incubated in 5% CO₂ at 35 ± 2°C for 20 to 24 hours.

A report of *Susceptible* indicates that the antimicrobial is likely to inhibit growth of the pathogen if the antimicrobial compound in the blood reaches the concentrations usually achievable. A report of *Intermediate* indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where a high dosage of drug can be used. This category also provides a buffer zone, which prevents small, uncontrolled technical factors from causing major discrepancies in interpretation. A report of *Resistant* indicates that the antimicrobial is not likely to inhibit growth of the pathogen if the antimicrobial compound in the blood reaches the concentrations usually achievable and other therapy should be selected.

Quality Control
 Standardized susceptibility test procedures require the use of quality control microorganisms to control the technical aspects of the test procedures³. Standard Erythromycin powder should provide the following range of values noted in Table 2.

QC Strain	Acceptable Quality Control Ranges	
	Minimum Inhibitory Concentration (mcg/mL)	Disk Diffusion (zone diameter in mm)
<i>Enterococcus faecalis</i> ATCC 29212	1 to 4	NA ^a
<i>Staphylococcus aureus</i> ATCC 29213	0.25 to 1	NA
<i>Staphylococcus aureus</i> ATCC 25923	NA	22 to 30
<i>Streptococcus pneumoniae</i> ATCC 49619 ^b	0.03 to 0.12 ^c	25 to 30 ^d

^a not applicable
^b This organism may be used for validation of susceptibility test results when testing *Streptococcus spp.* other than *S. pneumoniae*.
^c This quality control range for *S. pneumoniae* is applicable only to tests performed by broth microdilution using cation-adjusted Mueller-Hinton broth supplemented with 2 to 5% lysed horse blood inoculated with a direct colony suspension and incubated in ambient air at 35 ± 2°C for 20 to 24 hours.

^d This quality control zone diameter range is applicable only to tests performed using Mueller-Hinton agar supplemented with 5% defibrinated sheep blood inoculated with a direct colony suspension and incubated in 5% CO₂ at 35 ± 2°C for 20 to 24 hours.

INDICATIONS AND USAGE
 Erythrocin Lactobionate IV (erythromycin lactobionate for injection,