

Cleocin Phosphate (clindamycin phosphate) Injection, Solution
[Pharmacia & Upjohn, Co.]

To reduce the development of drug-resistant bacteria and maintain the effectiveness of CLEOCIN PHOSPHATE and other antibacterial drugs, CLEOCIN PHOSPHATE should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

Sterile Solution is for Intramuscular and Intravenous Use

CLEOCIN PHOSPHATE in the ADD-Vantage™ Vial is For Intravenous Use Only

WARNING

Clostridium difficile associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including CLEOCIN PHOSPHATE and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

Because CLEOCIN PHOSPHATE therapy has been associated with severe colitis which may end fatally, it should be reserved for serious infections where less toxic antimicrobial agents are inappropriate, as described in the **INDICATIONS AND USAGE** section. It should not be used in patients with nonbacterial infections such as most upper respiratory tract infections.

C. difficile produces toxins A and B which contribute to the development of CDAD.

Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.


DESCRIPTION

CLEOCIN PHOSPHATE Sterile Solution in vials contains clindamycin phosphate, a water soluble ester of clindamycin and phosphoric acid. Each mL contains the equivalent of 150 mg clindamycin, 0.5 mg disodium edetate and 9.45 mg benzyl alcohol added as preservative in each mL. Clindamycin is a semisynthetic antibiotic produced by a 7(S)-chloro-substitution of the 7(R)-hydroxyl group of the parent compound lincomycin.

The chemical name of clindamycin phosphate is L-threo- α -D-galacto- Octopyranoside, methyl 7-chloro-6,7,8-trideoxy-6-[[[(1-methyl-4-propyl-2- pyrrolidiny]l) carbonyl] amino]-1-thio-, 2-(dihydrogen phosphate), (2S-trans)-.

The molecular formula is C₁₈H₃₄ClN₂O₈PS and the molecular weight is 504.96.

The structural formula is represented below:



CLEOCIN PHOSPHATE in the ADD-Vantage Vial is intended for intravenous use only after further dilution with appropriate volume of ADD-Vantage diluent base solution.

CLEOCIN PHOSPHATE IV Solution in the Galaxy[®] plastic container for intravenous use is composed of clindamycin phosphate equivalent to 300, 600 and 900 mg of clindamycin premixed with 5% dextrose as a sterile solution. Disodium edetate has been added at a concentration of 0.04 mg/mL. The pH has been adjusted with sodium hydroxide and/or hydrochloric acid.

The plastic container is fabricated from a specially designed multilayer plastic, PL 2501. Solutions in contact with the plastic container can leach out certain of its chemical components in very small amounts within the expiration period. The suitability of the plastic has been confirmed in tests in animals according to the USP biological tests for plastic containers, as well as by tissue culture toxicity studies.

CLINICAL PHARMACOLOGY

Biologically inactive clindamycin phosphate is rapidly converted to active clindamycin.

By the end of short-term intravenous infusion, peak serum levels of active clindamycin are reached. Biologically inactive clindamycin phosphate disappears rapidly from the serum; the average elimination half-life is 6 minutes; however, the serum elimination half-life of active clindamycin is about 3 hours in adults and 2½ hours in pediatric patients.

After intramuscular injection of clindamycin phosphate, peak levels of active clindamycin are reached within 3 hours in adults and 1 hour in pediatric patients. Serum level curves may be constructed from IV peak serum levels as given in Table 1 by application of elimination half-lives listed above.

Serum levels of clindamycin can be maintained above the *in vitro* minimum inhibitory concentrations for most indicated organisms by administration of clindamycin phosphate every 8 to 12 hours in adults and every 6 to 8 hours in pediatric patients, or by continuous intravenous infusion. An equilibrium state is reached by the third dose.

The elimination half-life of clindamycin is increased slightly in patients with markedly reduced renal or hepatic function. Hemodialysis and peritoneal dialysis are not effective in removing clindamycin from the serum. Dosage schedules need not be modified in the presence of mild or moderate renal or hepatic disease.

No significant levels of clindamycin are attained in the cerebrospinal fluid even in the presence of inflamed meninges.

Pharmacokinetic studies in elderly volunteers (61–79 years) and younger adults (18–39 years) indicate that age alone does not alter clindamycin pharmacokinetics (clearance, elimination half-life, volume of distribution, and area under the serum concentration-time curve) after IV administration of clindamycin phosphate. After oral administration of clindamycin hydrochloride, elimination half-life is increased to approximately 4.0 hours (range 3.4–5.1 h) in the elderly compared to 3.2 hours (range 2.1– 4.2 h) in younger adults. The extent of absorption, however, is not different between age groups and no dosage alteration is necessary for the elderly with normal hepatic function and normal (age-adjusted) renal function¹.

Serum assays for active clindamycin require an inhibitor to prevent *in vitro* hydrolysis of clindamycin phosphate.

Table 1. Average Peak and Trough Serum Concentrations of Active Clindamycin After Dosing with Clindamycin Phosphate

Dosage Regimen	Peak mcg/mL	Trough mcg/mL
Healthy Adult Males (Post equilibrium)		
600 mg IV in 30 min q6h	10.9	2.0
600 mg IV in 30 min q8h	10.8	1.1
900 mg IV in 30 min q8h	14.1	1.7
600 mg IM q12h *	9	
Pediatric Patients (first dose) *		
5–7 mg/kg IV in 1 hour	10	
5–7 mg/kg IM	8	
3–5 mg/kg IM	4	

* Data in this group from patients being treated for infection.

Microbiology

Although clindamycin phosphate is inactive *in vitro*, rapid *in vivo* hydrolysis converts this compound to the antibacterially active clindamycin.

Clindamycin has been shown to have *in vitro* activity against isolates of the following organisms:

Aerobic gram positive cocci, including:

- Staphylococcus aureus (penicillinase and non-penicillinase producing strains). When tested by *in vitro* methods, some staphylococcal strains originally resistant to erythromycin rapidly develop resistance to clindamycin.
- Staphylococcus epidermidis (penicillinase and non-penicillinase producing strains). When tested by *in vitro* methods, some staphylococcal strains originally resistant to erythromycin rapidly develop resistance to clindamycin.
- Streptococci (except *Enterococcus faecalis*)
- Pneumococci

Anaerobic gram negative bacilli, including:

- *Bacteroides* species (including *Bacteroides fragilis* group and *Bacteroides melaninogenicus* group)
- *Fusobacterium* species

Anaerobic gram positive nonsporeforming bacilli, including:

- *Propionibacterium*
- *Eubacterium*
- *Actinomyces* species

Anaerobic and microaerophilic gram positive cocci, including:

- *Peptococcus* species
- *Peptostreptococcus* species
- Microaerophilic streptococci

Clostridia: Clostridia are more resistant than most anaerobes to clindamycin. Most *Clostridium perfringens* are susceptible, but other species, e.g., *Clostridium sporogenes* and *Clostridium tertium* are frequently resistant to clindamycin. Susceptibility testing should be done.

Cross resistance has been demonstrated between clindamycin and lincomycin.

Antagonism has been demonstrated between clindamycin and erythromycin.

In vitro Susceptibility Testing

Disk diffusion technique

Quantitative methods that require measurement of zone diameters give the most precise estimates of antibiotic susceptibility. One such procedure² has been recommended for use with disks to test susceptibility to clindamycin.

Reports from a laboratory using the standardized single-disk susceptibility test¹ with a 2 mcg clindamycin disk should be interpreted according to the following criteria:

Susceptible organisms produce zones of 17 mm or greater, indicating that the tested organism is likely to respond to therapy.

Organisms of intermediate susceptibility produce zones of 15–16 mm, indicating that the tested organism would be susceptible if a high dosage is used or if the infection is confined to tissues and fluids (e.g., urine), in which high antibiotic levels are attained.

Resistant organisms produce zones of 14 mm or less, indicating that other therapy should be selected.

Standardized procedures require the use of control organisms. The 2 mcg clindamycin disk should give a zone diameter between 24 and 30 mm for *S. aureus* ATCC 25923.

Dilution techniques

A bacterial isolate may be considered susceptible if the minimum inhibitory concentration (MIC) for clindamycin is not more than 1.6 mcg/mL. Organisms are considered moderately susceptible if the MIC is greater than 1.6 mcg/mL and less than or equal to 4.8 mcg/mL. Organisms are considered resistant if the MIC is greater than 4.8 mcg per mL.

The range of MICs for the control strains are as follows:

S. aureus ATCC 29213, 0.06—0.25 mcg/mL.

E. faecalis ATCC 29212, 4.0—16 mcg/mL.

For anaerobic bacteria the minimum inhibitory concentration (MIC) of clindamycin can be determined by agar dilution and broth dilution (including microdilution) techniques.³ If MICs are not determined routinely, the disk broth method is recommended for routine use. THE KIRBY-BAUER DISK DIFFUSION METHOD AND ITS INTERPRETIVE STANDARDS ARE NOT RECOMMENDED FOR ANAEROBES.

INDICATIONS AND USAGE

CLEOCIN PHOSPHATE products are indicated in the treatment of serious infections caused by susceptible anaerobic bacteria.

CLEOCIN PHOSPHATE products are also indicated in the treatment of serious infections due to susceptible strains of streptococci, pneumococci, and staphylococci. Its use should be reserved for penicillin-allergic patients or other patients for whom, in the judgment of the physician, a penicillin is inappropriate. Because of the risk of antibiotic-associated pseudomembranous colitis, as described in the WARNING box, before selecting clindamycin the physician should consider the nature of the infection and the suitability of less toxic alternatives (e.g., erythromycin).

Bacteriologic studies should be performed to determine the causative organisms and their susceptibility to clindamycin.

Indicated surgical procedures should be performed in conjunction with antibiotic therapy.

CLEOCIN PHOSPHATE is indicated in the treatment of serious infections caused by susceptible strains of the designated organisms in the conditions listed below:

Lower respiratory tract infections including pneumonia, empyema, and lung abscess caused by anaerobes, *Streptococcus pneumoniae*, other streptococci (except *E. faecalis*), and *Staphylococcus aureus*.

Skin and skin structure infections caused by *Streptococcus pyogenes*, *Staphylococcus aureus*, and anaerobes.

Gynecological infections including endometritis, non-gonococcal tubo-ovarian abscess, pelvic cellulitis, and postsurgical vaginal cuff infection caused by susceptible anaerobes.

Intra-abdominal infections including peritonitis and intra-abdominal abscess caused by susceptible anaerobic organisms.

Septicemia caused by *Staphylococcus aureus*, streptococci (except *Enterococcus faecalis*), and susceptible anaerobes.

Bone and joint infections including acute hematogenous osteomyelitis caused by *Staphylococcus aureus* and as adjunctive therapy in the surgical treatment of chronic bone and joint infections due to susceptible organisms.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of CLEOCIN PHOSPHATE and other antibacterial drugs, CLEOCIN PHOSPHATE should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

CONTRAINDICATIONS

This drug is contraindicated in individuals with a history of hypersensitivity to preparations containing clindamycin or lincomycin.

WARNINGS

See **WARNING** box.

Clostridium difficile associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including CLEOCIN PHOSPHATE, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

A careful inquiry should be made concerning previous sensitivities to drugs and other allergens.

This product contains benzyl alcohol as a preservative. Benzyl alcohol has been associated with a fatal "Gasping Syndrome" in premature infants. (See **PRECAUTIONS—Pediatric Use**.)

Usage in Meningitis—Since clindamycin does not diffuse adequately into the cerebrospinal fluid, the drug should not be used in the treatment of meningitis.

SERIOUS ANAPHYLACTOID REACTIONS REQUIRE IMMEDIATE EMERGENCY TREATMENT WITH EPINEPHRINE. OXYGEN AND INTRAVENOUS CORTICOSTEROIDS SHOULD ALSO BE ADMINISTERED AS INDICATED.

PRECAUTIONS

General

Review of experience to date suggests that a subgroup of older patients with associated severe illness may tolerate diarrhea less well. When clindamycin is indicated in these patients, they should be carefully monitored for change in bowel frequency.

CLEOCIN PHOSPHATE products should be prescribed with caution in individuals with a history of gastrointestinal disease, particularly colitis.

CLEOCIN PHOSPHATE should be prescribed with caution in atopic individuals.

Certain infections may require incision and drainage or other indicated surgical procedures in addition to antibiotic therapy.

The use of CLEOCIN PHOSPHATE may result in overgrowth of nonsusceptible organisms—particularly yeasts. Should superinfections occur, appropriate measures should be taken as indicated by the clinical situation.

CLEOCIN PHOSPHATE should not be injected intravenously undiluted as a bolus, but should be infused over at least 10–60 minutes as directed in the DOSAGE AND ADMINISTRATION section.

Clindamycin dosage modification may not be necessary in patients with renal disease. In patients with moderate to severe liver disease, prolongation of clindamycin half-life has been found. However, it was postulated from studies that when given every eight hours, accumulation should rarely occur. Therefore, dosage modification in patients with liver disease may not be necessary. However, periodic liver enzyme determinations should be made when treating patients with severe liver disease.

Prescribing CLEOCIN PHOSPHATE in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

Information for Patients

Patients should be counseled that antibacterial drugs including CLEOCIN PHOSPHATE should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When CLEOCIN PHOSPHATE is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by CLEOCIN PHOSPHATE or other antibacterial drugs in the future.

Diarrhea is a common problem caused by antibiotics which usually ends when the antibiotic is discontinued. Sometimes after starting treatment with antibiotics, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibiotic. If this occurs, patients should contact their physician as soon as possible.

Laboratory Tests

During prolonged therapy periodic liver and kidney function tests and blood counts should be performed.

Drug Interactions

Clindamycin has been shown to have neuromuscular blocking properties that may enhance the action of other neuromuscular blocking agents. Therefore, it should be used with caution in patients receiving such agents.

Antagonism has been demonstrated between clindamycin and erythromycin *in vitro*. Because of possible clinical significance, the two drugs should not be administered concurrently.

Carcinogenesis, Mutagenesis, Impairment Of Fertility

Long term studies in animals have not been performed with clindamycin to evaluate carcinogenic potential. Genotoxicity tests performed included a rat micronucleus test and an Ames Salmonella reversion test. Both tests were negative.

Fertility studies in rats treated orally with up to 300 mg/kg/day (approximately 1.1 times the highest recommended adult human dose based on mg/m²) revealed no effects on fertility or mating ability.

Pregnancy

Teratogenic effects

Pregnancy category B

Reproduction studies performed in rats and mice using oral doses of clindamycin up to 600 mg/kg/day (2.1 and 1.1 times the highest recommended adult human dose based on mg/m², respectively) or subcutaneous doses of clindamycin up to 250 mg/kg/day (0.9 and 0.5 times the highest recommended adult human dose based on mg/m², respectively) revealed no evidence of teratogenicity.

There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of the human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

Clindamycin has been reported to appear in breast milk in the range of 0.7 to 3.8 mcg/mL at dosages of 150 mg orally to 600 mg intravenously. Because of the potential for adverse reactions due to clindamycin in neonates (see **Pediatric Use**), the decision to discontinue the drug should be made, taking into account the importance of the drug to the mother.

Pediatric Use

When CLEOCIN PHOSPHATE Sterile Solution is administered to the pediatric population (birth to 16 years) appropriate monitoring of organ system functions is desirable.

Usage in Newborns and Infants

This product contains benzyl alcohol as a preservative. Benzyl alcohol has been associated with a fatal "Gasping Syndrome" in premature infants.

The potential for the toxic effect in the pediatric population from chemicals that may leach from the single dose premixed IV preparation in plastic has not been evaluated.

Geriatric Use

Clinical studies of clindamycin did not include sufficient numbers of patients age 65 and over to determine whether they respond differently from younger patients. However, other reported clinical experience indicates that antibiotic-associated colitis and diarrhea (due to *Clostridium difficile*) seen in association with most antibiotics occur more frequently in the elderly (>60 years) and may be more severe. These patients should be carefully monitored for the development of diarrhea.

Pharmacokinetic studies with clindamycin have shown no clinically important differences between young and elderly subjects with normal hepatic function and normal (age-adjusted) renal function after oral or intravenous administration.

ADVERSE REACTIONS

The following reactions have been reported with the use of clindamycin.

Gastrointestinal

Antibiotic-associated colitis (see **WARNINGS**), pseudomembranous colitis, abdominal pain, nausea, and vomiting. The onset of pseudomembranous colitis symptoms may occur during or after antibacterial treatment (see **WARNINGS**). An unpleasant or metallic taste occasionally has been reported after

intravenous administration of the higher doses of clindamycin phosphate.

Hypersensitivity Reactions

Maculopapular rash and urticaria have been observed during drug therapy. Generalized mild to moderate morbilliform-like skin rashes are the most frequently reported of all adverse reactions. Rare instances of erythema multiforme, some resembling Stevens-Johnson syndrome, have been associated with clindamycin. A few cases of anaphylactoid reactions have been reported. If a hypersensitivity reaction occurs, the drug should be discontinued. The usual agents (epinephrine, corticosteroids, antihistamines) should be available for emergency treatment of serious reactions.

Skin and Mucous Membranes

Pruritus, vaginitis, and rare instances of exfoliative dermatitis have been reported (see *Hypersensitivity Reactions*).

Liver

Jaundice and abnormalities in liver function tests have been observed during clindamycin therapy.

Renal

Although no direct relationship of clindamycin to renal damage has been established, renal dysfunction as evidenced by azotemia, oliguria, and/or proteinuria has been observed in rare instances.

Hematopoietic

Transient neutropenia (leukopenia) and eosinophilia have been reported. Reports of agranulocytosis and thrombocytopenia have been made. No direct etiologic relationship to concurrent clindamycin therapy could be made in any of the foregoing.

Local Reactions

Pain, induration and sterile abscess have been reported after intramuscular injection and thrombophlebitis after intravenous infusion. Reactions can be minimized or avoided by giving deep intramuscular injections and avoiding prolonged use of indwelling intravenous catheters.

Musculoskeletal

Rare instances of polyarthrititis have been reported.

Cardiovascular

Rare instances of cardiopulmonary arrest and hypotension have been reported following too rapid intravenous administration. (See **DOSAGE AND ADMINISTRATION** section.)

OVERDOSAGE

Significant mortality was observed in mice at an intravenous dose of 855 mg/kg and in rats at an oral or subcutaneous dose of approximately 2618 mg/kg. In the mice, convulsions and depression were observed.

Hemodialysis and peritoneal dialysis are not effective in removing clindamycin from the serum.

DOSAGE AND ADMINISTRATION

If diarrhea occurs during therapy, this antibiotic should be discontinued (see **WARNING** box).

Adults: Parenteral (IM or IV Administration): Serious infections due to aerobic gram-positive cocci and the more susceptible anaerobes (NOT generally including *Bacteroides fragilis*, *Peptococcus* species and *Clostridium* species other than *Clostridium perfringens*):

600–1200 mg/day in 2, 3 or 4 equal doses.

More severe infections, particularly those due to proven or suspected *Bacteroides fragilis*, *Peptococcus* species, or *Clostridium* species other than *Clostridium perfringens*:

1200–2700 mg/day in 2, 3 or 4 equal doses.

For more serious infections, these doses may have to be increased. In life-threatening situations due to either aerobes or anaerobes these doses may be increased. Doses of as much as 4800 mg daily have been given intra-venously to adults. See **Dilution and Infusion Rates** section below.

Single intramuscular injections of greater than 600 mg are not recommended.

Alternatively, drug may be administered in the form of a single rapid infusion of the first dose followed by continuous IV infusion as follows:

To maintain serum clindamycin levels	Rapid infusion rate	Maintenance infusion rate
Above 4 mcg/mL	10 mg/min for 30 min	0.75 mg/min
Above 5 mcg/mL	15 mg/min for 30 min	1.00 mg/min
Above 6 mcg/mL	20 mg/min for 30 min	1.25 mg/min

Neonates (less than 1 month):

15 to 20 mg/kg/day in 3 to 4 equal doses. The lower dosage may be adequate for small prematures.

Pediatric patients 1 month of age to 16 years: Parenteral (IM or IV) administration: 20 to 40 mg/kg/day in 3 or 4 equal doses. The higher doses would be used for more severe infections. As an alternative to dosing on a body weight basis, pediatric patients may be dosed on the basis of square meters body surface: 350 mg/m²/day for serious infections and 450 mg/m²/day for more severe infections.

Parenteral therapy may be changed to oral CLEOCIN PEDIATRIC® Flavored Granules (clindamycin palmitate hydrochloride) or CLEOCIN HCl® Capsules (clindamycin hydrochloride) when the condition warrants and at the discretion of the physician.

In cases of β -hemolytic streptococcal infections, treatment should be continued for at least 10 days.

Dilution and Infusion Rates

Clindamycin phosphate must be diluted prior to IV administration. The concentration of clindamycin in diluent for infusion should not exceed 18 mg per mL. Infusion rates should not exceed 30 mg per minute. The usual infusion dilutions and rates are as follows:

Dose	Diluent	Time
300 mg	50 mL	10 min
600 mg	50 mL	20 min
900 mg	50–100 mL	30 min
1200 mg	100 mL	40 min

Administration of more than 1200 mg in a single 1-hour infusion is not recommended.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Dilution and Compatibility: Physical and biological compatibility studies monitored for 24 hours at room temperature have demonstrated no inactivation or incompatibility with the use of CLEOCIN PHOSPHATE Sterile Solution (clindamycin phosphate) in IV solutions containing sodium chloride, glucose, calcium or potassium, and solutions containing vitamin B complex in concentrations usually used clinically. No incompatibility has been demonstrated with the antibiotics cephalothin, kanamycin, gentamicin, penicillin or carbenicillin.

The following drugs are physically incompatible with clindamycin phosphate: ampicillin sodium, phenytoin sodium, barbiturates, aminophylline, calcium gluconate, and magnesium sulfate.

The compatibility and duration of stability of drug admixtures will vary depending on concentration and other conditions. For current information regarding compatibilities of clindamycin phosphate under specific conditions, please contact the Medical and Drug Information Unit, Pharmacia & Upjohn Company (Division of Pfizer Inc).

Physico-Chemical Stability of diluted solutions of CLEOCIN PHOSPHATE

Room temperature: 6, 9 and 12 mg/mL (equivalent to clindamycin base) in dextrose injection 5%, sodium chloride injection 0.9%, or Lactated Ringers Injection in glass bottles or minibags, demonstrated physical and chemical stability for at least 16 days at 25°C. Also, 18 mg/mL (equivalent to clindamycin base) in dextrose injection 5%, in minibags, demonstrated physical and chemical stability for at least 16 days at 25°C.

Refrigeration: 6, 9 and 12 mg/mL (equivalent to clindamycin base) in dextrose injection 5%, sodium chloride injection 0.9%, or Lactated Ringers Injection in glass bottles or minibags, demonstrated physical and chemical stability for at least 32 days at 4°C.

IMPORTANT: This chemical stability information in no way indicates that it would be acceptable practice to use this product well after the preparation time. Good professional practice suggests that compounded admixtures should be administered as soon after preparation as is feasible.

Frozen: 6, 9 and 12 mg/mL (equivalent to clindamycin base) in dextrose injection 5%, sodium chloride injection 0.9%, or Lactated Ringers Injection in minibags demonstrated physical and chemical stability for at least eight weeks at -10°C.

Frozen solutions should be thawed at room temperature and not refrozen.

DIRECTIONS FOR DISPENSING

Pharmacy Bulk Package — Not for Direct Infusion

The Pharmacy Bulk Package is for use in a Pharmacy Admixture Service only under a laminar flow hood. Entry into the vial should be made with a small diameter sterile transfer set or other small diameter sterile dispensing device, and contents dispensed in aliquots using aseptic technique. Multiple entries with a needle and syringe are not recommended. AFTER ENTRY USE ENTIRE CONTENTS OF VIAL PROMPTLY. ANY UNUSED PORTION MUST BE DISCARDED WITHIN 24 HOURS AFTER INITIAL ENTRY.

DIRECTIONS FOR USE

CLEOCIN PHOSPHATE IV Solution in Galaxy Plastic Container

Premixed CLEOCIN PHOSPHATE IV Solution is for intravenous administration using sterile equipment. Check for minute leaks prior to use by squeezing bag firmly. If leaks are found, discard solution as sterility may be impaired. Do not add supplementary medication. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit. Do not use unless solution is clear and seal is intact.

Caution: Do not use plastic containers in series connections. Such use could result in air embolism due to residual air being drawn from the primary container before administration of the fluid from the secondary container is complete.

Preparation for Administration:

1. Suspend container from eyelet support.
2. Remove protector from outlet port at bottom of container.
3. Attach administration set. Refer to complete directions accompanying set.

Preparation of CLEOCIN PHOSPHATE in ADD-Vantage System—For IV Use Only. CLEOCIN PHOSPHATE 600 mg and 900 mg may be reconstituted in 50 mL or 100 mL, respectively, of Dextrose Injection 5% or Sodium Chloride Injection 0.9% in the ADD-diluent container. Refer to separate instructions for ADD-Vantage¹ System.

¹ ADD-Vantage is a registered trademark of Abbott Laboratories.

HOW SUPPLIED

Each mL of CLEOCIN PHOSPHATE Sterile Solution contains clindamycin phosphate equivalent to 150 mg clindamycin; 0.5 mg disodium edetate; 9.45 mg benzyl alcohol added as preservative. When necessary, pH is adjusted with sodium hydroxide and/or hydrochloric acid. CLEOCIN PHOSPHATE is available in the following packages:

25-2 mL vials	NDC 0009-0870-26
25-4 mL vials	NDC 0009-0775-26
25-6 mL vials	NDC 0009-0902-18
1-60 mL Pharmacy Bulk Package	NDC 0009-0728-05

CLEOCIN PHOSPHATE is supplied in ADD-Vantage vials as follows:

NDC	Vial Size	Total Clindamycin Phosphate/vial	Amount of Diluent
0009-3124-03	25-4 mL vials	600 mg	50 mL
0009-3447-03	25-6 mL vials	900 mg	100 mL

Store at controlled room temperature 20° to 25°C (68° to 77°F) [see USP]. CLEOCIN PHOSPHATE IV Solution in Galaxy plastic containers is a sterile solution of clindamycin phosphate with 5% dextrose. The single dose Galaxy plastic containers are available as follows:

24-300 mg/50 mL containers	NDC 0009-3381-02
----------------------------	------------------

Cleocin Phosphate®(clindamycin injection, USP) and(clindamycin injection in 5% dex... Page 13 of 22

24-600 mg/50 mL containers	NDC 0009-3375-02
24-900 mg/50 mL containers	NDC 0009-3382-02

Exposure of pharmaceutical products to heat should be minimized. It is recommended that Galaxy plastic containers be stored at room temperature (25°C). Avoid temperatures above 30°C.

ANIMAL TOXICOLOGY

One year oral toxicity studies in Spartan Sprague-Dawley rats and beagle dogs at dose levels up to 300 mg/kg/day (approximately 1.1 and 3.6 times the highest recommended adult human dose based on mg/m², respectively) have shown clindamycin to be well tolerated. No appreciable difference in pathological findings has been observed between groups of animals treated with clindamycin and comparable control groups. Rats receiving clindamycin hydrochloride at 600 mg/kg/day (approximately 2.1 times the highest recommended adult human dose based on mg/m²) for 6 months tolerated the drug well; however, dogs dosed at this level (approximately 7.2 times the highest recommended adult human dose based on mg/m²) vomited, would not eat, and lost weight.

REFERENCES

1. Smith RB, Phillips JP: Evaluation of CLEOCIN HCl and CLEOCIN Phosphate in an Aged Population. Upjohn TR 8147-82-9122-021, December 1982.
2. Bauer AW, Kirby WMM, Sherris JC, Turck M; Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Path.*, **45**:493–496, 1966. Standardized Disk Susceptibility Test, *Federal Register*, **37**:20527–29, 1972.
3. National Committee for Clinical Lab. Standards. Methods for Antimicrobial Susceptibility Testing of Anaerobic Bacteria—Second Edition; Tentative Standard. NCCLS publication M₁₁-T₂. Villanova, PA; NCCLS; 1988.

CLEOCIN PHOSPHATE IV Solution in the Galaxy plastic containers is manufactured for Pfizer Inc by Baxter Healthcare Corporation, Deerfield, IL 60015.

Galaxy® is a trademark of Baxter International, Inc.

Rx Only



Distributed by

Pharmacia & Upjohn Company
Division of Pfizer Inc, NY, NY 10017

LAB-0044-5.0

June 2007

Cleocin Phosphate®
clindamycin sterile solution for injection, USP
in ADD-Vantage™ Vial
For Intravenous Use Only
NOT FOR INTRAMUSCULAR USE

INSTRUCTIONS FOR USE FOR ADD-VANTAGE SYSTEM—FOR IV USE ONLY

To Open Diluent Container:

Peel overwrap from the corner and remove container. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually.

To Assemble Vial and Flexible Diluent Container:

(Use Aseptic Technique)

1. Remove the protective covers from the top of the vial and the vial port on the diluent container as follows:
 - a. To remove the breakaway vial cap, swing the pull ring over the top of the vial and pull down far enough to start the opening (SEE FIGURE 1.) then pull straight up to remove the cap. (SEE FIGURE 2.) **NOTE:** Once the breakaway cap has been removed, **DO NOT ACCESS VIAL WITH SYRINGE.**




Fig. 1



Fig. 2

- b. To remove the vial port cover, grasp the tab on the pull ring, pull up to break the three tie strings, then pull back to remove the cover. (SEE FIGURE 3.)
2. Screw the vial into the vial port until it will go no further. **THE VIAL MUST BE SCREWED IN TIGHTLY TO ASSURE A SEAL.** This occurs approximately $\frac{1}{2}$ turn (180°) after the first audible click. (SEE FIGURE 4.) The clicking sound does not assure a seal; the vial must be turned as far as it will go. **NOTE:** Once vial is seated, do not attempt to remove. (SEE FIGURE 4.)
3. Recheck the vial to assure that it is tight by trying to turn it further in the direction of assembly.
4. Label appropriately.



Fig. 3




Fig. 4

To Prepare Admixture:

1. Squeeze the bottom of the diluent container gently to inflate the portion of the container surrounding the end of the drug vial.
2. With the other hand, push the drug vial down into the container telescoping the walls of the container. Grasp the inner cap of the vial through the walls of the container. (SEE FIGURE 5.)
3. Pull the inner cap from the drug vial. (SEE FIGURE 6.) Verify that the rubber stopper has been pulled out, allowing the drug and diluent to mix.




Fig. 5




Fig. 6

4. Mix container contents thoroughly and use within the specified time.

Preparation for Administration:(Use Aseptic Technique)

1. Confirm the activation and admixture of vial contents.
2. Check for leaks by squeezing container firmly. If leaks are found, discard unit as sterility may be impaired.
3. Close flow control clamp of administration set.

4. Remove cover from outlet port at bottom of container.
5. Insert piercing pin of administration set into port with a twisting motion until the pin is firmly seated. NOTE: See full directions on administration set carton.
6. Lift the free end of the hanger loop on the bottom of the vial, breaking the two tie strings. Bend the loop outward to lock it in the upright position, then suspend container from hanger.
7. Squeeze and release drip chamber to establish proper fluid level in chamber.
8. Open flow control clamp and clear air from set. Close clamp.
9. Attach set to venipuncture device. If device is not indwelling, prime and make venipuncture.
10. Regulate rate of administration with flow control clamp.

WARNING: Do not use flexible container in series connections.



Distributed by

Pharmacia & Upjohn Company
Division of Pfizer Inc, NY, NY 10017

™ADD-Vantage is a registered trademark of Abbott Laboratories

LAB-0050-2.0

November 2005

Cleocin Phosphate (clindamycin phosphate)			
PRODUCT INFO			
Product Code	0009-0870	Dosage Form	INJECTION, SOLUTION
Route Of Administration	INTRAMUSCULAR	DEA Schedule	
INGREDIENTS			
Name (Active Moiety)	Type	Strength	
clindamycin phosphate (clindamycin)	Active	150 MILLIGRAM In 1 MILLILITER	
disodium edetate	Inactive		
benzyl alcohol	Inactive		
dextrose	Inactive		
sodium hydroxide	Inactive		
hydrochloric acid	Inactive		
IMPRINT INFORMATION			
Characteristic Appearance	Characteristic	Appearance	
Color	Score		
Shape	Symbol		
Imprint Code	Coating		

Size		
PACKAGING		
# NDC	Package Description	Multilevel Packaging
1 0009-0870-26	25 VIAL In 1 BAG	contains a VIAL
1	2 MILLILITER In 1 VIAL	This package is contained within the BAG (0009-0870-26)

Cleocin Phosphate (clindamycin phosphate)			
PRODUCT INFO			
Product Code	0009-0775	Dosage Form	INJECTION, SOLUTION
Route Of Administration	INTRAMUSCULAR	DEA Schedule	
INGREDIENTS			
Name (Active Moiety)	Type	Strength	
clindamycin phosphate (clindamycin)	Active	150 MILLIGRAM In 1 MILLILITER	
disodium edetate	Inactive		
benzyl alcohol	Inactive		
dextrose	Inactive		
sodium hydroxide	Inactive		
hydrochloric acid	Inactive		
IMPRINT INFORMATION			
Characteristic Appearance	Characteristic	Appearance	
Color	Score		
Shape	Symbol		
Imprint Code	Coating		
Size			
PACKAGING			
# NDC	Package Description	Multilevel Packaging	
1 0009-0775-26	25 VIAL In 1 BAG	contains a VIAL	
1	4 MILLILITER In 1 VIAL	This package is contained within the BAG (0009-0775-26)	

Cleocin Phosphate (clindamycin phosphate)
PRODUCT INFO

Product Code	0009-0902	Dosage Form	INJECTION, SOLUTION
Route Of Administration	INTRAMUSCULAR	DEA Schedule	

INGREDIENTS

Name (Active Moiety)	Type	Strength
clindamycin phosphate (clindamycin)	Active	150 MILLIGRAM In 1 MILLILITER
disodium edetate	Inactive	
benzyl alcohol	Inactive	
dextrose	Inactive	
sodium hydroxide	Inactive	
hydrochloric acid	Inactive	

IMPRINT INFORMATION

Characteristic Appearance	Characteristic	Appearance
Color	Score	
Shape	Symbol	
Imprint Code	Coating	
Size		

PACKAGING

#	NDC	Package Description	Multilevel Packaging
1	0009-0902-18	25 VIAL In 1 BAG	contains a VIAL
1		6 MILLILITER In 1 VIAL	This package is contained within the BAG (0009-0902-18)

Cleocin Phosphate (clindamycin phosphate)

PRODUCT INFO			
Product Code	0009-0728	Dosage Form	INJECTION, SOLUTION
Route Of Administration	INTRAMUSCULAR	DEA Schedule	

INGREDIENTS

Name (Active Moiety)	Type	Strength
clindamycin phosphate (clindamycin)	Active	150 MILLIGRAM In 1 MILLILITER
disodium edetate	Inactive	
benzyl alcohol	Inactive	
dextrose	Inactive	
sodium hydroxide	Inactive	
hydrochloric acid	Inactive	

IMPRINT INFORMATION

Characteristic	Appearance	Characteristic	Appearance
Color		Score	
Shape		Symbol	
Imprint Code		Coating	
Size			

PACKAGING

# NDC	Package Description	Multilevel Packaging
1 0009-0728-05	60 MILLILITER In 1 VIAL, PHARMACY BULK PACKAGE	None

Cleocin Phosphate (clindamycin phosphate)

PRODUCT INFO

Product Code	0009-3124	Dosage Form	INJECTION, SOLUTION
Route Of Administration	INTRAMUSCULAR	DEA Schedule	

INGREDIENTS

Name (Active Moiety)	Type	Strength
clindamycin phosphate (clindamycin)	Active	150 MILLIGRAM In 1 MILLILITER
disodium edetate	Inactive	
benzyl alcohol	Inactive	
dextrose	Inactive	
sodium hydroxide	Inactive	
hydrochloric acid	Inactive	

IMPRINT INFORMATION

Characteristic	Appearance	Characteristic	Appearance
Color		Score	
Shape		Symbol	
Imprint Code		Coating	
Size			

PACKAGING

# NDC	Package Description	Multilevel Packaging
1 0009-3124-03	25 VIAL In 1 PACKAGE	contains a VIAL
1	4 MILLILITER In 1 VIAL	This package is contained within the PACKAGE (0009-3124-03)

Cleocin Phosphate (clindamycin phosphate)

PRODUCT INFO

Product Code	0009-3447	Dosage Form	INJECTION, SOLUTION
Route Of Administration	INTRAMUSCULAR	DEA Schedule	

INGREDIENTS

Name (Active Moiety)	Type	Strength
clindamycin phosphate (clindamycin)	Active	150 MILLIGRAM In 1 MILLILITER
disodium edetate	Inactive	
benzyl alcohol	Inactive	
dextrose	Inactive	
sodium hydroxide	Inactive	
hydrochloric acid	Inactive	

IMPRINT INFORMATION

Characteristic Appearance	Characteristic	Appearance
Color	Score	
Shape	Symbol	
Imprint Code	Coating	
Size		

PACKAGING

# NDC	Package Description	Multilevel Packaging
1 0009-3447-03	25 VIAL In 1 PACKAGE	contains a VIAL
1	6 MILLILITER In 1 VIAL	This package is contained within the PACKAGE (0009-3447-03)

Cleocin Phosphate (clindamycin phosphate)

PRODUCT INFO

Product Code	0009-3381	Dosage Form	INJECTION, SOLUTION
Route Of Administration	INTRAVENOUS	DEA Schedule	

INGREDIENTS

Name (Active Moiety)	Type	Strength
clindamycin phosphate (clindamycin)	Active	150 MILLIGRAM In 1 MILLILITER
disodium edetate	Inactive	
benzyl alcohol	Inactive	
dextrose	Inactive	

sodium hydroxide	Inactive	
hydrochloric acid	Inactive	

IMPRINT INFORMATION

Characteristic	Appearance	Characteristic	Appearance
Color		Score	
Shape		Symbol	
Imprint Code		Coating	
Size			

PACKAGING

#	NDC	Package Description	Multilevel Packaging
1	0009-3381-02	24 CONTAINER In 1 PACKAGE	contains a VIAL, PLASTIC
1		50 MILLILITER In 1 VIAL, PLASTIC	This package is contained within the PACKAGE (0009-3381-02)

Cleocin Phosphate (clindamycin phosphate)																																																												
PRODUCT INFO <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%;">Product Code</td> <td style="width: 20%;">0009-3375</td> <td style="width: 20%;">Dosage Form</td> <td style="width: 30%;">INJECTION, SOLUTION</td> </tr> <tr> <td>Route Of Administration</td> <td>INTRAVENOUS</td> <td>DEA Schedule</td> <td></td> </tr> </table> INGREDIENTS <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 45%;">Name (Active Moiety)</th> <th style="width: 15%;">Type</th> <th style="width: 40%;">Strength</th> </tr> </thead> <tbody> <tr> <td>clindamycin phosphate (clindamycin)</td> <td>Active</td> <td>150 MILLIGRAM In 1 MILLILITER</td> </tr> <tr> <td>disodium edetate</td> <td>Inactive</td> <td></td> </tr> <tr> <td>benzyl alcohol</td> <td>Inactive</td> <td></td> </tr> <tr> <td>dextrose</td> <td>Inactive</td> <td></td> </tr> <tr> <td>sodium hydroxide</td> <td>Inactive</td> <td></td> </tr> <tr> <td>hydrochloric acid</td> <td>Inactive</td> <td></td> </tr> </tbody> </table> IMPRINT INFORMATION <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 40%;">Characteristic</th> <th style="width: 20%;">Appearance</th> <th style="width: 40%;">Characteristic</th> <th style="width: 20%;">Appearance</th> </tr> </thead> <tbody> <tr> <td>Color</td> <td></td> <td>Score</td> <td></td> </tr> <tr> <td>Shape</td> <td></td> <td>Symbol</td> <td></td> </tr> <tr> <td>Imprint Code</td> <td></td> <td>Coating</td> <td></td> </tr> <tr> <td>Size</td> <td></td> <td></td> <td></td> </tr> </tbody> </table> PACKAGING <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 10%;">#</th> <th style="width: 15%;">NDC</th> <th style="width: 40%;">Package Description</th> <th style="width: 35%;">Multilevel Packaging</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>				Product Code	0009-3375	Dosage Form	INJECTION, SOLUTION	Route Of Administration	INTRAVENOUS	DEA Schedule		Name (Active Moiety)	Type	Strength	clindamycin phosphate (clindamycin)	Active	150 MILLIGRAM In 1 MILLILITER	disodium edetate	Inactive		benzyl alcohol	Inactive		dextrose	Inactive		sodium hydroxide	Inactive		hydrochloric acid	Inactive		Characteristic	Appearance	Characteristic	Appearance	Color		Score		Shape		Symbol		Imprint Code		Coating		Size				#	NDC	Package Description	Multilevel Packaging				
Product Code	0009-3375	Dosage Form	INJECTION, SOLUTION																																																									
Route Of Administration	INTRAVENOUS	DEA Schedule																																																										
Name (Active Moiety)	Type	Strength																																																										
clindamycin phosphate (clindamycin)	Active	150 MILLIGRAM In 1 MILLILITER																																																										
disodium edetate	Inactive																																																											
benzyl alcohol	Inactive																																																											
dextrose	Inactive																																																											
sodium hydroxide	Inactive																																																											
hydrochloric acid	Inactive																																																											
Characteristic	Appearance	Characteristic	Appearance																																																									
Color		Score																																																										
Shape		Symbol																																																										
Imprint Code		Coating																																																										
Size																																																												
#	NDC	Package Description	Multilevel Packaging																																																									

1	0009-3375-02	24 CONTAINER In 1 PACKAGE	contains a VIAL, PLASTIC
1		50 MILLILITER In 1 VIAL, PLASTIC	This package is contained within the PACKAGE (0009-3375-02)

Cleocin Phosphate (clindamycin phosphate)

PRODUCT INFO

Product Code	0009-3382	Dosage Form	INJECTION, SOLUTION
Route Of Administration	INTRAVENOUS	DEA Schedule	

INGREDIENTS

Name (Active Moiety)	Type	Strength
clindamycin phosphate (clindamycin)	Active	150 MILLIGRAM In 1 MILLILITER
disodium edetate	Inactive	
benzyl alcohol	Inactive	
dextrose	Inactive	
sodium hydroxide	Inactive	
hydrochloric acid	Inactive	

IMPRINT INFORMATION

Characteristic	Appearance	Characteristic	Appearance
Color		Score	
Shape		Symbol	
Imprint Code		Coating	
Size			

PACKAGING

#	NDC	Package Description	Multilevel Packaging
1	0009-3382-02	1 CONTAINER In 1 PACKAGE	contains a VIAL, PLASTIC
1		50 MILLILITER In 1 VIAL, PLASTIC	This package is contained within the PACKAGE (0009-3382-02)

Revised: 03/2007

Pharmacia & Upjohn, Co.