

**This label may not be the latest approved by FDA.
Labeling information, please visit <https://www.fda.gov>**

**Cetirizine Hydrochloride
and Pseudoephedrine
Hydrochloride**

**Extended Release Tablets
5 mg / 120 mg**

TO OPEN USE SCISSORS

DISTRIBUTED BY PERRIGO
ALLEGAN, MI 49010 USA

MADE IN INDIA

LOT EXP.

**Cetirizine Hydrochloride
and Pseudoephedrine
Hydrochloride**

**Extended Release Tablets
5 mg / 120 mg**

TO OPEN USE SCISSORS

DISTRIBUTED BY PERRIGO
ALLEGAN, MI 49010 USA

MADE IN INDIA

LOT EXP.
17600 00 X1

**Cetirizine Hydrochloride
and Pseudoephedrine
Hydrochloride**

**Extended Release Tablets
5 mg / 120 mg**

TO OPEN USE SCISSORS

DISTRIBUTED BY PERRIGO
ALLEGAN, MI 49010 USA

MADE IN INDIA

LOT EXP.

**Cetirizine Hydrochloride
and Pseudoephedrine
Hydrochloride**

**Extended Release Tablets
5 mg / 120 mg**

TO OPEN USE SCISSORS

DISTRIBUTED BY PERRIGO
ALLEGAN, MI 49010 USA

MADE IN INDIA

LOT EXP.

**Cetirizine Hydrochloride
and Pseudoephedrine
Hydrochloride**

**Extended Release Tablets
5 mg / 120 mg**

TO OPEN USE SCISSORS

DISTRIBUTED BY PERRIGO
ALLEGAN, MI 49010 USA

MADE IN INDIA

LOT EXP.

**Cetirizine Hydrochloride
and Pseudoephedrine
Hydrochloride**

**Extended Release Tablets
5 mg / 120 mg**

TO OPEN USE SCISSORS

DISTRIBUTED BY PERRIGO
ALLEGAN, MI 49010 USA

MADE IN INDIA

LOT EXP.

EFB1021
FA17653C1

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease ■ thyroid disease ■ diabetes
- glaucoma ■ high blood pressure
- trouble urinating due to an enlarged prostate gland or liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

- **do not use more than directed**
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask doctor if

- an allergic reaction to this product occurs. Seek medical help right away.

Other information

- store between 20° to 25°C (68° to 77°F)

Inactive ingredients

coloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

Directions

- do not break or chew tablet; swallow tablet whole

adults and children take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.

adults 65 years and over ask a doctor and over

children under 12 years of age ask a doctor

consumers with liver or kidney disease ask a doctor

Drug Facts (continued)

Warnings

- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever
- if pregnant or breast-feeding: not recommended
- if pregnant: ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not break or chew tablet; swallow tablet whole

adults and children take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.

adults 65 years and over ask a doctor and over

children under 12 years of age ask a doctor

consumers with liver or kidney disease ask a doctor

Other information

- store between 20° to 25°C (68° to 77°F)

Inactive ingredients

coloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

Stop use and ask doctor if

- an allergic reaction to this product occurs. Seek medical help right away.

When using this product

- **do not use more than directed**
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask doctor if

- an allergic reaction to this product occurs. Seek medical help right away.

Drug Facts (continued)

Active ingredients

Purpose

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: fever, runny nose, sneezing, itchy, watery eyes, itching of the nose or throat, nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Drug Facts (continued)

Active ingredients

Purpose

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: fever, runny nose, sneezing, itchy, watery eyes, itching of the nose or throat, nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Original Prescription Strength

Cetirizine Hydrochloride and Pseudoephedrine Hydrochloride Extended Release Tablets 5 mg / 120 mg Antihistamine/Nasal Decongestant

Indoor & Outdoor Allergies

ALLERGY & CONGESTION

12 Hour Relief of:

- Sneezing
- Runny Nose
- Sinus Pressure
- Itchy, Watery Eyes
- Itchy Throat or Nose
- Nasal Congestion

12 EXTENDED RELEASE TABLETS

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

QUESTIONS? If you have questions of a medical nature, please contact your pharmacist, doctor or health care professional.

Made in India

DISTRIBUTED BY

PERNICO

ALLEGAN, MI 49010

0 000000000000 0

0000000000

0000000000

CODE AREA

1 TABLETS FA C1

Cetirizine Hydrochloride and Pseudoephedrine Hydrochloride Extended Release Tablets 5 mg/120 mg Antihistamine/Nasal Decongestant

ALLERGY & CONGESTION

FORMAT: STANDARD

TYPE SIZES:

TITLE: 9 pt Helvetica Bold Cond. Oblique

TITLE (continued): 8 pt Helvetica Bold Cond. Oblique

(continued) 8 pt Helvetica Cond. Medium

HEADINGS: 8 pt Helvetica Bold Cond. Oblique

SUBHEADINGS: 6 pt Helvetica Bold Cond.

TEXT: 6 pt Helvetica Medium Cond.

LEADING: minimum 6.3 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

EFB1021
FA17662C1

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease ■ thyroid disease ■ diabetes ■ glaucoma ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

- do not use more than directed
- drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.

Inactive ingredients
colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

Other information

- store between 20° to 25°C (68° to 77°F)

Directions

adults and children take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.

adults 65 years and over ask a doctor

children under 12 years of age ask a doctor

consumers with liver or kidney disease ask a doctor

Drug Facts (continued)

■ you get nervous, dizzy, or sleepless ■ symptoms do not improve within 7 days or are accompanied by fever

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts

Active ingredients
(in each extended release tablet)

Cetirizine HCl 5 mg, Pseudoephedrine HCl 120 mg, Antihistamine Pseudoephedrine HCl 120 mg, Nasal Decongestant

Purpose
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: itchy, watery eyes; runny nose; sneezing; itching of the nose or throat; nasal congestion; reduces swelling of nasal passages; temporarily relieves sinus congestion and pressure; temporarily restores free breathing through the nose

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: itchy, watery eyes; runny nose; sneezing; itching of the nose or throat; nasal congestion; reduces swelling of nasal passages; temporarily relieves sinus congestion and pressure; temporarily restores free breathing through the nose

Warnings

■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease ■ thyroid disease ■ diabetes ■ glaucoma ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

- do not use more than directed
- drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.

Inactive ingredients
colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

Other information

- store between 20° to 25°C (68° to 77°F)

Directions

adults and children take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.

adults 65 years and over ask a doctor

children under 12 years of age ask a doctor

consumers with liver or kidney disease ask a doctor

Drug Facts (continued)

■ you get nervous, dizzy, or sleepless ■ symptoms do not improve within 7 days or are accompanied by fever

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Original Prescription Strength

Cetirizine Hydrochloride and Pseudoephedrine Hydrochloride Extended Release Tablets 5 mg/120 mg Antihistamine/Nasal Decongestant

Indoor & Outdoor Allergies

ALLERGY & CONGESTION

12 Hour Relief of:

- Sneezing
- Runny Nose
- Sinus Pressure
- Itchy, Watery Eyes
- Itchy Throat or Nose
- Nasal Congestion


24 EXTENDED RELEASE TABLETS

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

QUESTIONS? If you have questions of a medical nature, please contact your pharmacist, doctor or health care professional.

Made in India

DISTRIBUTED BY
PERIGO
ALLEGAN, MI 49010



FORMAT: STANDARD

TYPE SIZES:

TITLE: 9 pt Helvetica Bold Cond. Oblique

TITLE (continued): 8 pt Helvetica Bold Cond. Oblique

(continued) 8 pt Helvetica Cond. Medium

HEADINGS: 8 pt Helvetica Bold Cond. Oblique

SUBHEADINGS: 6 pt Helvetica Bold Cond.

TEXT: 6 pt Helvetica Medium Cond.

LEADING: minimum 6.3 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt