

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DULOXETINE DELAYED-RELEASE CAPSULES safely and effectively. See full prescribing information for DULOXETINE DELAYED-RELEASE CAPSULES.

DULOXETINE delayed-release capsules USP for oral use.

Initial U.S. Approval: 2004

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS
See full prescribing information for complete boxed warning.
Increased risk of suicidal thinking and behavior in children, adolescents, and young adults taking antidepressants (5.1)
Monitor for worsening and emergence of suicidal thoughts and behaviors (5.1)

RECENT MAJOR CHANGES
Boxed Warning: Suicidal Thoughts and Behaviors 10/2014
Indications and Usage (1) 10/2014
Dosage for Treatment of Generalized Anxiety Disorder (2.2) 10/2014
Contraindications:
Uncontrolled Narrow-Angle Glaucoma (4.2) Removed 07/2014
Warnings and Precautions:
Orthostatic Hypotension, Falls and Syncope (5.3) 11/2014
Angle-Closure Glaucoma (5.9) 07/2014

INDICATIONS AND USAGE
Duloxetine delayed-release capsules are a serotonin and norepinephrine reuptake inhibitor (SNRI) indicated for the treatment of:
Major Depressive Disorder (MDD) (1)
Generalized Anxiety Disorder (GAD) (1)
Diabetic Peripheral Neuropathic Pain (DPNP) (1)
Chronic Musculoskeletal Pain (1)

DOSE AND ADMINISTRATION
Take duloxetine delayed-release capsules once daily, with or without food. Swallow duloxetine delayed-release capsules whole. Do not crush or chew. Do not open capsules. Take missed doses as soon as it is remembered. Do not take two doses of duloxetine delayed-release capsules at the same time. (2)

Table with 4 columns: Indication, Starting Dose, Target Dose, Maximum Dose. Rows include MDD (2.1), GAD (2.2), Adults, DPNP (2.3), Chronic Musculoskeletal Pain (2.5).

ADVERSE REACTIONS
Most common adverse reactions (≥5% and at least twice the incidence of placebo): nausea, dry mouth, somnolence, constipation, decreased appetite, and hyperhidrosis (8.3).

DRUG INTERACTIONS
Some patients may benefit from starting at 30 mg once daily (2)
There is no evidence that doses greater than 60 mg daily confers additional benefit. Some adverse reactions were observed to be dose-dependent (2)
Discontinuing duloxetine delayed-release capsules: Gradually reduce dosage to avoid discontinuation symptoms (2.7, 5.7)
Hepatic Impairment: Avoid use in patients with chronic liver disease or cirrhosis (5.14)

USE IN SPECIFIC POPULATIONS
Pregnancy: Based on animal data may cause fetal harm (8.1)
Nursing Mothers: Exercise caution when administering to a nursing woman (8.3)

CONTRAINDICATIONS
Serotonin Syndrome and MAOIs: Do not use MAOIs intended to treat psychiatric disorders with duloxetine delayed-release capsules or within 5 days of stopping treatment with duloxetine delayed-release capsules. Do not use duloxetine delayed-release capsules within 14 days of stopping an MAOI used to treat psychiatric disorders.

FULL PRESCRIBING INFORMATION: CONTENTS
WARNING: SUICIDAL THOUGHTS AND BEHAVIORS
INDICATIONS AND USAGE
DOSE AND ADMINISTRATION
2.1 Indications and Usage
2.2 Dosage for Treatment of Major Depressive Disorder
2.3 Dosage for Treatment of Generalized Anxiety Disorder
2.4 Dosage for Treatment of Diabetic Peripheral Neuropathic Pain
2.5 Dosage for Treatment of Chronic Musculoskeletal Pain
2.6 Dosing in Specific Populations
2.7 Discontinuing Duloxetine Delayed-Release Capsules
2.8 Switching a Patient to or from a Monoamine Oxidase Inhibitor (MAOI) Intended to Treat Psychiatric Disorders
2.9 Use of Duloxetine Delayed-Release Capsules with Other MAOIs such as Linezolid or Methylene Blue
3 DOSE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
5.1 Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults
5.2 Hepatotoxicity
5.3 Orthostatic Hypotension, Falls and Syncope
5.4 Pregnancy
5.5 Abnormal Bleeding
5.6 Severe Skin Reactions
5.7 Discontinuation of Treatment with Duloxetine Delayed-Release Capsules
5.8 Activation of Mania/Hypomania
5.9 Angle-Closure Glaucoma
5.10 Seizures
5.11 Effect on Blood Pressure
5.12 Clinically Important Drug Interactions
5.13 Hypomania
5.14 Use in Patients with Concomitant Illness
5.15 Urinary Hesitation and Retention
5.16 Laboratory Tests
5.17 Signs and Symptoms
6 GENERAL REACTIONS AND BEHAVIORS IN CHILDREN, ADOLESCENTS, AND YOUNG ADULTS
6.1 Clinical Trial Data Sources
6.2 Adverse Reactions Reported as Reasons for Discontinuation of Treatment in Adult Placebo-Controlled Trials
6.3 Most Common Adult Adverse Reactions
6.4 Adverse Reactions Occurring at an Incidence of 5% or More Among Duloxetine Delayed-Release Capsule-Treated Patients in Adult Placebo-Controlled Trials
6.5 Adverse Reactions Occurring at an Incidence of 2% or More Among Duloxetine Delayed-Release Capsule-Treated Patients in Adult Placebo-Controlled Trials

6.6 Effects on Male and Female Sexual Function in Adults
6.7 Vital Sign Changes in Adults
6.8 Laboratory Changes in Adults
6.9 Electrocardiogram Changes in Adults
6.10 Other Adverse Reactions Observed During the Premarketing and Postmarketing Clinical Trial Evaluation of Duloxetine Delayed-Release Capsules in Adults
6.11 Adverse Reactions Observed in Children and Adolescent Placebo-Controlled Trials
6.12 Postmarketing Spontaneous Reports

6.13 Pharmacokinetics
6.14 Signs and Symptoms
6.15 Laboratory Tests
6.16 Signs and Symptoms
6.17 Signs and Symptoms
6.18 Signs and Symptoms
6.19 Signs and Symptoms
6.20 Signs and Symptoms
6.21 Signs and Symptoms
6.22 Signs and Symptoms
6.23 Signs and Symptoms
6.24 Signs and Symptoms
6.25 Signs and Symptoms
6.26 Signs and Symptoms
6.27 Signs and Symptoms
6.28 Signs and Symptoms
6.29 Signs and Symptoms
6.30 Signs and Symptoms
6.31 Signs and Symptoms
6.32 Signs and Symptoms
6.33 Signs and Symptoms
6.34 Signs and Symptoms
6.35 Signs and Symptoms
6.36 Signs and Symptoms
6.37 Signs and Symptoms
6.38 Signs and Symptoms
6.39 Signs and Symptoms
6.40 Signs and Symptoms
6.41 Signs and Symptoms
6.42 Signs and Symptoms
6.43 Signs and Symptoms
6.44 Signs and Symptoms
6.45 Signs and Symptoms
6.46 Signs and Symptoms
6.47 Signs and Symptoms
6.48 Signs and Symptoms
6.49 Signs and Symptoms
6.50 Signs and Symptoms
6.51 Signs and Symptoms
6.52 Signs and Symptoms
6.53 Signs and Symptoms
6.54 Signs and Symptoms
6.55 Signs and Symptoms
6.56 Signs and Symptoms
6.57 Signs and Symptoms
6.58 Signs and Symptoms
6.59 Signs and Symptoms
6.60 Signs and Symptoms
6.61 Signs and Symptoms
6.62 Signs and Symptoms
6.63 Signs and Symptoms
6.64 Signs and Symptoms
6.65 Signs and Symptoms
6.66 Signs and Symptoms
6.67 Signs and Symptoms
6.68 Signs and Symptoms
6.69 Signs and Symptoms
6.70 Signs and Symptoms
6.71 Signs and Symptoms
6.72 Signs and Symptoms
6.73 Signs and Symptoms
6.74 Signs and Symptoms
6.75 Signs and Symptoms
6.76 Signs and Symptoms
6.77 Signs and Symptoms
6.78 Signs and Symptoms
6.79 Signs and Symptoms
6.80 Signs and Symptoms
6.81 Signs and Symptoms
6.82 Signs and Symptoms
6.83 Signs and Symptoms
6.84 Signs and Symptoms
6.85 Signs and Symptoms
6.86 Signs and Symptoms
6.87 Signs and Symptoms
6.88 Signs and Symptoms
6.89 Signs and Symptoms
6.90 Signs and Symptoms
6.91 Signs and Symptoms
6.92 Signs and Symptoms
6.93 Signs and Symptoms
6.94 Signs and Symptoms
6.95 Signs and Symptoms
6.96 Signs and Symptoms
6.97 Signs and Symptoms
6.98 Signs and Symptoms
6.99 Signs and Symptoms
6.100 Signs and Symptoms

6.101 Signs and Symptoms
6.102 Signs and Symptoms
6.103 Signs and Symptoms
6.104 Signs and Symptoms
6.105 Signs and Symptoms
6.106 Signs and Symptoms
6.107 Signs and Symptoms
6.108 Signs and Symptoms
6.109 Signs and Symptoms
6.110 Signs and Symptoms
6.111 Signs and Symptoms
6.112 Signs and Symptoms
6.113 Signs and Symptoms
6.114 Signs and Symptoms
6.115 Signs and Symptoms
6.116 Signs and Symptoms
6.117 Signs and Symptoms
6.118 Signs and Symptoms
6.119 Signs and Symptoms
6.120 Signs and Symptoms
6.121 Signs and Symptoms
6.122 Signs and Symptoms
6.123 Signs and Symptoms
6.124 Signs and Symptoms
6.125 Signs and Symptoms
6.126 Signs and Symptoms
6.127 Signs and Symptoms
6.128 Signs and Symptoms
6.129 Signs and Symptoms
6.130 Signs and Symptoms
6.131 Signs and Symptoms
6.132 Signs and Symptoms
6.133 Signs and Symptoms
6.134 Signs and Symptoms
6.135 Signs and Symptoms
6.136 Signs and Symptoms
6.137 Signs and Symptoms
6.138 Signs and Symptoms
6.139 Signs and Symptoms
6.140 Signs and Symptoms
6.141 Signs and Symptoms
6.142 Signs and Symptoms
6.143 Signs and Symptoms
6.144 Signs and Symptoms
6.145 Signs and Symptoms
6.146 Signs and Symptoms
6.147 Signs and Symptoms
6.148 Signs and Symptoms
6.149 Signs and Symptoms
6.150 Signs and Symptoms
6.151 Signs and Symptoms
6.152 Signs and Symptoms
6.153 Signs and Symptoms
6.154 Signs and Symptoms
6.155 Signs and Symptoms
6.156 Signs and Symptoms
6.157 Signs and Symptoms
6.158 Signs and Symptoms
6.159 Signs and Symptoms
6.160 Signs and Symptoms
6.161 Signs and Symptoms
6.162 Signs and Symptoms
6.163 Signs and Symptoms
6.164 Signs and Symptoms
6.165 Signs and Symptoms
6.166 Signs and Symptoms
6.167 Signs and Symptoms
6.168 Signs and Symptoms
6.169 Signs and Symptoms
6.170 Signs and Symptoms
6.171 Signs and Symptoms
6.172 Signs and Symptoms
6.173 Signs and Symptoms
6.174 Signs and Symptoms
6.175 Signs and Symptoms
6.176 Signs and Symptoms
6.177 Signs and Symptoms
6.178 Signs and Symptoms
6.179 Signs and Symptoms
6.180 Signs and Symptoms
6.181 Signs and Symptoms
6.182 Signs and Symptoms
6.183 Signs and Symptoms
6.184 Signs and Symptoms
6.185 Signs and Symptoms
6.186 Signs and Symptoms
6.187 Signs and Symptoms
6.188 Signs and Symptoms
6.189 Signs and Symptoms
6.190 Signs and Symptoms
6.191 Signs and Symptoms
6.192 Signs and Symptoms
6.193 Signs and Symptoms
6.194 Signs and Symptoms
6.195 Signs and Symptoms
6.196 Signs and Symptoms
6.197 Signs and Symptoms
6.198 Signs and Symptoms
6.199 Signs and Symptoms
6.200 Signs and Symptoms

6.201 Signs and Symptoms
6.202 Signs and Symptoms
6.203 Signs and Symptoms
6.204 Signs and Symptoms
6.205 Signs and Symptoms
6.206 Signs and Symptoms
6.207 Signs and Symptoms
6.208 Signs and Symptoms
6.209 Signs and Symptoms
6.210 Signs and Symptoms
6.211 Signs and Symptoms
6.212 Signs and Symptoms
6.213 Signs and Symptoms
6.214 Signs and Symptoms
6.215 Signs and Symptoms
6.216 Signs and Symptoms
6.217 Signs and Symptoms
6.218 Signs and Symptoms
6.219 Signs and Symptoms
6.220 Signs and Symptoms
6.221 Signs and Symptoms
6.222 Signs and Symptoms
6.223 Signs and Symptoms
6.224 Signs and Symptoms
6.225 Signs and Symptoms
6.226 Signs and Symptoms
6.227 Signs and Symptoms
6.228 Signs and Symptoms
6.229 Signs and Symptoms
6.230 Signs and Symptoms
6.231 Signs and Symptoms
6.232 Signs and Symptoms
6.233 Signs and Symptoms
6.234 Signs and Symptoms
6.235 Signs and Symptoms
6.236 Signs and Symptoms
6.237 Signs and Symptoms
6.238 Signs and Symptoms
6.239 Signs and Symptoms
6.240 Signs and Symptoms
6.241 Signs and Symptoms
6.242 Signs and Symptoms
6.243 Signs and Symptoms
6.244 Signs and Symptoms
6.245 Signs and Symptoms
6.246 Signs and Symptoms
6.247 Signs and Symptoms
6.248 Signs and Symptoms
6.249 Signs and Symptoms
6.250 Signs and Symptoms
6.251 Signs and Symptoms
6.252 Signs and Symptoms
6.253 Signs and Symptoms
6.254 Signs and Symptoms
6.255 Signs and Symptoms
6.256 Signs and Symptoms
6.257 Signs and Symptoms
6.258 Signs and Symptoms
6.259 Signs and Symptoms
6.260 Signs and Symptoms
6.261 Signs and Symptoms
6.262 Signs and Symptoms
6.263 Signs and Symptoms
6.264 Signs and Symptoms
6.265 Signs and Symptoms
6.266 Signs and Symptoms
6.267 Signs and Symptoms
6.268 Signs and Symptoms
6.269 Signs and Symptoms
6.270 Signs and Symptoms
6.271 Signs and Symptoms
6.272 Signs and Symptoms
6.273 Signs and Symptoms
6.274 Signs and Symptoms
6.275 Signs and Symptoms
6.276 Signs and Symptoms
6.277 Signs and Symptoms
6.278 Signs and Symptoms
6.279 Signs and Symptoms
6.280 Signs and Symptoms
6.281 Signs and Symptoms
6.282 Signs and Symptoms
6.283 Signs and Symptoms
6.284 Signs and Symptoms
6.285 Signs and Symptoms
6.286 Signs and Symptoms
6.287 Signs and Symptoms
6.288 Signs and Symptoms
6.289 Signs and Symptoms
6.290 Signs and Symptoms
6.291 Signs and Symptoms
6.292 Signs and Symptoms
6.293 Signs and Symptoms
6.294 Signs and Symptoms
6.295 Signs and Symptoms
6.296 Signs and Symptoms
6.297 Signs and Symptoms
6.298 Signs and Symptoms
6.299 Signs and Symptoms
6.300 Signs and Symptoms

6.301 Signs and Symptoms
6.302 Signs and Symptoms
6.303 Signs and Symptoms
6.304 Signs and Symptoms
6.305 Signs and Symptoms
6.306 Signs and Symptoms
6.307 Signs and Symptoms
6.308 Signs and Symptoms
6.309 Signs and Symptoms
6.310 Signs and Symptoms
6.311 Signs and Symptoms
6.312 Signs and Symptoms
6.313 Signs and Symptoms
6.314 Signs and Symptoms
6.315 Signs and Symptoms
6.316 Signs and Symptoms
6.317 Signs and Symptoms
6.318 Signs and Symptoms
6.319 Signs and Symptoms
6.320 Signs and Symptoms
6.321 Signs and Symptoms
6.322 Signs and Symptoms
6.323 Signs and Symptoms
6.324 Signs and Symptoms
6.325 Signs and Symptoms
6.326 Signs and Symptoms
6.327 Signs and Symptoms
6.328 Signs and Symptoms
6.329 Signs and Symptoms
6.330 Signs and Symptoms
6.331 Signs and Symptoms
6.332 Signs and Symptoms
6.333 Signs and Symptoms
6.334 Signs and Symptoms
6.335 Signs and Symptoms
6.336 Signs and Symptoms
6.337 Signs and Symptoms
6.338 Signs and Symptoms
6.339 Signs and Symptoms
6.340 Signs and Symptoms
6.341 Signs and Symptoms
6.342 Signs and Symptoms
6.343 Signs and Symptoms
6.344 Signs and Symptoms
6.345 Signs and Symptoms
6.346 Signs and Symptoms
6.347 Signs and Symptoms
6.348 Signs and Symptoms
6.349 Signs and Symptoms
6.350 Signs and Symptoms
6.351 Signs and Symptoms
6.352 Signs and Symptoms
6.353 Signs and Symptoms
6.354 Signs and Symptoms
6.355 Signs and Symptoms
6.356 Signs and Symptoms
6.357 Signs and Symptoms
6.358 Signs and Symptoms
6.359 Signs and Symptoms
6.360 Signs and Symptoms
6.361 Signs and Symptoms
6.362 Signs and Symptoms
6.363 Signs and Symptoms
6.364 Signs and Symptoms
6.365 Signs and Symptoms
6.366 Signs and Symptoms
6.367 Signs and Symptoms
6.368 Signs and Symptoms
6.369 Signs and Symptoms
6.370 Signs and Symptoms
6.371 Signs and Symptoms
6.372 Signs and Symptoms
6.373 Signs and Symptoms
6.374 Signs and Symptoms
6.375 Signs and Symptoms
6.376 Signs and Symptoms
6.377 Signs and Symptoms
6.378 Signs and Symptoms
6.379 Signs and Symptoms
6.380 Signs and Symptoms
6.381 Signs and Symptoms
6.382 Signs and Symptoms
6.383 Signs and Symptoms
6.384 Signs and Symptoms
6.385 Signs and Symptoms
6.386 Signs and Symptoms
6.387 Signs and Symptoms
6.388 Signs and Symptoms
6.389 Signs and Symptoms
6.390 Signs and Symptoms
6.391 Signs and Symptoms
6.392 Signs and Symptoms
6.393 Signs and Symptoms
6.394 Signs and Symptoms
6.395 Signs and Symptoms
6.396 Signs and Symptoms
6.397 Signs and Symptoms
6.398 Signs and Symptoms
6.399 Signs and Symptoms
6.400 Signs and Symptoms

psychiatric disorders. In addition, do not start duloxetine delayed-release capsule in a patient who is being treated with linezolid or intravenous methylene blue (4)

WARNINGS AND PRECAUTIONS
Hepatotoxicity: Hepatic failure has been reported in patients treated with duloxetine delayed-release capsules. Duloxetine delayed-release capsules should be discontinued in patients who develop jaundice or other evidence of clinically significant liver dysfunction and should not be resumed unless another cause can be established. Duloxetine delayed-release capsules should not be prescribed to patients with substantial alcohol use or evidence of chronic liver disease (5.2)
Orthostatic Hypotension, Falls and Syncope: Cases have been reported with duloxetine delayed-release capsules (5.3)
Serotonin Syndrome: Serotonin syndrome has been reported with SSRIs and SNRIs, including with duloxetine delayed-release capsules, both when taken alone, but especially when coadministered with other serotonergic agents, including triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, tryptophan, buspirone and St. John's Wort. If such symptoms occur, discontinue duloxetine delayed-release capsules and initiate supportive treatment. If concomitant use of duloxetine delayed-release capsules with other serotonergic drugs is clinically warranted, patients should be made aware of a potential increased risk for serotonin syndrome, particularly during treatment initiation and dose increases (5.4)
Abnormal Bleeding: Duloxetine delayed-release capsules may increase the risk of bleeding events. Patients should be cautioned about the risk of bleeding associated with the concomitant use of duloxetine delayed-release capsules and NSAIDs, aspirin, or other drugs that affect coagulation (5.5, 7.4)
Severe Skin Reactions: Severe skin reactions, including erythema multiforme and Stevens-Johnson Syndrome (SJS), can occur with duloxetine delayed-release capsules. Duloxetine delayed-release capsules should be discontinued at the first appearance of blisters, peeling rash, mucosal erosions, or any other sign of hypersensitivity if no other etiology can be identified (5.6)
Discontinuation: May result in symptoms, including dizziness, headache, nausea, diarrhea, paresthesia, irritability, vomiting, insomnia, anxiety, hyperhidrosis, and fatigue (5.7)
Activation of mania or hypomania has been reported in a small proportion of patients who were treated with other marketed drugs effective in the treatment of major depressive disorder. In such cases, duloxetine delayed-release capsules should be used cautiously in patients with a history of manic or hypomanic episodes (5.8)
Angle-Closure Glaucoma: Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)
Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)

WARNINGS AND PRECAUTIONS
Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults: Patients with major depressive disorder (MDD), both adult and pediatric, may experience worsening of suicidal thoughts and actions, including suicidal ideation, thoughts of self-harm, or suicide, either before or after they are taking antidepressant medication, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain psychiatric disorders, and these risks may be increased by antidepressant treatment. There has been one report of suicidal ideation and attempts in patients treated with duloxetine delayed-release capsules. Patients and caregivers should be monitored for worsening of depression and the emergence of suicidal thoughts and actions during early treatment and during dose changes (5.1)
Discontinuation: May result in symptoms, including dizziness, headache, nausea, diarrhea, paresthesia, irritability, vomiting, insomnia, anxiety, hyperhidrosis, and fatigue (5.7)
Activation of mania or hypomania has been reported in a small proportion of patients who were treated with other marketed drugs effective in the treatment of major depressive disorder. In such cases, duloxetine delayed-release capsules should be used cautiously in patients with a history of manic or hypomanic episodes (5.8)
Angle-Closure Glaucoma: Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)
Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)

ADVERSE REACTIONS
Most common adverse reactions (≥5% and at least twice the incidence of placebo): nausea, dry mouth, somnolence, constipation, decreased appetite, and hyperhidrosis (8.3).
To report SUSPECTED ADVERSE REACTIONS, contact Par Pharmaceutical, Inc. at 1-800-828-9339 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS
Some patients may benefit from starting at 30 mg once daily (2)
There is no evidence that doses greater than 60 mg daily confers additional benefit. Some adverse reactions were observed to be dose-dependent (2)
Discontinuing duloxetine delayed-release capsules: Gradually reduce dosage to avoid discontinuation symptoms (2.7, 5.7)
Hepatic Impairment: Avoid use in patients with chronic liver disease or cirrhosis (5.14)

USE IN SPECIFIC POPULATIONS
Pregnancy: Based on animal data may cause fetal harm (8.1)
Nursing Mothers: Exercise caution when administering to a nursing woman (8.3)

CONTRAINDICATIONS
Serotonin Syndrome and MAOIs: Do not use MAOIs intended to treat psychiatric disorders with duloxetine delayed-release capsules or within 5 days of stopping treatment with duloxetine delayed-release capsules. Do not use duloxetine delayed-release capsules within 14 days of stopping an MAOI used to treat psychiatric disorders.

INDICATIONS AND USAGE
Duloxetine delayed-release capsules are a serotonin and norepinephrine reuptake inhibitor (SNRI) indicated for the treatment of:
Major Depressive Disorder (MDD) (1)
Generalized Anxiety Disorder (GAD) (1)
Diabetic Peripheral Neuropathic Pain (DPNP) (1)
Chronic Musculoskeletal Pain (1)

DOSE AND ADMINISTRATION
Take duloxetine delayed-release capsules once daily, with or without food. Swallow duloxetine delayed-release capsules whole. Do not crush or chew. Do not open capsules. Take missed doses as soon as it is remembered. Do not take two doses of duloxetine delayed-release capsules at the same time. (2)

ADVERSE REACTIONS
Most common adverse reactions (≥5% and at least twice the incidence of placebo): nausea, dry mouth, somnolence, constipation, decreased appetite, and hyperhidrosis (8.3).

DRUG INTERACTIONS
Some patients may benefit from starting at 30 mg once daily (2)
There is no evidence that doses greater than 60 mg daily confers additional benefit. Some adverse reactions were observed to be dose-dependent (2)
Discontinuing duloxetine delayed-release capsules: Gradually reduce dosage to avoid discontinuation symptoms (2.7, 5.7)
Hepatic Impairment: Avoid use in patients with chronic liver disease or cirrhosis (5.14)

USE IN SPECIFIC POPULATIONS
Pregnancy: Based on animal data may cause fetal harm (8.1)
Nursing Mothers: Exercise caution when administering to a nursing woman (8.3)

some cases, a patient already receiving duloxetine delayed-release capsules may require urgent treatment with linezolid or intravenous methylene blue. If acceptable alternatives to linezolid or intravenous methylene blue are not available and the potential benefits of duloxetine delayed-release capsules outweigh the risks, duloxetine delayed-release capsules should be used cautiously in patients with a history of manic or hypomanic episodes (5.8)
Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)

DOSE FORMS AND STRENGTHS
Duloxetine Delayed-Release Capsules, USP are available as delayed-release capsules:
20 mg hard gelatin capsules with dark orange cap and light orange body, imprinted with 'A156' on the cap and '156' on the body.
30 mg hard gelatin capsules with yellow cap and yellow body, imprinted with 'A157' on the cap and '30' on the body, containing white to off white spherical coated pellets.
40 mg hard gelatin capsules with yellow cap and light orange body, imprinted with 'A158' on the cap and '40' on the body, containing white to off white spherical coated pellets.
60 mg hard gelatin capsules with yellow cap and light orange body, imprinted with 'A159' on the cap and '60' on the body, containing white to off white spherical coated pellets.

CONTRAINDICATIONS
Serotonin Syndrome and MAOIs: Do not use MAOIs intended to treat psychiatric disorders with duloxetine delayed-release capsules or within 5 days of stopping treatment with duloxetine delayed-release capsules. Do not use duloxetine delayed-release capsules within 14 days of stopping an MAOI intended to treat psychiatric disorders. It is also contraindicated to start duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)
Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)

WARNINGS AND PRECAUTIONS
Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults: Patients with major depressive disorder (MDD), both adult and pediatric, may experience worsening of suicidal thoughts and actions, including suicidal ideation, thoughts of self-harm, or suicide, either before or after they are taking antidepressant medication, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain psychiatric disorders, and these risks may be increased by antidepressant treatment. There has been one report of suicidal ideation and attempts in patients treated with duloxetine delayed-release capsules. Patients and caregivers should be monitored for worsening of depression and the emergence of suicidal thoughts and actions during early treatment and during dose changes (5.1)
Discontinuation: May result in symptoms, including dizziness, headache, nausea, diarrhea, paresthesia, irritability, vomiting, insomnia, anxiety, hyperhidrosis, and fatigue (5.7)
Activation of mania or hypomania has been reported in a small proportion of patients who were treated with other marketed drugs effective in the treatment of major depressive disorder. In such cases, duloxetine delayed-release capsules should be used cautiously in patients with a history of manic or hypomanic episodes (5.8)
Angle-Closure Glaucoma: Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)
Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)

ADVERSE REACTIONS
Most common adverse reactions (≥5% and at least twice the incidence of placebo): nausea, dry mouth, somnolence, constipation, decreased appetite, and hyperhidrosis (8.3).
To report SUSPECTED ADVERSE REACTIONS, contact Par Pharmaceutical, Inc. at 1-800-828-9339 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS
Some patients may benefit from starting at 30 mg once daily (2)
There is no evidence that doses greater than 60 mg daily confers additional benefit. Some adverse reactions were observed to be dose-dependent (2)
Discontinuing duloxetine delayed-release capsules: Gradually reduce dosage to avoid discontinuation symptoms (2.7, 5.7)
Hepatic Impairment: Avoid use in patients with chronic liver disease or cirrhosis (5.14)

USE IN SPECIFIC POPULATIONS
Pregnancy: Based on animal data may cause fetal harm (8.1)
Nursing Mothers: Exercise caution when administering to a nursing woman (8.3)

CONTRAINDICATIONS
Serotonin Syndrome and MAOIs: Do not use MAOIs intended to treat psychiatric disorders with duloxetine delayed-release capsules or within 5 days of stopping treatment with duloxetine delayed-release capsules. Do not use duloxetine delayed-release capsules within 14 days of stopping an MAOI used to treat psychiatric disorders.

INDICATIONS AND USAGE
Duloxetine delayed-release capsules are a serotonin and norepinephrine reuptake inhibitor (SNRI) indicated for the treatment of:
Major Depressive Disorder (MDD) (1)
Generalized Anxiety Disorder (GAD) (1)
Diabetic Peripheral Neuropathic Pain (DPNP) (1)
Chronic Musculoskeletal Pain (1)

DOSE AND ADMINISTRATION
Take duloxetine delayed-release capsules once daily, with or without food. Swallow duloxetine delayed-release capsules whole. Do not crush or chew. Do not open capsules. Take missed doses as soon as it is remembered. Do not take two doses of duloxetine delayed-release capsules at the same time. (2)

ADVERSE REACTIONS
Most common adverse reactions (≥5% and at least twice the incidence of placebo): nausea, dry mouth, somnolence, constipation, decreased appetite, and hyperhidrosis (8.3).

DOSE FORMS AND STRENGTHS
Duloxetine Delayed-Release Capsules, USP are available as delayed-release capsules:
20 mg hard gelatin capsules with dark orange cap and light orange body, imprinted with 'A156' on the cap and '156' on the body.
30 mg hard gelatin capsules with yellow cap and yellow body, imprinted with 'A157' on the cap and '30' on the body, containing white to off white spherical coated pellets.
40 mg hard gelatin capsules with yellow cap and light orange body, imprinted with 'A158' on the cap and '40' on the body, containing white to off white spherical coated pellets.
60 mg hard gelatin capsules with yellow cap and light orange body, imprinted with 'A159' on the cap and '60' on the body, containing white to off white spherical coated pellets.

CONTRAINDICATIONS
Serotonin Syndrome and MAOIs: Do not use MAOIs intended to treat psychiatric disorders with duloxetine delayed-release capsules or within 5 days of stopping treatment with duloxetine delayed-release capsules. Do not use duloxetine delayed-release capsules within 14 days of stopping an MAOI intended to treat psychiatric disorders. It is also contraindicated to start duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)
Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)

WARNINGS AND PRECAUTIONS
Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults: Patients with major depressive disorder (MDD), both adult and pediatric, may experience worsening of suicidal thoughts and actions, including suicidal ideation, thoughts of self-harm, or suicide, either before or after they are taking antidepressant medication, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain psychiatric disorders, and these risks may be increased by antidepressant treatment. There has been one report of suicidal ideation and attempts in patients treated with duloxetine delayed-release capsules. Patients and caregivers should be monitored for worsening of depression and the emergence of suicidal thoughts and actions during early treatment and during dose changes (5.1)
Discontinuation: May result in symptoms, including dizziness, headache, nausea, diarrhea, paresthesia, irritability, vomiting, insomnia, anxiety, hyperhidrosis, and fatigue (5.7)
Activation of mania or hypomania has been reported in a small proportion of patients who were treated with other marketed drugs effective in the treatment of major depressive disorder. In such cases, duloxetine delayed-release capsules should be used cautiously in patients with a history of manic or hypomanic episodes (5.8)
Angle-Closure Glaucoma: Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)
Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)

ADVERSE REACTIONS
Most common adverse reactions (≥5% and at least twice the incidence of placebo): nausea, dry mouth, somnolence, constipation, decreased appetite, and hyperhidrosis (8.3).

DOSE FORMS AND STRENGTHS
Duloxetine Delayed-Release Capsules, USP are available as delayed-release capsules:
20 mg hard gelatin capsules with dark orange cap and light orange body, imprinted with 'A156' on the cap and '156' on the body.
30 mg hard gelatin capsules with yellow cap and yellow body, imprinted with 'A157' on the cap and '30' on the body, containing white to off white spherical coated pellets.
40 mg hard gelatin capsules with yellow cap and light orange body, imprinted with 'A158' on the cap and '40' on the body, containing white to off white spherical coated pellets.
60 mg hard gelatin capsules with yellow cap and light orange body, imprinted with 'A159' on the cap and '60' on the body, containing white to off white spherical coated pellets.

CONTRAINDICATIONS
Serotonin Syndrome and MAOIs: Do not use MAOIs intended to treat psychiatric disorders with duloxetine delayed-release capsules or within 5 days of stopping treatment with duloxetine delayed-release capsules. Do not use duloxetine delayed-release capsules within 14 days of stopping an MAOI intended to treat psychiatric disorders. It is also contraindicated to start duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)
Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)

WARNINGS AND PRECAUTIONS
Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults: Patients with major depressive disorder (MDD), both adult and pediatric, may experience worsening of suicidal thoughts and actions, including suicidal ideation, thoughts of self-harm, or suicide, either before or after they are taking antidepressant medication, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain psychiatric disorders, and these risks may be increased by antidepressant treatment. There has been one report of suicidal ideation and attempts in patients treated with duloxetine delayed-release capsules. Patients and caregivers should be monitored for worsening of depression and the emergence of suicidal thoughts and actions during early treatment and during dose changes (5.1)
Discontinuation: May result in symptoms, including dizziness, headache, nausea, diarrhea, paresthesia, irritability, vomiting, insomnia, anxiety, hyperhidrosis, and fatigue (5.7)
Activation of mania or hypomania has been reported in a small proportion of patients who were treated with other marketed drugs effective in the treatment of major depressive disorder. In such cases, duloxetine delayed-release capsules should be used cautiously in patients with a history of manic or hypomanic episodes (5.8)
Angle-Closure Glaucoma: Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)
Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)

ADVERSE REACTIONS
Most common adverse reactions (≥5% and at least twice the incidence of placebo): nausea, dry mouth, somnolence, constipation, decreased appetite, and hyperhidrosis (8.3).

DOSE FORMS AND STRENGTHS
Duloxetine Delayed-Release Capsules, USP are available as delayed-release capsules:
20 mg hard gelatin capsules with dark orange cap and light orange body, imprinted with 'A156' on the cap and '156' on the body.
30 mg hard gelatin capsules with yellow cap and yellow body, imprinted with 'A157' on the cap and '30' on the body, containing white to off white spherical coated pellets.
40 mg hard gelatin capsules with yellow cap and light orange body, imprinted with 'A158' on the cap and '40' on the body, containing white to off white spherical coated pellets.
60 mg hard gelatin capsules with yellow cap and light orange body, imprinted with 'A159' on the cap and '60' on the body, containing white to off white spherical coated pellets.

CONTRAINDICATIONS
Serotonin Syndrome and MAOIs: Do not use MAOIs intended to treat psychiatric disorders with duloxetine delayed-release capsules or within 5 days of stopping treatment with duloxetine delayed-release capsules. Do not use duloxetine delayed-release capsules within 14 days of stopping an MAOI intended to treat psychiatric disorders. It is also contraindicated to start duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)
Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)

WARNINGS AND PRECAUTIONS
Suicidal Thoughts and Behaviors in Children, Adolescents, and Young Adults: Patients with major depressive disorder (MDD), both adult and pediatric, may experience worsening of suicidal thoughts and actions, including suicidal ideation, thoughts of self-harm, or suicide, either before or after they are taking antidepressant medication, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain psychiatric disorders, and these risks may be increased by antidepressant treatment. There has been one report of suicidal ideation and attempts in patients treated with duloxetine delayed-release capsules. Patients and caregivers should be monitored for worsening of depression and the emergence of suicidal thoughts and actions during early treatment and during dose changes (5.1)
Discontinuation: May result in symptoms, including dizziness, headache, nausea, diarrhea, paresthesia, irritability, vomiting, insomnia, anxiety, hyperhidrosis, and fatigue (5.7)
Activation of mania or hypomania has been reported in a small proportion of patients who were treated with other marketed drugs effective in the treatment of major depressive disorder. In such cases, duloxetine delayed-release capsules should be used cautiously in patients with a history of manic or hypomanic episodes (5.8)
Angle-Closure Glaucoma: Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)
Starting duloxetine delayed-release capsules in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome (5.9)

ADVERSE REACTIONS
Most common adverse reactions (≥5% and at least twice the incidence of placebo): nausea, dry mouth, somnolence, constipation, decreased appetite, and hyperhidrosis (8.3).

DOSE FORMS AND STRENGTHS
Duloxetine Delayed-Release Capsules, USP are available as delayed-release capsules:
20 mg hard gelatin capsules with dark orange cap and light orange body, imprinted with 'A156' on the cap and '156' on the body.
30 mg hard gelatin capsules with yellow cap and yellow body, imprinted with 'A157' on the cap and '30' on the body, containing white to off white spherical coated pellets.
40 mg hard gelatin capsules with yellow cap and light orange body, imprinted with 'A158' on the cap and '40' on the body, containing white to off white spherical coated pellets.
60 mg hard gelatin capsules with yellow cap and light orange body, imprinted with 'A159' on the cap and '60' on the

