

Children's Allegra Allergy 12 Hour Oral Suspension Carton

05.02.16

ALG Kids_12hr_4oz_Liquid CTN_2015_30pct



Children's Allegra[®]
ALLERGY 12 HR

NON-DROWSY

NDC XXXXX-XXXX-X

NON-DROWSY
2 Years & Older

Children's Allegra[®]
ALLERGY 12 HR

Ages 2 Years & Older

Drug Facts (continued)

Other information

- each 5 mL contains: **sodium 18 mg**
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)

Inactive ingredients

butylparaben, edetate disodium, flavor, poloxamer 407, propylene glycol, propylparaben, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

Questions or comments?

call toll-free 1-800-633-1610 or www.allegra.com

The makers of Allegra[®] do not make store brand products. The trade dress of this Allegra package is subject to trademark protection.
 Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2015
 Origin Germany

XXXXXXXX-XX

Children's Allegra[®]
ALLERGY 12 HR
 fexofenadine HCl oral suspension
30 mg/5 ml antihistamine

INDOOR / OUTDOOR ALLERGY RELIEF

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

LIQUID

Oral Suspension
 alcohol free / dye free

4 fl. oz. (120 ml)

Children's Allegra[®]
ALLERGY 12 HR

Ages 2 Years & Older



berry
 FLAVOR

Use Only With Enclosed Dosing Cup
 Wash and let air dry after each use

alcohol free / dye free

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Fexofenadine HCl 30 mg	Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake well before using

- use only with enclosed dosing cup

adults and children 12 years of age and over	take 10 mL every 12 hours; do not take more than 20 mL in 24 hours
children 2 to under 12 years of age	take 5 mL every 12 hours; do not take more than 10 mL in 24 hours
children under 2 years of age	ask a doctor
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Note: mL = milliliters

**UPC
 FPO**

LOT:

EXP:

Children's Allegra Allergy 12 Hour Oral Suspension Label

12.02.15

ALG Kids_12hr_4oz_LBL_2015_r2

NDC XXXXX-XXXX-X

NON-DROWSY
2 Years & Older

Children's Allegra[®] ALLERGY 12 HR
fexofenadine HCl oral suspension
30 mg/5 ml antihistamine

INDOOR / OUTDOOR ALLERGY RELIEF
Dosing Cup Included
Wash and let air dry after each use

Oral Suspension 4 fl. oz. (120 ml)

Active ingredient (in each 5 mL)
Fexofenadine HCl 30 mg.....Antihistamine
Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat
Warnings **Do not use** if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have kidney disease.** Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
Directions ■ shake well before using ▶

Purpose ■ use only with enclosed dosing cup ■ **adults and children 12 years of age and over** take 10 mL every 12 hours; do not take more than 20 mL in 24 hours ■ **children 2 to under 12 years of age** take 5 mL every 12 hours; do not take more than 10 mL in 24 hours ■ **children under 2 years of age** ask a doctor ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor
Note: mL = milliliters **Other information**
■ each 5 mL contains: **sodium 18 mg** ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F)
Inactive ingredients butylparaben, edetate disodium, flavor, poloxamer 407, propylene glycol, propylparaben, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol **Questions or comments?** call toll-free **1-800-633-1610** or www.allegra.com

Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
The trade dress of this Allegra[®] package is subject to trademark protection. © 2015 Origin Germany

XXXXXXX
LOT
EXP

Children's Allegra Allergy 12 Hour Oral Suspension Carton

06.10.16

ALG Kids_12hr_8oz_Liquid CTN_2015_30pct_r1

Children's Allegra ALLERGY 12 HR
NON-DROWSY

Children's Allegra ALLERGY 12 HR
Ages 2 Years & Older

Children's Allegra ALLERGY 12 HR
Ages 2 Years & Older

Children's Allegra ALLERGY 12 HR
Ages 2 Years & Older

Children's Allegra ALLERGY 12 HR
fexofenadine HCl oral suspension
30 mg/5 ml antihistamine

INDOOR / OUTDOOR ALLERGY RELIEF

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

LIQUID Oral Suspension
alcohol free / dye free
8 fl. oz. (240 ml)

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Fexofenadine HCl 30 mg	Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake well before using
■ use only with enclosed dosing cup

adults and children 12 years of age and over	take 10 mL every 12 hours; do not take more than 20 mL in 24 hours
children 2 to under 12 years of age	take 5 mL every 12 hours; do not take more than 10 mL in 24 hours
children under 2 years of age	ask a doctor
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Note: mL = milliliters

Other information

- each 5 mL contains: sodium 18 mg
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)

Inactive ingredients
butylparaben, edetate disodium, flavor, poloxamer 407, propylene glycol, propylparaben, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

Questions or comments?
call toll-free 1-800-633-1610 or www.allegra.com

The makers of Allegra® do not make store brand products. The trade dress of this Allegra package is subject to trademark protection.
Dist. By: Chatterm, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2015 Origin Germany

UPC FPO

LOT:
EXP:

Bullets:
6 point
Universal News CommPi

Annotations:

- Title: 14 point Helvetica Bold italic
- Headings: 8 point Helvetica Bold
- Subheadings: 6 point Helvetica Bold
- Body Text: 6 point Helvetica Regular
- 11 point Helvetica Rounded Bold
- 6 point Helvetica Neue Condensed
- 9 point MotherCare Reg
- 8 point MotherCare Reg
- 9 point SharpSansNo1 Book Itc
- table format for 3 or more dosages
- 0.5 point hairline
- 2.5 point box barline
- 2.5 point barline
- 6 point Helvetica Neue Condensed

Children's Allegra Allergy 12 Hour Oral Suspension Label

06.09.16

ALG Kids_12hr_8oz_LBL_2015

NDC XXXXX-XXXX-X

NON-DROWSY
2 Years & Older

Children's Allegra[®] ALLERGY 12 HR
fexofenadine HCl oral suspension
30 mg/5 ml antihistamine

INDOOR / OUTDOOR ALLERGY RELIEF
Dosing Cup Included
Wash and let air dry after each use

Oral Suspension 8 fl. oz. (240 ml)

Active ingredient (in each 5 mL)
Fexofenadine HCl 30 mg.....Antihistamine
Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat
Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.
Directions ■ shake well before using

Purpose
■ use only with enclosed dosing cup ■ **adults and children 12 years of age and over** take 10 mL every 12 hours; do not take more than 20 mL in 24 hours ■ **children 2 to under 12 years of age** take 5 mL every 12 hours; do not take more than 10 mL in 24 hours ■ **children under 2 years of age** ask a doctor ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor Note: mL = milliliters **Other information** ■ each 5 mL contains: **sodium 18 mg** ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F)
Inactive ingredients butylparaben, edetate disodium, flavor, poloxamer 407, propylene glycol, propylparaben, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol **Questions or comments?** call toll-free **1-800-633-1610** or www.allegra.com


XXXXXX-XX
LOT
EXP

Dist. By: Chatterm, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
The trade dress of this Allegra[®] package is subject to trademark protection. © 2015 Origin Germany

Children's Allegra Allergy 12 Hour Oral Suspension Dosing Cup

12.2.15

ALG Kids_12hr_4oz_Liquid_Dosing Cup_2015_r1



This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
06/15/2016