

873293



**Children's Allegra<sup>®</sup>**  
**ALLERGY**

**12 HR**

NON-DROWSY

NDC 41167-4244-4

**2 Years & Older**

**Children's Allegra<sup>®</sup>**  
**ALLERGY**

**12 HR**

**Ages 2 Years & Older**

**Drug Facts (continued)**

**Other information**

- each 5 mL contains: potassium 5 mg and sodium 19 mg
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- color change from white to beige is expected over time. This does not impact product effectiveness and safety.

**Inactive ingredients**

edetate disodium, flavor, poloxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xyllitol

**Questions or comments?**  
call 1-866-844-2798

The makers of Allegra<sup>®</sup> do not make store brand products. The trade dress of this Allegra<sup>®</sup> package is subject to trademark protection.

Dist. By: Chatterm, Inc., a Sanofi Company  
Chattanooga, TN 37409-0219 ©2022  
Origin Germany

Alcohol & Dye  
**FREE**



**Children's Allegra<sup>®</sup>**  
**ALLERGY**

fexofenadine HCl oral suspension  
30 mg/5 mL antihistamine

**NON-DROWSY**  
Liquid

**12 HR**

INDOOR / OUTDOOR  
ALLERGY RELIEF

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

**4 fl. oz. (120 mL)**



**Berry**  
FLAVOR

**FREE OF:**  
ALCOHOL  
& DYES

**Use Only With**  
**Enclosed Dosing Cup**  
Wash and let air dry after each use

**Children's Allegra<sup>®</sup>**  
**ALLERGY**

**12 HR**

**Ages 2 Years & Older**

**Drug Facts**

Active ingredient (in each 5 mL)	Purpose
Fexofenadine HCl 30 mg	Antihistamine

**Uses**  
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

**Warnings**  
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.  
Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

**When using this product**

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ shake well before using  
■ use only with enclosed dosing cup

adults and children 12 years of age and over	take 10 mL every 12 hours; do not take more than 20 mL in 24 hours
children 2 to under 12 years of age	take 5 mL every 12 hours; do not take more than 10 mL in 24 hours
children under 2 years of age	ask a doctor
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Note: mL = milliliters



0 41167 42444 5

LOT:

EXP:



873293

**Children's Allegra Allergy 12 Hour 4oz Berry Liquid**  
**Liquid 4oz Carton**  
**12.01.22**

Allegra\_Children's\_Allergy\_12hr\_Berry\_4oz\_CTN\_Germany\_FDA\_873293\_DSN.ai  
(PDF of art being shown is submitted at 100% actual size)

# Children's Allegra Allergy 12 Hour 4 oz Berry Flavor Liquid Liquid 4 oz Bottle Label

06.02.22

Allegra\_Children's\_Allergy\_12hr\_Berry\_4oz\_Wrap\_845643-01\_DSN4.ai  
(PDF of art being shown is submitted at 100% actual size)

NDC 41167-4244-4

**NON-DROWSY**  
**2 Years & Older**

Alcohol & Dye **FREE**

**Children's Allegra**  
**ALLERGY**

fexofenadine HCl oral suspension  
**30 mg/5 mL antihistamine**

**12 HR**  
**Berry**  
FLAVOR

**INDOOR / OUTDOOR ALLERGY RELIEF**

Dosing Cup Included  
Wash & let air dry after each use

**4 fl. oz. (120 mL)**

**Active ingredient (in each 5 mL)**  
Fexofenadine HCl 30 mg.....Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  
■ runny nose ■ itchy, watery eyes ■ sneezing  
■ itching of the nose or throat

**Warnings Do not use** if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions)

**Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** ■ shake well before using ■ use only with enclosed dosing cup ■ **adults and children 12 years of age and over** take 10 mL every 12 hours; do not take more

**Purpose** than 20 mL in 24 hours ■ **children 2 to under 12 years of age** take 5 mL every 12 hours; do not take more than 10 mL in 24 hours ■ **children under 2 years of age** ask a doctor ■ **adults 65 years of age and older** ask a doctor

■ **consumers with kidney disease** ask a doctor.  
Note: mL = milliliters

**Other information**  
■ each 5 mL contains: **potassium 5 mg and sodium 19 mg** ■ safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ color change from white to beige is expected over time. This does not impact product effectiveness and safety.

**Inactive ingredients** edetate disodium, flavor, poloxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

**Questions or comments?** call 1-866-844-2798

Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219  
The trade dress of this Allegra® package is subject to trademark protection. ©2022 Origin Germany

845643-01  
LOT  
EXP

873286

**Children's Allegra<sup>®</sup>**  
**ALLERGY**  
**12 HR**  
NON-DROWSY

**Children's Allegra<sup>®</sup>**  
**ALLERGY**  
**12 HR**  
Ages 2 Years & Older



**Berry**  
FLAVOR

**FREE OF:**  
ALCOHOL  
& DYES

Use Only With  
Enclosed Dosing Cup  
Wash and let air dry after each use

**Children's Allegra<sup>®</sup>**  
**ALLERGY**  
**12 HR**  
NON-DROWSY

**VALUE SIZE!**  
Compare to 4 ounce

**2 Years & Older**

NDC 41167-4244-1

**Alcohol & Dye FREE**



**Children's Allegra<sup>®</sup>**  
**ALLERGY**  
**12 HR**  
Ages 2 Years & Older

fexofenadine HCl oral suspension  
30 mg/5 mL antihistamine

**NON-DROWSY**  
**Liquid**

**12 HR**  
INDOOR / OUTDOOR  
ALLERGY RELIEF

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat

**8 fl. oz. (240 mL)**



**Berry**  
FLAVOR

Use Only With  
Enclosed Dosing Cup  
Wash and let air dry after each use

**Children's Allegra<sup>®</sup>**  
**ALLERGY**  
**12 HR**  
NON-DROWSY

**Children's Allegra<sup>®</sup>**  
**ALLERGY**  
**12 HR**  
Ages 2 Years & Older



**Berry**  
FLAVOR

**FREE OF:**  
ALCOHOL  
& DYES

Use Only With  
Enclosed Dosing Cup  
Wash and let air dry after each use

**Drug Facts**

Active ingredient (in each 5 mL)	Purpose
Fexofenadine HCl 30 mg	Antihistamine

**Uses**  
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**  
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.  
Ask a doctor before use if you have kidney disease. Your doctor should determine if you need a different dose.

**When using this product**

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ shake well before using  
■ use only with enclosed dosing cup

adults and children 12 years of age and over	take 10 mL every 12 hours; do not take more than 20 mL in 24 hours
children 2 to under 12 years of age	take 5 mL every 12 hours; do not take more than 10 mL in 24 hours
children under 2 years of age	ask a doctor
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Note: mL = milliliters

**Other information**

- each 5 mL contains: potassium 5 mg and sodium 19 mg
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- color change from white to beige is expected over time. This does not impact product effectiveness and safety.

**Inactive ingredients**  
edetate disodium, flavor, poloxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

**Questions or comments?**  
call 1-866-844-2798

The makers of Allegra<sup>®</sup> do not make store brand products.  
The trade dress of this Allegra<sup>®</sup> package is subject to trademark protection.  
Dist. By: Chatterm, Inc., a Sanofi Company  
Chattanooga, TN 37409-0219 ©2022 Origin Germany

873286



0 41167 42441 4

LOT:  
EXP:



873286

**Children's Allegra Allergy 12 Hour 8oz Berry Liquid**  
**Liquid 8oz Carton**  
**12.01.22**

Allegra\_Children's\_Allergy\_12hr\_Berry\_8oz\_CTN\_Germany\_FDA\_873286\_DSN.ai  
(PDF of art being shown is submitted at 100% actual size)

# Children's Allegra Allergy 12 Hour 8 oz Berry Flavor Liquid

## Liquid 8 oz Bottle Label

05.21.22


Allegra\_Children's\_Allergy\_12hr\_Berry\_8oz\_Lbl\_845645-01\_DSN3.ai

(PDF of art being shown is submitted at 100% actual size)

NDC 41167-4244-1

**NON-DROWSY**  
2 Years & Older

Alcohol & Dye  
**FREE**



**Children's Allegra**  
**ALLERGY**

fexofenadine HCl oral suspension  
30 mg/5 mL antihistamine

**12 HR**  
**Berry**  
FLAVOR

**INDOOR / OUTDOOR ALLERGY RELIEF**

Dosing Cup Included  
Wash & let air dry after each use

8 fl. oz. (240 mL)

**Active ingredient (in each 5 mL)**  
Fexofenadine HCl 30 mg .....Antihistamine

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  
■ runny nose ■ itchy, watery eyes ■ sneezing  
■ itching of the nose or throat

**Warnings Do not use** if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions)


**Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** ■ shake well before using ■ use only with enclosed dosing cup ■ **adults and children 12 years of age and over** take 10 mL every 12 hours; do not take more

**Purpose** than 20 mL in 24 hours ■ **children 2 to under 12 years of age** take 5 mL every 12 hours; do not take more than 10 mL in 24 hours ■ **children under 2 years of age** ask a doctor ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor. Note: mL = milliliters

**Other information**  
■ each 5 mL contains: **potassium 5 mg and sodium 19 mg** ■ safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ color change from white to beige is expected over time. This does not impact product effectiveness and safety.

**Inactive ingredients** edetate disodium, flavor, poloxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

**Questions or comments?** call 1-866-844-2798



845645-01

LOT  
EXP

Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2022  
The trade dress of this Allegra® package is subject to trademark protection. Origin Germany



Children's Allegra Hives 12 Hour 4oz Berry Liquid  
Liquid 4oz Carton  
12.01.22

Allegra\_Children's\_Hives\_12hr\_Berry\_4oz\_CTN\_845652-01\_DSN8.ai  
(PDF of art being shown is submitted at 100% actual size)

# Children's Allegra Hives 12 Hour 4 oz Berry Flavor Liquid

## Liquid 4 oz Bottle Label

06.02.22

Allegra\_Children's\_Hives\_12hr\_Berry\_4oz\_Lbl\_845653-01\_DSN4.ai

(PDF of art being shown is submitted at 100% actual size)

NDC 41167-4226-1

Alcohol & Dye FREE

NON-DROWSY  
6 Years & Older

Children's Allegra Hives

fexofenadine HCl oral suspension 12 HR  
30 mg/5 mL antihistamine

REDUCES HIVES and RELIEVES ITCHING due to hives

Dosing Cup Included  
Wash & let air dry after each use

Berry FLAVOR

4 fl. oz. (120 mL)

**Active ingredient (in each 5 mL)**  
Fexofenadine HCl 30 mg ..... Antihistamine

**Purpose**  
Antihistamine

**Uses** reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

**Warnings** **Severe Allergy Warning:** Get emergency help immediately if you have hives along with any of the following symptoms: ■ swelling of tongue ■ trouble swallowing ■ swelling in or around mouth ■ trouble speaking ■ wheezing or problems breathing ■ drooling ■ dizziness or loss of consciousness These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. **Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times. **Do not use** ■ to prevent hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause. ■ if you have ever had an allergic reaction to this product or any of its ingredients **Ask a doctor before use if you have** ■ kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch **When using this** ▶

**product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** ■ an allergic reaction to this product occurs. Seek medical help right away. ■ symptoms do not improve after 3 days of treatment ■ the hives have lasted more than 6 weeks **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** ■ shake well before using ■ use only with enclosed dosing cup ■ adults and children 12 years of age and over take 10 mL every 12 hours; do not take more than 20 mL in 24 hours ■ children 6 to under 12 years of age take 5 mL every 12 hours; do not take more than 10 mL in 24 hours ■ children under 6 years of age ask a doctor ■ adults 65 years of age and older ask a doctor ■ consumers with kidney disease ask a doctor. Note: mL = milliliters **Other information** ■ each 5 mL contains: potassium 5 mg and sodium 19 mg ■ safety sealed; do not use if carton is opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ color change from white to beige is expected over time. This does not impact product effectiveness and safety. **Inactive ingredients** edetate disodium, flavor, polyoxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol **Questions or comments?** call 1-866-844-2798

Dist. By: Chatterm, Inc., a Sanofi Company, Chattanooga, TN 37409-0219  
The trade dress of this Allegra® package is subject to trademark protection. ©2022 Origin Germany

845653-01  
LOT  
EXP

**Children's Allegra Hives**  
12 HR  
NON-DROWSY

---

**Children's Allegra Hives**  
12 HR

**Berry**  
FLAVOR

**FREE OF:  
ALCOHOL  
& DYES**

Use Only With  
Enclosed Dosing Cup  
Wash and let air dry after each use

**Children's Allegra Hives**  
12 HR  
NON-DROWSY

**6 Years & Older**

**Alcohol & Dye FREE**

**Children's Allegra Hives**  
fexofenadine HCl oral suspension  
30 mg/5 mL antihistamine

**12 HOUR  
NON-DROWSY  
Liquid**

**REDUCES HIVES and  
RELIEVES ITCHING  
due to hives**

**8 fl. oz. (240 mL)**

**Berry**  
FLAVOR

**Children's Allegra Hives**  
12 HR

**Berry**  
FLAVOR

Use Only With Enclosed Dosing Cup  
Wash and let air dry after each use

845654-01



0 41167 42262 5

LOT:  
EXP:

**Drug Facts**

**Active ingredient (in each 5 mL)** Purpose  
Fexofenadine HCl 30 mg ..... Antihistamine

**Uses**  
reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

**Warnings**  
**Severe Allergy Warning:** Get emergency help immediately if you have hives along with any of the following symptoms:  

- swelling of tongue
- wheezing or problems breathing
- dizziness or loss of consciousness
- trouble swallowing
- trouble speaking
- drooling

 These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

**Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

**Do not use**

- to prevent hives from any known cause such as:
  - foods
  - insect stings
  - medicines
  - latex or rubber gloves
- hives that are an unusual color, look bruised or blistered
- hives that do not itch

**When using this product**

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.  
Distributed By: Chatterm, Inc., a Sanofi Company  
Chattanooga, TN 37409-0219 ©2022 Origin Germany

**Drug Facts (continued)**

**Directions**

adults and children 12 years of age and over	take 10 mL every 12 hours; do not take more than 20 mL in 24 hours
children 6 to under 12 years of age	take 5 mL every 12 hours; do not take more than 10 mL in 24 hours
children under 6 years of age	ask a doctor
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Note: mL = milliliters

**Other information**

- each 5 mL contains: potassium 5 mg and sodium 19 mg
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- color change from white to beige is expected over time. This does not impact product effectiveness and safety.

**Inactive ingredients**  
edetate disodium, flavor, poloxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

**Questions or comments?**  
call 1-866-844-2798

**Children's Allegra Hives 12 Hour 8oz Berry Liquid  
Liquid 8oz Carton  
12.01.22**

Allegra\_Children's\_Hives\_12hr\_Berry\_8oz\_Ctn\_845654-01\_DSN10.ai  
(PDF of art being shown is submitted at 100% actual size)

# Children's Allegra Hives 12 Hour 8 oz Berry Flavor Liquid

## Liquid 8 oz Bottle Label

### 06.04.22

Allegra\_Children's\_Hives\_12hr\_Berry\_8oz\_Lbl\_845640-01\_DSN4.ai  
(PDF of art being shown is submitted at 100% actual size)

NDC 41167-4226-2

**Alcohol & Dye FREE**

**NON-DROWSY**  
**6 Years & Older**

**Children's Allegra<sup>®</sup> Hives**  
fexofenadine HCl oral suspension  
**30 mg/5 mL antihistamine**

**12 HR**  
**Berry FLAVOR**

**REDUCES HIVES and RELIEVES ITCHING due to hives**

Dosing Cup Included  
Wash & let air dry after each use

**8 fl. oz. (240 mL)**

**Active ingredient (in each 5 mL)**  
Fexofenadine HCl 30 mg

**Purpose**  
Antihistamine

**Uses** reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

**Warnings Severe Allergy Warning:** Get emergency help **immediately** if you have hives along with any of the following symptoms: ■ swelling of tongue ■ trouble swallowing ■ swelling in or around mouth ■ trouble speaking ■ wheezing or problems breathing ■ drooling ■ dizziness or loss of consciousness. These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. **Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times. **Do not use** ■ to prevent hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause. ■ if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** ■ kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch. **When using this product** ■ do not take more than directed ■ do not take at

the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** ■ an allergic reaction to this product occurs. Seek medical help right away. ■ symptoms do not improve after 3 days of treatment ■ the hives have lasted more than 6 weeks. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** ■ shake well before using ■ use only with enclosed dosing cup ■ **adults and children 12 years of age and over** take 10 mL every 12 hours; do not take more than 20 mL in 24 hours ■ **children 6 to under 12 years of age** take 5 mL every 12 hours; do not take more than 10 mL in 24 hours ■ **children under 6 years of age** ask a doctor ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor. Note: mL = milliliters. **Other information** ■ each 5 mL contains: **potassium 5 mg and sodium 19 mg** ■ safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ color change from white to beige is expected over time. This does not impact product effectiveness and safety. **Inactive ingredients**

Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219  
The trade dress of this Allegra<sup>®</sup> package is subject to trademark protection. ©2022 Origin Germany

845640-01  
LOT  
EXP

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

NUSHIN F TODD  
12/16/2022 05:01:26 PM