



Children's Allegra Allergy 12 Hour 4oz Grape Liquid
Liquid 4oz Carton
12.01.22

Allegra_Children's_Allergy_12hr_Grape_4oz_CTN_Germany_FDA_873295_DSN.ai
(PDF of art being shown is submitted at 100% actual size)

Children's Allegra Allergy 12 Hour 4 oz Grape Flavor Liquid

Liquid 4 oz Bottle Label

06.02.22

Allegra_Children's_Allergy_12hr_Grape_4oz_Wrap_845638-01_DSN5.ai
(PDF of art being shown is submitted at 100% actual size)

NDC 41167-4246-4

NON-DROWSY
2 Years & Older

Alcohol & Dye **FREE**

Children's Allegra
ALLERGY

fexofenadine HCl oral suspension
30 mg/5 mL antihistamine

12 HR Grape
FLAVOR

INDOOR / OUTDOOR ALLERGY RELIEF

Dosing Cup Included
Wash & let air dry after each use

4 fl. oz. (120 mL)

Active ingredient (in each 5 mL)
Fexofenadine HCl 30 mg.....Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat

Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** ■ shake well before using ■ use only with enclosed dosing cup ■ **adults and children 12 years of age and over** take 10 mL every 12 hours; do not take more

Purpose than 20 mL in 24 hours ■ **children 2 to under 12 years of age** take 5 mL every 12 hours; do not take more than 10 mL in 24 hours ■ **children under 2 years of age** ask a doctor ■ **adults 65 years of age and older** ask a doctor

■ **consumers with kidney disease** ask a doctor. Note: mL = milliliters

Other information
■ each 5 mL contains: **potassium 5 mg and sodium 19 mg** ■ safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ color change from white to beige is expected over time. This does not impact product effectiveness and safety.

Inactive ingredients edetate disodium, flavor, poloxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

Questions or comments? call 1-866-844-2798

Dist. By: Chatterm, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
The trade dress of this Allegra® package is subject to trademark protection. ©2022 Origin Germany

845638-01
LOT
EXP

873298



Children's Allegra Allergy 12 Hour 8oz Grape Liquid
Liquid 8oz Carton
12.01.22

Allegra_Children's_Allergy_12hr_Grape_8oz_CTN_Germany_FDA_873298_DSN2.ai
(PDF of art being shown is submitted at 100% actual size)

Children's Allegra Allergy 12 Hour 8 oz Grape Flavor Liquid

Liquid 8 oz Bottle Label

05.21.22

Allegra_Children's_Allergy_12hr_Grape_8oz_Lbl_845640-01_DSN4.ai
(PDF of art being shown is submitted at 100% actual size)

NDC 41167-4246-2

Alcohol & Dye FREE

NON-DROWSY
2 Years & Older

Children's Allegra[®]
ALLERGY

fexofenadine HCl oral suspension
30 mg/5 mL antihistamine

12 HR
Grape
FLAVOR

INDOOR / OUTDOOR ALLERGY RELIEF

Dosing Cup Included
Wash & let air dry after each use

8 fl. oz. (240 mL)

Active ingredient (in each 5 mL)
Fexofenadine HCl 30 mgAntihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose ■ itchy, watery eyes ■ sneezing
■ itching of the nose or throat

Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions)

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** ■ shake well before using ■ use only with enclosed dosing cup ■ **adults and children 12 years of age and over** take 10 mL every 12 hours; do not take more

Purpose than 20 mL in 24 hours ■ **children 2 to under 12 years of age** take 5 mL every 12 hours; do not take more than 10 mL in 24 hours
■ **children under 2 years of age** ask a doctor
■ **adults 65 years of age and older** ask a doctor
■ **consumers with kidney disease** ask a doctor.
Note: mL = milliliters **Other information**
■ each 5 mL contains: **potassium 5 mg and sodium 19 mg** ■ safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ color change from white to beige is expected over time. This does not impact product effectiveness and safety.

Inactive ingredients edetate disodium, flavor, poloxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

Questions or comments? call 1-866-844-2798

Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2022
The trade dress of this Allegra[®] package is subject to trademark protection. Origin Germany

845640-01
LOT
EXP

Ages 6 Years & Older

Children's Allegra Hives

12 HR



Grape FLAVOR

**FREE OF:
ALCOHOL
& DYES**

Use Only With
Enclosed Dosing Cup
Wash and let air dry after each use

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.

Dist. By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 ©2022
Origin Germany

Children's Allegra Hives

12 HR
NON-DROWSY

NDC 41167-4227-0

6 Years & Older

Alcohol & Dye FREE



Children's Allegra Hives

fexofenadine HCl oral suspension
30 mg/5 mL antihistamine

**12 HOUR
NON-DROWSY
Liquid**

**REDUCES HIVES and
RELIEVES ITCHING
due to hives**



Grape FLAVOR

4 fl. oz. (120 mL)



Drug Facts

Active ingredient (in each 5 mL)	Purpose
Fexofenadine HCl 30 mg	Antihistamine

Uses
reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings
Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:

- swelling in or around mouth
- trouble swallowing
- wheezing or problems breathing
- drooling
- dizziness or loss of consciousness

 These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

Do not use

- to prevent hives from any known cause such as:
 - foods
 - insect stings
 - medicines
 - latex or rubber gloves

 because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch

Drug Facts (continued)

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- shake well before using
- use only with enclosed dosing cup

adults and children 12 years of age and over	take 10 mL every 12 hours; do not take more than 20 mL in 24 hours
children 6 to under 12 years of age	take 5 mL every 12 hours; do not take more than 10 mL in 24 hours
children under 6 years of age	ask a doctor
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Note: mL = milliliters

Other information

- each 5 mL contains: potassium 5 mg and sodium 19 mg
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- color change from white to beige is expected over time. This does not impact product effectiveness and safety.

Inactive ingredients
 edetate disodium, flavor, poloxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

Questions or comments? call 1-866-844-2798

845647-01



0 41167 42270 0

LOT:
EXP:



**Children's Allegra Hives 12 Hour 4oz Grape Liquid
Liquid 4oz Carton
12.01.22**

Allegra_Children's_Hives_12hr_Grape_4oz_CTN_845647-01_DSN6.ai
(PDF of art being shown is submitted at 100% actual size)

Children's Allegra Hives 12 Hour 4 oz Grape Flavor Liquid Liquid 4 oz Bottle Label

06.02.22

Allegra_Children's_Hives_12hr_Grape_4oz_Lbl_845648-01_DSN4.ai
(PDF of art being shown is submitted at 100% actual size)

Alcohol & Dye FREE

NDC 41167-4227-0

NON-DROWSY

6 Years & Older

Children's Allegra Hives

fexofenadine HCl oral suspension **12 HR**

30 mg/5 mL antihistamine

Grape FLAVOR

4 fl. oz. (120 mL)

REDUCES HIVES and RELIEVES ITCHING due to hives

Dosing Cup Included
Wash & let air dry after each use

Active ingredient (in each 5 mL)
Fexofenadine HCl 30 mg

Purpose
Antihistamine

Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings **Severe Allergy Warning:** Get emergency help immediately if you have hives along with any of the following symptoms: ■ swelling of tongue ■ trouble swallowing ■ swelling in or around mouth ■ trouble speaking ■ wheezing or problems breathing ■ drooling ■ dizziness or loss of consciousness. These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. **Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times. **Do not use** ■ to prevent hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause. ■ if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** ■ kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch. **When using this**

product ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** ■ an allergic reaction to this product occurs. Seek medical help right away. ■ symptoms do not improve after 3 days of treatment. ■ the hives have lasted more than 6 weeks. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** ■ shake well before using ■ use only with enclosed dosing cup ■ **adults and children 12 years of age and over** take 10 mL every 12 hours; do not take more than 20 mL in 24 hours ■ **children 6 to under 12 years of age** take 5 mL every 12 hours; do not take more than 10 mL in 24 hours ■ **children under 6 years of age** ask a doctor ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor. Note: mL = milliliters. **Other information** ■ each 5 mL contains: potassium 5 mg and sodium 19 mg ■ safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ color change from white to beige is expected over time. This does not impact product effectiveness and safety. **Inactive ingredients** edetate disodium, flavox, poloxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol. **Questions or comments?** call 1-866-844-2798

Dist. By: Chatterm, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
The trade dress of this Allegra® package is subject to trademark protection. ©2022 Origin Germany

845648-01
LOT
EXP

Children's Allegra Hives
12 HR
NON-DROWSY

Ages 6 Years & Older

Children's Allegra Hives
12 HR



Grape
FLAVOR

FREE OF:
ALCOHOL
& DYES

Use Only With
Enclosed Dosing Cup
Wash and let air dry after each use

845650-01

Children's Allegra Hives
12 HR
NON-DROWSY

NDC 41167-4227-1
6 Years & Older

Alcohol & Dye FREE

Children's Allegra Hives
fexofenadine HCl oral suspension
30 mg/5 mL antihistamine

12 HOUR NON-DROWSY Liquid

REDUCES HIVES and RELIEVES ITCHING due to hives

8 fl. oz. (240 mL)

Grape
FLAVOR

LOT:
EXP:

Children's Allegra Hives
12 HR

Grape
FLAVOR

Use Only With Enclosed Dosing Cup
Wash and let air dry after each use

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Fexofenadine HCl 30 mg	Antihistamine

Uses
reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings
Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:

- swelling of tongue ■ trouble swallowing
- swelling in or around mouth ■ trouble speaking
- wheezing or problems breathing ■ drooling
- dizziness or loss of consciousness

 These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

Do not use

- to prevent hives from any known cause such as:
- foods ■ insect stings ■ medicines ■ latex or rubber gloves

 because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause.

■ If you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have

- kidney disease. Your doctor should determine if you need a different dose.
- hives that are an unusual color, look bruised or blistered
- hives that do not itch

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

The makers of Allegra® do not make store brand products. The trade dress of this Allegra® package is subject to trademark protection.

Distributed By: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219
©2022 Origin Germany

Drug Facts (continued)

Directions

- shake well before using
- use only with enclosed dosing cup

adults and children 12 years of age and over	take 10 mL every 12 hours; do not take more than 20 mL in 24 hours
children 6 to under 12 years of age	take 5 mL every 12 hours; do not take more than 10 mL in 24 hours
children under 6 years of age	ask a doctor
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

Note: mL = milliliters

Other information

- each 5 mL contains: potassium 5 mg and sodium 19 mg
- safety sealed: do not use if carton is opened or if printed foil inner seal on bottle is torn or missing
- store between 20° and 25°C (68° and 77°F)
- color change from white to beige is expected over time. This does not impact product effectiveness and safety.

Inactive ingredients
edetate disodium, flavor, poloxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

Questions or comments?
call 1-866-844-2798

**Children's Allegra Hives 12 Hour 8oz Grape Liquid
Liquid 8oz Carton
12.01.22**

Allegra_Children's_Hives_12hr_Grape_8oz_CTN_845650-01_DSN6.ai
(PDF of art being shown is submitted at 100% actual size)

Children's Allegra Hives 12 Hour 8 oz Grape Flavor Liquid

Liquid 8 oz Bottle Label

06.04.22

Allegra_Children's_Hives_12hr_Grape_8oz_Lbl_845651-01_DSN4.ai
(PDF of art being shown is submitted at 100% actual size)

NDC 41167-4227-1

Alcohol & Dye FREE

NON-DROWSY
6 Years & Older

Children's Allegra[®] Hives
fexofenadine HCl oral suspension
30 mg/5 mL antihistamine

12 HR
Grape FLAVOR

REDUCES HIVES and RELIEVES ITCHING due to hives

Dosing Cup Included
Wash & let air dry after each use

8 fl. oz. (240 mL)

Active ingredient (in each 5 mL)
Fexofenadine HCl 30 mg.....Antihistamine
Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring. **Warnings Severe Allergy Warning:** Get emergency help **immediately** if you have hives along with any of the following symptoms: ■ swelling of tongue ■ trouble swallowing ■ swelling in or around mouth ■ trouble speaking ■ wheezing or problems breathing ■ drooling ■ dizziness or loss of consciousness. These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional **immediately**. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. **Not a Substitute for Epinephrine.** If your doctor has prescribed an epinephrine injector for "anaphylaxis" or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times. **Do not use** ■ to prevent hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious. If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause. ■ if you have ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** ■ kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch. **When using this product** ■ do not take more than directed ■ do not take at

Purpose the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor if** ■ an allergic reaction to this product occurs. Seek medical help right away. ■ symptoms do not improve after 3 days of treatment. ■ the hives have lasted more than 6 weeks. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** ■ shake well before using ■ use only with enclosed dosing cup ■ **adults and children 12 years of age and over** take 10 mL every 12 hours; do not take more than 20 mL in 24 hours ■ **children 6 to under 12 years of age** take 5 mL every 12 hours; do not take more than 10 mL in 24 hours ■ **children under 6 years of age** ask a doctor ■ **adults 65 years of age and older** ask a doctor ■ **consumers with kidney disease** ask a doctor. **Note:** mL = milliliters. **Other information** ■ each 5 mL contains: **potassium 5 mg and sodium 19 mg** ■ safety sealed; do not use if carton is opened or if printed foil inner seal on bottle is torn or missing ■ store between 20° and 25°C (68° and 77°F) ■ color change from white to beige is expected over time. This does not impact product effectiveness and safety. **Inactive ingredients** edetate disodium, flavor, poloxamer 407, potassium sorbate, propylene glycol, purified water, sodium phosphate dibasic heptahydrate, sodium phosphate monobasic monohydrate, sucrose, titanium dioxide, xanthan gum, xylitol

Questions or comments?
call 1-866-844-2798

Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
The trade dress of this Allegra[®] package is subject to trademark protection. ©2022 Origin Germany

845651-01

LOT
EXP

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
12/27/2022 03:23:45 PM