

BACK




SEAL AREA FOR LOT & EXP
25 mm x 4.5 mm

AREA FOR LOT AND EXP
TO BE PRINTED IN DARK COLOR
26 mm x 5.5 mm

000068426

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	N/A



READ AND KEEP CARD FOR COMPLETE WARNINGS AND INFORMATION

Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or licensor
Trademarks owned or licensed by GSK
For most recent product information, visit www.Advil.com
Patents: www.productpats.com


FPO
3 0573 0133 09 8

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Ibuprofen 200 mg (provided as Ibuprofen sodium 256 mg) (NSAID)* Pain reliever/ *nonsteroidal anti-inflammatory drug Fever reducer</p> <p>Uses temporarily relieves minor aches and pains due to: headache toothache backache menstrual cramps the common cold muscular aches minor pain of arthritis temporarily reduces fever</p> <p>Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery</p> <p>Ask a doctor before use if stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke you are taking a diuretic</p> <p>Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug</p>	<p>Drug Facts (continued)</p> <p>When using this product take with food or milk if stomach upset occurs</p> <p>Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better you have symptoms of heart problems or stroke: chest pain trouble breathing weakness in one part or side of body slurred speech leg swelling pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions do not take more than directed the smallest effective dose should be used adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist if pain or fever does not respond to 1 tablet, 2 tablets may be used do not exceed 6 tablets in 24 hours, unless directed by a doctor children under 12 years: ask a doctor</p> <p>Other information each tablet contains: sodium 22 mg read all warnings and directions before use. Keep card. store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F)</p> <p>Inactive ingredients acesulfame potassium, caramel color, carnauba wax, colloidal silicon dioxide, copovidone, ferric oxide, hypromellose, mannitol, medium-chain triglycerides, microcrystalline cellulose, natural and artificial flavor, pharmaceutical ink, polyethylene glycol, propylene glycol, sodium lauryl sulfate, sucralose, titanium dioxide</p> <p>Questions or comments? Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL</p> <p>000068429</p>
--	--

↑
LOT & EXP AREA

000068429

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



000068428

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (provided as Pain reliever/
ibuprofen sodium 256 mg) (NSAID)*Fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
 - temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
 - adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
 - children under 12 years: ask a doctor

Other information

- each tablet contains: sodium 22 mg
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

acesulfame potassium, caramel color, carnauba wax, colloidal silicon dioxide, copovidone, ferric oxide, hypromellose, mannitol, medium-chain triglycerides, microcrystalline cellulose, natural and artificial flavor, pharmaceutical ink, polyethylene glycol, propylene glycol, sodium lauryl sulfate, sucralose, titanium dioxide

Questions or comments?

Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

For most recent product information, visit www.Advil.com
Dist. by: GSK CH, Warren, NJ 07059
©2021 GSK or licensor
Trademarks owned or licensed by GSK
Patents: www.productpats.com

000072134



UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

000072134

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if the sealed child resistant pouch is broken or torn.

Drug Facts

Active Ingredients (in each tablet): Ibuprofen sodium 200 mg (provided as Ibuprofen Sodium 256 mg) and sodium salicylate 56 mg (provided as Sodium Salicylate 71.68 mg).

Purpose: Pain reliever, fever reducer.

Uses: Ibuprofen sodium 200 mg (NSAID) and sodium salicylate 56 mg (NSAID) are used to relieve pain and reduce fever.

Warnings: Allergy Alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, difficulty breathing, swelling of the face, lips, tongue, or throat. Stop use and seek medical help right away. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach Bleeding Warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: take a blood thinning (anticoagulant) or steroid drug; take a blood thinning (anticoagulant) or steroid drug; take aspirin, ibuprofen, naproxen, or other NSAID; have 3 or more alcoholic drinks every day while using this product; have a history of stomach problems, such as heartburn, ulcers, or bleeding from taking pain relievers; or have a history of stomach problems, such as heartburn, ulcers, or bleeding from taking pain relievers.

Directions: Adults and children 12 years and over: Take 1 tablet every 4 to 6 hours. The smallest effective dose should be used. Children under 12 years: Ask a doctor. Do not use for more than 10 days unless directed by a doctor. Do not use for more than 10 days unless directed by a doctor. Do not use for more than 10 days unless directed by a doctor.

Other Information: Each label contains sodium 22 mg. Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).

Questions or Comments? Call 1-800-842-ADVIL. For more recent product information, visit www.Advii.com. For US Patient or Application status see www.products.fda.com. Distributed by: GSK Consumer Healthcare, Warren, NJ 07059. Trademarks are owned by or licensed to the GSK group of companies. ©2021 GSK group of companies or its licensor.

Advil FILM-COATED IBUPROFEN SODIUM

Ibuprofen Tablets, 200 mg (Provided as Ibuprofen Sodium 256 mg) Pain Reliever / Fever Reducer (NSAID)

Advil Tablets

Advil FILM-COATED IBUPROFEN SODIUM

Ibuprofen Tablets, 200 mg (Provided as Ibuprofen Sodium 256 mg) Pain Reliever / Fever Reducer (NSAID)

Advil Tablets

Advil FILM-COATED IBUPROFEN SODIUM

Ibuprofen Tablets, 200 mg (Provided as Ibuprofen Sodium 256 mg) Pain Reliever / Fever Reducer (NSAID)

Advil Tablets

Headache
 Muscular Aches
 Minor Pain of Arthritis
 Menstrual Cramps
 Toothache
 Fever

See Full Drug Facts information on the back of this dispenser

100 TABLETS (50 PACKETS of 2 TABLETS EACH)

000068427

FPO

3 0573 0133 02 9

LASER ETCHING AREA FOR LOT & EXP. PRINT A SOLID DARK COLOR.

000068427

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	7 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, 5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



WHITE AREA HOLD OUT FOR
LOT CODE & EXPIRATION DATE
(NO VARNISH)

000072137

DRUG FACTS TEXT DEFINED	TYPE SIZE	DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt	• # OF CHARACTERS PER INCH	<39
• DRUG FACTS CONTINUED	8 pt	• BULLETS	5 pt
• HEADINGS	8 pt	• SPACE BEFORE BULLET	2 ems
• SUBHEADINGS/BODY TEXT	6 pt	• BARLINES, HAIRLINES	1.5 pt, .5 pt
• LEADING	6.5 pt	• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

BACK



SEAL AREA FOR LOT & EXP
25 mm x 4.5 mm

AREA FOR LOT AND EXP
TO BE PRINTED IN DARK COLOR
26 mm x 5.5 mm

NPA006058

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	N/A

Drug Facts
Active ingredient (in each tablet)
Ibuprofen 200 mg (provided as ibuprofen sodium 256 mg) (NSAID)* ... Fever reducer
Purpose
Pain reliever/
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - backache
 - headache
 - muscular aches
 - the common cold
 - toothache
 - minor pain of arthritis
- temporarily reduces fever

Drug Facts (continued)
Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

NPA006090

LIFT HERE

3/16" x 3/16" DATA MATRIX CODE (ECC 200), PRINTED IN A DARK COLOR ON UNPRINTED WHITE BACKGROUND

PANEL 1

Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you

Drug Facts (continued)

- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

PANEL 2

Drug Facts (continued)
Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

BASE PANEL


NPA006090

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.4 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

FRONT



BACK



PRINTED AREA FOR ADVIL LOT & EXP

NPA006065

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do not use if seal under bottle cap imprinted with -SEALED BY YOUR PROTECTION- is broken or missing.
READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Active ingredient (in each tablet)
Ibuprofen 200 mg (provided as ibuprofen sodium 256 mg) (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose
Pain relief/fever reducer

Uses
■ temporarily relieves minor aches and pains due to:
■ menstrual cramps ■ backache ■ headache
■ muscular aches ■ the common cold ■ toothache
■ minor pain of arthritis ■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ skin redness ■ rash
■ dizziness ■ an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)
Stomach bleeding warning: This product contains an NSAID, which may cause serious stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or over-the-counter NSAIDs, aspirin, ibuprofen, naproxen, or other ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery.
Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic.
Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug.

Drug Facts (continued)
When using this product ■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ heartburn ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ fever ■ gets worse or lasts more than 10 days ■ fever ■ gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

For most recent product information, visit www.Advils.com
Dist. by: GSK CH, Warren, NJ 07059
©2021 GSK or licensee
Trademarks owned or licensed by GSK

FPO

3 0573 0230 40 7

Drug Facts (continued)
Other information
■ each tablet contains sodium 22 mg
■ read all warnings and directions before use; keep carton.
■ avoid excessive heat above 40°C (104°F)
Inactive ingredients: acetylsalicylic acid, potassium, caramel color, carnauba wax, colloidal silicon dioxide, copovidone, ferric oxide, hypromellose, mannitol, medium-chain triglycerides, microcrystalline cellulose, natural and artificial flavor, pharmaceutical ink, polyethylene glycol, propylene glycol, sodium lauryl sulfate, sucralose, titanium dioxide

Questions or comments?
Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL
NPA006057 Patents: www.productpatents.com

Advil
menstrual PAIN
Ibuprofen Tablets, 200 mg
(Provided as Ibuprofen Sodium 256 mg)
Pain Reliever/Fever Reducer (NSAID)
Advil msp **Tablets**
40 Tablets

NPA006057

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.3 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
02/08/2022 02:38:27 PM