

LOT: \_\_\_\_\_

EXP: \_\_\_\_\_

**IMPRINT AREA  
NO COPY**

**NO COPY AREA  
NO VARNISH**

**Quillivant XR<sup>®</sup>**  
methylphenidate HCl  
for extended-release oral suspension

**300 mg/60 mL total volume**  
(When reconstituted with 53 mL of water)

**25 mg/5 mL**   
(5 mg/mL) when reconstituted

**SHAKE WELL FOR AT LEAST  
10 SECONDS BEFORE EACH USE**

**Pharmacist Information**  
**Usual dosage:** Once daily. See package insert for dosage information.  
**Prepare suspension at time of dispensing as follows:**  
 Tip bottle until powder flows freely. Add 53 mL of water for reconstitution, insert bottle adapter into neck of bottle. Replace bottle cap. Shake with vigorous back and forth motion for at least 10 seconds to prepare suspension.  
**Dispense in original packaging (bottle in carton) with bottle adapter inserted and with enclosed oral dosing dispenser.**  
 Each 1 mL of reconstituted suspension contains 5 mg of Methylphenidate Hydrochloride. Each bottle contains 300 mg of Methylphenidate Hydrochloride.  
**Store at 25°C (77°F); excursions permitted from 15°C to 30°C (59°F to 86°F). [See USP controlled room temperature.]**

**Quillivant XR<sup>®</sup>**  
methylphenidate HCl  
for extended-release oral suspension

**300 mg/60 mL total volume**  
(When reconstituted with 53 mL of water)

**25 mg/5 mL**   
(5 mg/mL) when reconstituted

**SHAKE WELL FOR AT LEAST  
10 SECONDS BEFORE EACH USE**

**Pharmacist: Quillivant XR must be reconstituted with 53 mL of water prior to dispensing.**

KEEP OUT OF THE REACH OF CHILDREN

Distributed by  
**NextWave Pharmaceuticals, Inc.**  
 A subsidiary of Pfizer Inc., New York, NY 10007

**Manufactured by:**  
 Tris Pharma, Inc.  
 Monmouth Junction, NJ 08852

**LiquiXR<sup>®</sup>**  
ORAL DOSING DISPENSER

**Quillivant XR<sup>®</sup>**  
methylphenidate HCl  
for extended-release oral suspension

**300 mg/60 mL total volume**  
(When reconstituted with 53 mL of water)

**25 mg/5 mL**   
(5 mg/mL) when reconstituted

**SHAKE WELL FOR AT LEAST  
10 SECONDS BEFORE EACH USE**

**Patient Information**  
 Check and make sure that the Quillivant XR bottle contains liquid medicine. If Quillivant XR is in powder form, do not use it. Return it to your pharmacist.  
 Keep bottle tightly closed. Discard any unused portion after 120 days.  
**Instructions for Using Enclosed Oral Dosing Dispenser**

- Use only with the oral dosing dispenser provided with this product.
- Shake the bottle well (up and down) for at least 10 seconds before each use.
- Remove bottle cap.
- Confirm that bottle adapter has been inserted into top of the bottle.
- Insert tip of oral dosing dispenser provided with this product into bottle adapter.
- Turn bottle upside down and withdraw prescribed amount of QUILLIVANT XR into oral dosing dispenser.
- Remove filled oral dosing dispenser from bottle and dispense QUILLIVANT XR directly into mouth.
- Replace bottle cap and store bottle as directed.
- Wash oral dosing dispenser after each use (components are dishwasher-safe).

**NO COPY AREA  
NO VARNISH**

14340500

FPO - NDC/HRI

3 1

**24478-19010**

LB8315  
Rev 01  
09/14

FPO - 2D 20ML  
LB8315 Rev 01 09/14

Rx only

**NO COPY AREA  
NO VARNISH**

090177e1864db9c8\Approved\Approved On: 02-Apr-2015 13:30

(b) (4)

LOT: IMPRINT AREA  
 EXP: NO COPY

**Quillivant XR<sup>®</sup>**  
 methylphenidate HCl  
 for extended-release oral suspension

**600 mg/120 mL total volume**  
 (When reconstituted with 105 mL of water)

Pfizer NDC 24478-200-20  
 Pharmacist: Dispense the enclosed Medication Guide to each patient.

**Quillivant XR<sup>®</sup>**  
 methylphenidate HCl  
 for extended-release oral suspension

**600 mg/120 mL total volume**  
 (When reconstituted with 105 mL of water)

**25 mg/5 mL** (II)  
 (5 mg/mL) when reconstituted

**SHAKE WELL FOR AT LEAST 10 SECONDS BEFORE EACH USE**

**Pharmacist: Quillivant XR must be reconstituted with 105 mL of water prior to dispensing.**

KEEP OUT OF THE REACH OF CHILDREN


Distributed by:  
 Pfizer NextWave Pharmaceuticals, Inc.  
 A subsidiary of Pfizer Inc., New York, NY 10017

Manufactured by:  
 TriS Pharma, Inc.  
 Monmouth Junction, NJ 08852

**LiquiXR<sup>®</sup>**

Rx only

14340800

FPO - NDC/HRI  
  
 3 24478-20020 6

LB8316  
 Rev 01  
 09/14

FPO - 20 20ML  
 LB8316 Rev 01 09/14

NO COPY AREA  
 NO VARNISH

NO COPY AREA - NO VARNISH

NO COPY AREA  
 NO VARNISH

090177e1864db8ca Approved/Approved On: 02-Apr-2015 13:30



(b) (4)

LOT: IMPRINT AREA  
 EXP: NO COPY

**Quillivant XR®**  
 methylphenidate HCl  
 for extended-release oral suspension

**750 mg/150 mL total volume**  
 (When reconstituted with 131 mL of water)

Pfizer NDC 24478-205-25  
 Pharmacist: Dispense the enclosed Medication Guide to each patient.

**Quillivant XR®**  
 methylphenidate HCl  
 for extended-release oral suspension

**750 mg/150 mL total volume**  
 (When reconstituted with 131 mL of water)

**25 mg/5 mL**  
 (5 mg/mL) when reconstituted

SHAKE WELL FOR AT LEAST 10 SECONDS BEFORE EACH USE

Pharmacist Information  
 Usual dosage: Once daily. See package insert for dosage information.  
 Prepare suspension at time of dispensing as follows:  
 Tap bottle until powder flows freely. Add 131 mL of water for reconstitution. Insert bottle adapter into neck of bottle. Replace bottle cap. Shake with vigorous back and forth motion for at least 10 seconds to prepare suspension.  
 Dispense in original packaging (bottle in carton) with bottle adapter inserted and with enclosed oral dosing dispenser.  
 Each 1 mL of reconstituted suspension contains 5 mg of Methylphenidate Hydrochloride. Each bottle contains 750 mg of Methylphenidate Hydrochloride.  
 Store at 25°C (77°F); excursions permitted from 15°C to 30°C (59°F to 86°F). [See USP controlled room temperature.]

Pharmacist: Quillivant XR must be reconstituted with 131 mL of water prior to dispensing.

KEEP OUT OF THE REACH OF CHILDREN


Distributed by  
 Pfizer NextWave Pharmaceuticals, Inc.  
 A subsidiary of Pfizer Inc, New York, NY 10101

Manufactured by:  
 Trix Pharma, Inc.  
 Monmouth Junction, NJ 08852

LiquiXR™  
 TECHNOLOGY

Rx only

14340600

FPO - NDC/HRI  
  
 24478-20525 6

LB8317  
 Rev 01  
 09/14

FPO- 20 20ML  
 LB8317 Rev 01 09/14

NO COPY AREA  
 NO VARNISH

NO COPY AREA - NO VARNISH

090177e1864db9cc Approved On: 02-Apr-2015 13:30

(b) (4)

LOT: IMPRINT AREA  
EXP: NO COPY

**Quillivant XR®**  
methylphenidate HCl  
for extended-release oral suspension

900 mg/180 mL total volume  
(When reconstituted with 158 mL of water)

**Quillivant XR®**  
methylphenidate HCl  
for extended-release oral suspension

900 mg/180 mL total volume  
(When reconstituted with 158 mL of water)

**Quillivant XR®**  
methylphenidate HCl  
for extended-release oral suspension

900 mg/180 mL total volume  
(When reconstituted with 158 mL of water)

**Quillivant XR®**  
methylphenidate HCl  
for extended-release oral suspension

900 mg/180 mL total volume  
(When reconstituted with 158 mL of water)

**25 mg/5 mL** (5 mg/mL) when reconstituted  
**SHAKE WELL FOR AT LEAST 10 SECONDS BEFORE EACH USE**

**25 mg/5 mL** (5 mg/mL) when reconstituted  
**SHAKE WELL FOR AT LEAST 10 SECONDS BEFORE EACH USE**

**25 mg/5 mL** (5 mg/mL) when reconstituted  
**SHAKE WELL FOR AT LEAST 10 SECONDS BEFORE EACH USE**

**25 mg/5 mL** (5 mg/mL) when reconstituted  
**SHAKE WELL FOR AT LEAST 10 SECONDS BEFORE EACH USE**

**Pharmacist: Quillivant XR must be reconstituted with 158 mL of water prior to dispensing.**

KEEP OUT OF THE REACH OF CHILDREN

**Pharmacist Information**  
Usual dosage: Once daily. See package insert for dosage information.  
Prepare suspension at time of dispensing as follows:  
Tap bottle until powder flows freely. Add 158 mL of water for reconstitution. Insert bottle adapter into neck of bottle. Replace bottle cap. Shake with vigorous back and forth motion for at least 10 seconds to prepare suspension.  
Dispense in original packaging (bottle in carton) with bottle adapter inserted and with enclosed oral dosing dispenser.  
Each 1 mL of reconstituted suspension contains 5 mg of Methylphenidate Hydrochloride.  
Each bottle contains 900 mg of Methylphenidate Hydrochloride.  
Store at 25°C (77°F); excursions permitted from 15°C to 30°C (59°F to 86°F). [See USP controlled room temperature.]

**Instructions for Using Enclosed Oral Dosing Dispenser**

- Use only with the oral dosing dispenser provided with this product.
- Shake the bottle well (up and down) for at least 10 seconds before each use.
- Remove bottle cap.
- Confirm that bottle adapter has been inserted into top of the bottle.
- Insert tip of oral dosing dispenser provided with this product into bottle adapter.
- Turn bottle upside down and withdraw prescribed amount of QUILLIVANT XR into oral dosing dispenser.
- Remove filled oral dosing dispenser from bottle and dispense QUILLIVANT XR directly into mouth.
- Replace bottle cap and store bottle as directed.
- Wash oral dosing dispenser after each use (components are dishwasher-safe).

14340700

FPO - NDC/HRI  
3 24478-21030 4  
L88318 Rev 01 09/14  
FPO - 20 20ML L88318 Rev 01 09/14

Rx only

LIQUIXR TECHNOLOGY

NO COPY AREA NO VARNISH

NO COPY AREA NO VARNISH

NO COPY AREA - NO VARNISH

090177e1864db9ceApproved On: 02-Apr-2015 13:30

(b) (4)

090177e1864db9c7\Approved\Approved On: 02-Apr-2015 13:30

Pharmacist Information: Usual dosage: Once daily. See package insert for dosage information. Prepare suspension at time of dispensing as follows: Tap bottle until powder flows freely. Add 53 mL of water for reconstitution. Insert bottle adapter into neck of bottle. Replace bottle cap. Shake with vigorous back and forth motion for at least 10 seconds to prepare suspension. Dispense in original packaging (bottle in carton) with bottle adapter inserted and with enclosed oral dosing dispenser. Each 1 mL of reconstituted suspension contains 5 mg of Methylphenidate Hydrochloride. Each bottle contains 300 mg of Methylphenidate Hydrochloride. Store at 25°C (77°F); excursions permitted from 15°C to 30°C (59°F to 86°F). [See USP controlled room temperature.] LB8212 Rev 02 09/14

NDC 24478-190-10  
Pharmacist: Dispense the enclosed Medication Guide to each patient.

**Quillivant XR®**  
methylphenidate HCl  
for extended-release oral suspension

**300 mg/60 mL total volume**  
(When reconstituted with 53 mL of water)

**25 mg/5 mL**  
(5 mg/mL) when reconstituted

Shake well for at least 10 seconds before using.

LOT: [REDACTED]  
EXP: [REDACTED]

IMPRINT AREA reads this way NO VARNISH

FPO UPC (NDC4HRI) @ 100%  
24478-19010 1

LiquiXR®  
Rx only

14362400



090177e1864db9c9\Approved\Approved On: 02-Apr-2015 13:30

Pharmacist Information: Usual dosage: Once daily. See package insert for dosage information. Prepare suspension at time of dispensing as follows: Tap bottle until powder flows freely. Add 105 mL of water for reconstitution. Insert bottle adapter into neck of bottle. Replace bottle cap. Shake with vigorous back and forth motion for at least 10 seconds to prepare suspension. Dispense in original packaging (bottle in carton) with bottle adapter inserted and with enclosed oral dosing dispenser. Each 1 mL of reconstituted suspension contains 5 mg of Methylphenidate Hydrochloride. Each bottle contains 600 mg of Methylphenidate Hydrochloride. Store at 25°C (77°F); excursions permitted from 15°C to 30°C (59°F to 86°F). [See USP controlled room temperature.] LB8215 Rev 02 09/14

NDC 24478-200-20  
Pharmacist: Dispense the enclosed Medication Guide to each patient.

**Quillivant XR®**  
methylphenidate HCl  
for extended-release oral suspension

**600 mg/120 mL total volume**  
(When reconstituted with 105 mL of water)

**25 mg/5 mL**  
(5 mg/mL) when reconstituted

Shake well for at least 10 seconds before using.

LOT: [REDACTED]  
EXP: [REDACTED]

IMPRINT AREA reads this way NO VARNISH

FPO UPC (NDC4HRI) @ 100%  
24478-20020 6

LiquiXR®  
Rx only

14343700



090177e1864db9cb\Approved\Approved On: 02-Apr-2015 13:30

**Pharmacist Information:**  
**Usual dosage:** Once daily. See package insert for dosage information.  
**Prepare suspension at time of dispensing as follows:** Tap bottle until powder flows freely. **Add 131 mL of water for reconstitution.** Insert bottle adapter into neck of bottle. Replace bottle cap. **Shake with vigorous back and forth motion for at least 10 seconds to prepare suspension.**  
**Dispense in original packaging (bottle in carton) with bottle adapter inserted and with enclosed oral dosing dispenser.**  
Each 1 mL of reconstituted suspension contains 5 mg of Methylphenidate Hydrochloride. Each bottle contains 750 mg of Methylphenidate Hydrochloride. **Store at 25°C (77°F); excursions permitted from 15°C to 30°C (59°F to 86°F).** [See USP controlled room temperature.]  
LB8300 Rev 02 09/14

**Pfizer** Pharmacist: Dispense the enclosed Medication Guide to each patient.  
**NDC 24478-205-25**

**Quillivant XR®**  
**methylphenidate HCl**  
for extended-release oral suspension

**750 mg/150 mL total volume**  
(When reconstituted with 131 mL of water)

**25 mg/5 mL**  
(5 mg/mL) when reconstituted

Shake well for at least 10 seconds before using.

**LiquiXR™** Rx only

**Shake liquid well before using. If in powder form, do not use, return to Pharmacist.**  
Keep bottle tightly closed.  
Discard any unused portion of the reconstituted suspension after 120 days.  
**KEEP OUT OF THE REACH OF CHILDREN**  
**Distributed by:** NextWave Pharmaceuticals, Inc.  
A subsidiary of Pfizer Inc., New York, NY 10017  
**Manufactured by:** Tri-Tri Pharma, Inc.,  
Morris Plains, NJ 08852

LOT: **IMPRINT AREA reads this way**  
EXP: **NO VARNISH**

**FPO UPC (NDC/HRI) @ 100%**  
N 3 24478-20525 6

14345600



090177e1864db9cd\Approved\Approved On: 02-Apr-2015 13:30

**Pharmacist Information:**  
**Usual dosage:** Once daily. See package insert for dosage information.  
**Prepare suspension at time of dispensing as follows:** Tap bottle until powder flows freely. **Add 158 mL of water for reconstitution.** Insert bottle adapter into neck of bottle. Replace bottle cap. **Shake with vigorous back and forth motion for at least 10 seconds to prepare suspension.**  
**Dispense in original packaging (bottle in carton) with bottle adapter inserted and with enclosed oral dosing dispenser.**  
Each 1 mL of reconstituted suspension contains 5 mg of Methylphenidate Hydrochloride. Each bottle contains 900 mg of Methylphenidate Hydrochloride. **Store at 25°C (77°F); excursions permitted from 15°C to 30°C (59°F to 86°F).** [See USP controlled room temperature.]  
LB8218 Rev 02 09/14

**Pfizer** Pharmacist: Dispense the enclosed Medication Guide to each patient.  
**NDC 24478-210-30**

**Quillivant XR®**  
**methylphenidate HCl**  
for extended-release oral suspension

**900 mg/180 mL total volume**  
(When reconstituted with 158 mL of water)

**25 mg/5 mL**  
(5 mg/mL) when reconstituted

Shake well for at least 10 seconds before using.

**LiquiXR™** Rx only

**Shake liquid well before using. If in powder form, do not use, return to Pharmacist.**  
Keep bottle tightly closed.  
Discard any unused portion of the reconstituted suspension after 120 days.  
**KEEP OUT OF THE REACH OF CHILDREN**  
**Distributed by:** NextWave Pharmaceuticals, Inc.  
A subsidiary of Pfizer Inc., New York, NY 10017  
**Manufactured by:** Tri-Tri Pharma, Inc.,  
Morris Plains, NJ 08852

LOT: **IMPRINT AREA reads this way**  
EXP: **NO VARNISH**

**FPO UPC (NDC/HRI) @ 100%**  
N 3 24478-21030 4

14345600



---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/  
-----

MITCHELL V Mathis  
07/13/2015