

6.2 Postmarketing Experience
The following adverse reactions have been identified during post approval use of aripiprazole...

7. DRUG INTERACTIONS
7.1 Drugs Having Clinically Important Interactions with Aripiprazole
Table 16: Clinically Important Drug Interactions with Aripiprazole:

Table 16: Clinically Important Drug Interactions with Aripiprazole. Columns: Concomitant Drug Name or Drug Class, Clinical Rationale, Clinical Recommendation.

7.2 Drugs Having No Clinically Important Interactions with Aripiprazole
Based on pharmacokinetic studies, no dosage adjustment of aripiprazole is required when administered concomitantly with lamotrigine, valproate, lithium, lorazepam...

8. USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
Teratogenic Effects
Pregnancy Exposure Registry

8.2 Lactation
Aripiprazole is excreted in human breast milk. Because of the potential for serious adverse reactions in nursing infants from aripiprazole, a decision should be made whether to discontinue nursing or to discontinue the drug...

8.3 Nursing Mothers
Aripiprazole is present in human breast milk. Because of the potential for serious adverse reactions in nursing infants from aripiprazole, a decision should be made whether to discontinue nursing or to discontinue the drug...

8.4 Pediatric Use
The pharmacokinetics of aripiprazole and dehydro-aripiprazole in pediatric patients 10 to 17 years of age were similar to those in adults after correcting for the differences in body weight...

8.5 Geriatric Use
Safety and effectiveness in pediatric patients with schizophrenia were established in a 6-week, placebo-controlled clinical trial in 202 pediatric patients aged 13 to 17 years...

8.6 Hepatic Use
The pharmacokinetics of aripiprazole and dehydro-aripiprazole in pediatric patients 10 to 17 years of age were similar to those in adults after correcting for the differences in body weight...

8.7 Renal Use
The pharmacokinetics of aripiprazole and dehydro-aripiprazole in pediatric patients 10 to 17 years of age were similar to those in adults after correcting for the differences in body weight...

10.1 Human Experience
In clinical trials and in postmarketing experience, adverse reactions of deliberate or accidental overdose with aripiprazole have been reported worldwide. These include overdoses with oral aripiprazole alone and in combination with other substances...

10.2 Management of Overdose
No specific information is available on the treatment of overdose with aripiprazole. An electrocardiogram should be obtained in case of overdose and a QT interval prolongation is present...

10.3 Hemodialysis
Aripiprazole is not dialyzable. Hemodialysis is unlikely to be useful in overdose treatment since aripiprazole is highly bound to plasma proteins.

10.4 Description
Aripiprazole is a psychotropic drug that is available as Aripiprazole Tablets. Aripiprazole is 1-(4-(2,3-dichlorophenyl)-1-piperazinyl)butyl)-3,4-dihydroquinoline...

10.5 Pharmacodynamics
Aripiprazole exhibits high affinity for dopamine D1 and D2, serotonin 5-HT1A, and 5-HT2 receptors. It also has moderate affinity for dopamine D4, serotonin 5-HT2A, and 5-HT2B receptors...

10.6 Pharmacokinetics
Aripiprazole is primarily eliminated through hepatic metabolism involving two P450 isoenzymes, CYP2D6 and CYP3A4. For CYP2D6 poor metabolizers, the mean elimination half-life for aripiprazole is about 14 hours...

10.7 Oral Administration
Aripiprazole is well absorbed after administration of the tablet, with peak plasma concentrations occurring within 3 to 5 hours. The absolute oral bioavailability of the tablet formulation is 87%...

10.8 Distribution
The steady-state volume of distribution of aripiprazole following intravenous administration is high (404 L or 4.9 L/kg), indicating extensive extravascular distribution. All therapeutic concentrations of aripiprazole and its major metabolite are greater than 99% bound to serum proteins...

10.9 Elimination
The mean elimination half-lives are about 75 hours and 94 hours for aripiprazole and dehydro-aripiprazole, respectively. Steady-state concentrations are attained within 14 days of dosing for both active moieties...

10.10 Metabolism and Elimination
Aripiprazole is metabolized primarily by three biotransformation pathways: dehydrogenation, hydroxylation, and N-dealkylation. Based on in vitro studies, CYP3A4 and CYP2D6 enzymes are the major metabolic pathways for the dehydrogenation and hydroxylation of aripiprazole...

10.11 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.12 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.13 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.14 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.15 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.16 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.17 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.18 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.19 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.20 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.21 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.22 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.23 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...

10.24 Clinical Studies
The effects of aripiprazole on the exposures of aripiprazole and dehydro-aripiprazole are summarized in Figure 1 and Figure 2, respectively. Based on simulation, a 1.5-fold increase in mean Cmax and AUC values at steady-state is expected when extensive metabolizers of CYP2D6 are administered with both strong CYP2D6 and CYP3A4 inhibitors...




Figure 2: The effects of aripiprazole on the exposures of other drugs are summarized in Figure 2. The effects of aripiprazole on the exposures of other drugs are summarized in Figure 2.




Figure 3: The effects of aripiprazole on the exposures of other drugs are summarized in Figure 3. The effects of aripiprazole on the exposures of other drugs are summarized in Figure 3.




Figure 4: The effects of aripiprazole on the exposures of other drugs are summarized in Figure 4. The effects of aripiprazole on the exposures of other drugs are summarized in Figure 4.




Figure 5: The effects of aripiprazole on the exposures of other drugs are summarized in Figure 5. The effects of aripiprazole on the exposures of other drugs are summarized in Figure 5.




Figure 6: The effects of aripiprazole on the exposures of other drugs are summarized in Figure 6. The effects of aripiprazole on the exposures of other drugs are summarized in Figure 6.




Figure 7: The effects of aripiprazole on the exposures of other drugs are summarized in Figure 7. The effects of aripiprazole on the exposures of other drugs are summarized in Figure 7.




Figure 8: The effects of aripiprazole on the exposures of other drugs are summarized in Figure 8. The effects of aripiprazole on the exposures of other drugs are summarized in Figure 8.




Figure 9: The effects of aripiprazole on the exposures of other drugs are summarized in Figure 9. The effects of aripiprazole on the exposures of other drugs are summarized in Figure 9.




Figure 10: The effects of aripiprazole on the exposures of other drugs are summarized in Figure 10. The effects of aripiprazole on the exposures of other drugs are summarized in Figure 10.

In the four positive trials (n=414) comparing two fixed doses of aripiprazole (15 or 30 mg/day) to placebo, both doses of aripiprazole were superior to placebo in the PANSS total score (Study 1 in Table 16). PANSS positive subscale, and CGI-severity score. In addition, the 15 mg dose was superior to placebo in the PANSS negative subscale.

In a 4-week trial (n=404) comparing two fixed doses of aripiprazole (20 or 30 mg/day) to placebo, both doses of aripiprazole were superior to placebo in the PANSS total score (Study 2 in Table 16). PANSS positive subscale, PANSS negative subscale, and CGI-severity score.

In a 6-week trial (n=420) comparing three fixed doses of aripiprazole (10, 15, or 20 mg/day) to placebo, all three doses of aripiprazole were superior to placebo in the PANSS total score (Study 3 in Table 16). PANSS positive subscale, PANSS negative subscale, and CGI-severity score.

In a 6-week trial (n=367) comparing three fixed doses of aripiprazole (2, 5 or 10 mg/day) to placebo, the 10 mg dose of aripiprazole was superior to placebo in the PANSS total score (Study 4 in Table 16). The primary outcome measure of the study. The 2 and 5 mg doses did not demonstrate superiority to placebo on the primary outcome measure.

An examination of population subgroups did not reveal any clear evidence of differential responsiveness on the basis of age, gender, or race. A longer-term trial enrolled 310 inpatients or outpatients meeting DSM-IV criteria for schizophrenia who were, by history, symptomatically stable on either antipsychotic medications for periods of 3 months or longer. These patients were randomized to aripiprazole 15 mg/day or placebo for up to 26 weeks of observation for relapse.

The efficacy of aripiprazole in the treatment of schizophrenia in pediatric patients (13 to 17 years of age) was evaluated in one 6-week, placebo-controlled trial of outpatients who met DSM-IV criteria for schizophrenia and had a PANSS total score >10 at baseline. In this trial (n=302) comparing two fixed doses of aripiprazole (10 or 30 mg/day) to placebo, aripiprazole was superior to placebo in the PANSS total score at 5 and 11 days in the 10 mg/day treatment arm and in 11 days in the 30 mg/day treatment arm.

The efficacy of aripiprazole in the treatment of schizophrenia in pediatric patients (13 to 17 years of age) was evaluated in one 6-week, placebo-controlled trial of outpatients who met DSM-IV criteria for schizophrenia and had a PANSS total score >10 at baseline. In this trial (n=302) comparing two fixed doses of aripiprazole (10 or 30 mg/day) to placebo, aripiprazole was superior to placebo in the PANSS total score at 5 and 11 days in the 10 mg/day treatment arm and in 11 days in the 30 mg/day treatment arm.

Table 16: Schizophrenia Studies. Columns: Study Number, Treatment Group, Primary Efficacy Measure: PANSS (Mean Baseline Score (SD), LS Mean Change from Baseline (SE), Placebo-subtracted Difference* (95% CI)).

SD: standard deviation; SE: standard error; LS: Least-squares mean; CI: unadjusted confidence interval. *Difference (drug minus placebo) in least-squares mean change from baseline.

Figure 7: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 7). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 8: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 8). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 9: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 9). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 10: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 10). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 11: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 11). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 12: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 12). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 13: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 13). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 14: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 14). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 15: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 15). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 16: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 16). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 17: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 17). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 18: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 18). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 19: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 19). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 20: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 20). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 21: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 21). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Table 17: Bipolar Studies. Columns: Study Number, Treatment Group, Primary Efficacy Measure: Y-MRS (Mean Baseline Score (SD), LS Mean Change from Baseline (SE), Placebo-subtracted Difference* (95% CI)).

SD: standard deviation; SE: standard error; LS: Least-squares mean; CI: unadjusted confidence interval. *Difference (drug minus placebo) in least-squares mean change from baseline.

Figure 7: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 7). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 8: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 8). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 9: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 9). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 10: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 10). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 11: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 11). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 12: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 12). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 13: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 13). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 14: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 14). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 15: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 15). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 16: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 16). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 17: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 17). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 18: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 18). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 19: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 19). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 20: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 20). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 21: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 21). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 22: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 22). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 23: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 23). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 24: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 24). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 25: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 25). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 26: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 26). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 27: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 27). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

Figure 28: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Relapse (Bipolar Study 28). The plot shows the cumulative proportion of patients with relapse over time for aripiprazole and placebo groups.

16.2 Storage
Tablets
Store at 20° to 25° C (68° to 77° F); excursions permitted to 15°-30° C (59-86° F) [See USP Controlled Room Temperature].

17. PATIENT COUNSELING INFORMATION
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.1 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.2 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.3 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.4 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.5 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.6 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.7 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.8 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.9 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.10 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.11 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.12 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.13 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.14 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.15 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.16 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.17 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.18 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.19 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.20 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.21 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.22 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.23 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.24 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.25 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.26 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.27 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.28 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.29 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.30 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.31 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.32 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.33 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.34 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.35 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.36 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.37 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.38 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.39 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.40 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

17.41 Patient Counseling Information
See Medication Guide.
Discuss the following issues with patients prescribed aripiprazole:

Perforation

15 mm

297 mm

Medication Guide

Aripiprazole (AR-i-PIP-ra-zole) Tablets

Read this Medication Guide before you start taking aripiprazole and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or treatment.

What is the most important information I should know about aripiprazole?
(For other side effects, also see “What are the possible side effects of aripiprazole?”).

Serious side effects may happen when you take aripiprazole, including:

- **Increased risk of death in elderly patients with dementia-related psychosis:** Medicines like aripiprazole can raise the risk of death in elderly people who have lost touch with reality (psychosis) due to confusion and memory loss (dementia). Aripiprazole is not approved for the treatment of patients with dementia-related psychosis.
- **Risk of suicidal thoughts or actions:** Antidepressant medicines, depression and other serious mental illnesses, and suicidal thoughts or actions:
 1. **Antidepressant medicines may increase suicidal thoughts or actions in some children, teenagers, and young adults within the first few months of treatment.**
 2. **Depression and other serious mental illnesses are the most important causes of suicidal thoughts and actions. Some people may have a particularly high risk of having suicidal thoughts or actions.** These include people who have (or have a family history of) bipolar illness (also called manic-depressive illness) or suicidal thoughts or actions.
 3. **How can I watch for and try to prevent suicidal thoughts and actions in myself or a family member?**
 - Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is very important when an antidepressant medicine is started or when the dose is changed.
 - Call the healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings.
 - Keep all follow-up visits with the healthcare provider as scheduled. Call the healthcare provider between visits as needed, especially if you have concerns about symptoms.

Call a healthcare provider right away if you or your family member has any of the following symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- attempts to commit suicide
- new or worse depression
- new or worse anxiety
- feeling very agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

What else do I need to know about antidepressant medicines?

- **Never stop an antidepressant medicine without first talking to a healthcare provider.** Stopping an antidepressant medicine suddenly can cause other symptoms.

- **Antidepressants are medicines used to treat depression and other illnesses.** It is important to discuss all the risks of treating depression and also the risks of not treating it. Patients and their families or other caregivers should discuss all treatment choices with the healthcare provider, not just the use of antidepressants.
- **Antidepressant medicines have other side effects.** Talk to the healthcare provider about the side effects of the medicine prescribed for you or your family member.
- **Antidepressant medicines can interact with other medicines.** Know all of the medicines that you or your family member takes. Keep a list of all medicines to show the healthcare provider. Do not start new medicines without first checking with your healthcare provider.
- **Not all antidepressant medicines prescribed for children are FDA approved for use in children.** Talk to your child's healthcare provider for more information.

What is aripiprazole?

Aripiprazole is a prescription medicine used to treat:

- schizophrenia
- manic or mixed episodes that happen with bipolar I disorder

It is not known if aripiprazole is safe or effective in children:

- under 13 years of age with schizophrenia
- under 10 years of age with bipolar I disorder

Who should not take aripiprazole?

Do not take aripiprazole if you are allergic to aripiprazole or any of the ingredients in aripiprazole. See the end of this Medication Guide for a [complete list of ingredients in aripiprazole](#).

What should I tell my healthcare provider before taking aripiprazole?

Before taking aripiprazole, tell your healthcare provider if you have or had:

- diabetes or high blood sugar in you or your family; your healthcare provider should check your blood sugar before you start aripiprazole and also during therapy.
- seizures (convulsions).
- low or high blood pressure.
- heart problems or stroke.
- pregnancy or plans to become pregnant. It is not known if aripiprazole will harm your unborn baby.
- breast-feeding or plans to breast-feed. Aripiprazole can pass into your breast milk and may harm your baby. Talk to your healthcare provider about the best way to feed your baby if you receive aripiprazole.
- low white blood cell count.
- any other medical conditions.

Tell your healthcare provider about all the medicines that you take, including prescription medicines and over-the-counter medicines vitamins and herbal supplements.

Aripiprazole and other medicines may affect each other causing possible serious side effects. Aripiprazole may affect the way other medicines work, and other medicines may affect how aripiprazole works.

Your healthcare provider can tell you if it is safe to take aripiprazole with your other medicines. Do not start or stop any medicines while taking aripiprazole without talking to your healthcare provider first. Know the medicines you take. Keep a list of your medicines to show your healthcare provider and pharmacist when you get a new medicine.

How should I take aripiprazole?

- Take aripiprazole exactly as your healthcare provider tells you to take it. Do not change the dose or stop taking aripiprazole yourself.

210 mm

Perforation

15 mm

- Aripiprazole can be taken with or without food.
- Aripiprazole tablets should be swallowed whole.
- If you miss a dose of aripiprazole, take the missed dose as soon as you remember. If it is almost time for the next dose, just skip the missed dose and take your next dose at the regular time. Do not take two doses of aripiprazole at the same time.
- If you take too much aripiprazole, call your healthcare provider or poison control center at 1-800-222-1222 right away, or go to the nearest hospital emergency room.

What should I avoid while taking aripiprazole?

- Do not drive, operate heavy machinery, or do other dangerous activities until you know how aripiprazole affects you. Aripiprazole may make you drowsy.
- Avoid getting over-heated or dehydrated.
 - Do not over-exercise.
 - In hot weather, stay inside in a cool place if possible.
 - Stay out of the sun. Do not wear too much or heavy clothing.
 - Drink plenty of water.

What are the possible side effects of aripiprazole?

Aripiprazole may cause serious side effects, including:

See “What is the most important information I should know about aripiprazole?”

- **Stroke in elderly people (cerebrovascular problems) that can lead to death**
- **Neuroleptic malignant syndrome (NMS).** Tell your healthcare provider right away if you have some or all of the following symptoms: high fever, stiff muscles, confusion, sweating, changes in pulse, heart rate, and blood pressure. These may be symptoms of a rare and serious condition that can lead to death. Call your healthcare provider right away if you have any of these symptoms.
- **Uncontrolled body movements (tardive dyskinesia).** Aripiprazole may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop receiving aripiprazole. Tardive dyskinesia may also start after you stop receiving aripiprazole.
- **Problems with your metabolism such as:**
- **High blood sugar (hyperglycemia) and diabetes.** Increases in blood sugar can happen in some people who take aripiprazole. Extremely high blood sugar can lead to coma or death. If you have diabetes or risk factors for diabetes (such as being overweight or a family history of diabetes), your healthcare provider should check your blood sugar before you start aripiprazole and during your treatment.

Call your healthcare provider if you have any of these symptoms of high blood sugar while taking aripiprazole:

- feel very thirsty
- need to urinate more than usual
- feel very hungry
- feel weak or tired
- feel sick to your stomach
- feel confused, or your breath smells fruity
- **increased fat levels (cholesterol and triglycerides) in your blood.**
- **weight gain.** You and your healthcare provider should check your weight regularly.

- **Orthostatic hypotension (decreased blood pressure).** Lightheadedness or fainting when rising too quickly from a sitting or lying position.

- **Low white blood cell count**
- **Seizures (convulsions)**
- **problems with control of your body temperature especially when you exercise a lot or are in an area that is very hot. It is important for you to drink water to avoid dehydration.** See “What should I avoid while receiving aripiprazole?”
- **difficulty swallowing that can cause food or liquid to get into your lungs.**

The most common side effects of aripiprazole in adults include:

- nausea
- vomiting
- constipation
- headache
- blurred vision
- dizziness
- anxiety
- insomnia
- restlessness
- inner sense of restlessness/need to move (akathisia)
- upper respiratory illness

The most common side effects of aripiprazole in children include:

- feeling sleepy
- headache
- vomiting
- fatigue
- increased or decreased appetite
- insomnia
- nausea
- stuffy nose
- weight gain
- uncontrolled movement such as restlessness, tremor, muscle stiffness

- increased saliva or drooling

These are not all the possible side effects of aripiprazole. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store aripiprazole?

- Store aripiprazole at 20° to 25°C (68° to 77°F); excursions permitted to 15-30°C (59-86°F) [See USP Controlled Room Temperature].

Keep aripiprazole and all medicines out of the reach of children.

General information about the safe and effective use of aripiprazole

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use aripiprazole for a condition for which it was not prescribed. Do not give aripiprazole to other people, even if they have the same condition. It may harm them.

This Medication Guide summarizes the most important information about aripiprazole. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about aripiprazole that was written for healthcare professionals.

For more information about aripiprazole call Trigen Laboratories, LLC at 888 9 TRIGEN (888-987-4436).

What are the ingredients in aripiprazole tablets?

Active ingredient: aripiprazole

Inactive ingredients: corn starch, hydroxypropyl cellulose, lactose monohydrate, crospovidone, colloidal silicon dioxide, magnesium stearate and microcrystalline cellulose. Colorants include ferric oxide (yellow or red).

Additional pediatric use information is approved for Otsuka America Pharmaceutical, Inc.’s ABILIFY® (aripiprazole) product. However, due to Otsuka America Pharmaceutical, Inc.’s marketing exclusivity rights, this drug product is not labeled with that information.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

ABILIFY® is a trademark of Otsuka Pharmaceutical Company.

Made in India.

Manufactured for:
Trigen Laboratories, LLC
Sayreville, NJ 08872
www.trigenlab.com

TRIGEN
LABORATORIES

20014082

Issued: 04/2015

210 mm

297 mm