

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use ORENITRAM™ safely and effectively. See Full Prescribing Information for ORENITRAM.

**ORENITRAM (treprostinil) Extended Release Tablets for oral administration**  
Initial U.S. Approval: 2002

**INDICATIONS AND USAGE**

Orenitram is a prostacyclin vasodilator indicated for:

- Treatment of pulmonary arterial hypertension (PAH) (WHO Group 1) to improve exercise capacity. The study that established effectiveness included predominately patients with WHO functional class II-III symptoms and etiologies of idiopathic or heritable PAH (75%) or PAH associated with connective tissue disease (19%). (1.1)

As the sole vasodilator, the effect on exercise is small. Orenitram has not been shown to add to other vasodilator therapy. (1.1)

**DOSAGE AND ADMINISTRATION**

- Give with food. Swallow tablets whole; use only intact tablets. (2.1)
- Starting dose: 0.25 mg BID. (2.1)
- Titrate by 0.25 mg or 0.5 mg BID or 0.125 mg TID, not more than every 3 to 4 days as tolerated. (2.1)
- Maximum dose is determined by tolerability. (2.1)
- Mild hepatic impairment (Child Pugh Class A): Initiate at 0.125 mg BID. Increment at 0.125 mg BID every 3 to 4 days. (2.1)
- Avoid use in patients with moderate hepatic impairment. (2.1)

**DOSAGE FORMS AND STRENGTHS**

Extended Release Tablets: 0.125 mg, 0.25 mg, 1 mg and 2.5 mg. (3)

**CONTRAINDICATIONS**

- Severe hepatic impairment (Child Pugh Class C). (4)

**WARNINGS AND PRECAUTIONS**

- Do not abruptly discontinue dosing. (2.2, 5.1)
- Increased risk of bleeding, particularly in patients receiving anticoagulants. (5.2)
- Do not take Orenitram with alcohol (5.3)
- In patients with diverticulosis Orenitram tablets can become lodged in a diverticulum. (5.4)

**ADVERSE REACTIONS**

Most common adverse reactions (incidence >10%) reported in clinical studies with Orenitram are headache, nausea, and diarrhea. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact United Therapeutics Corp. at 1-866-458-6479 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

**DRUG INTERACTIONS**

- Blood pressure lowering drugs (e.g., diuretics, antihypertensive agents, or vasodilators): Risk of hypotension (7.1)
- When co-administered with strong CYP2C8 inhibitors the initial dose is 0.125 mg BID with 0.125 mg BID dose increments every 3 to 4 days. (7.3)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 12/2013

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1. INDICATIONS AND USAGE**

1.1 Pulmonary Arterial Hypertension

**2. DOSAGE AND ADMINISTRATION**

2.1 Recommended Dosing

2.2 Interruptions and Discontinuation

**3. DOSAGE FORMS AND STRENGTHS**

**4. CONTRAINDICATIONS**

**5. WARNINGS AND PRECAUTIONS**

5.1 Worsening PAH Symptoms upon Abrupt Withdrawal

5.2 Risk of Bleeding

5.3 Increased Exposure with Alcohol

5.4 Use in Patients with Blind-end Pouches

**6. ADVERSE REACTIONS**

6.1 Clinical Trials Experience

**7. DRUG INTERACTIONS**

7.1 Antihypertensive Agents or Other Vasodilators

7.2 Anticoagulants

7.3 Effect of CYP2C8 Inhibitors

**8. USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Labor and Delivery

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Patients with Hepatic Impairment

8.7 Patients with Renal Impairment

**10. OVERDOSAGE**

**11. DESCRIPTION**

**12. CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**14. CLINICAL STUDIES**

14.1 Clinical Trials in Pulmonary Arterial Hypertension (PAH)

**16. HOW SUPPLIED / STORAGE AND HANDLING**

16.1 How Supplied

16.2 Storage

**17. PATIENT COUNSELING INFORMATION**

\*Sections or subsections omitted from the full prescribing information are not listed

## FULL PRESCRIBING INFORMATION

### 1. INDICATIONS AND USAGE

#### 1.1 Pulmonary Arterial Hypertension

Orenitram is indicated for the treatment of pulmonary arterial hypertension (PAH) (WHO Group 1) to improve exercise capacity. The study that established effectiveness included predominately patients with WHO functional class II-III symptoms and etiologies of idiopathic or heritable PAH (75%) or PAH associated with connective tissue disease (19%).

When used as the sole vasodilator, the effect of Orenitram on exercise is about 10% of the deficit, and the effect, if any, on a background of another vasodilator is probably less than this. Orenitram is probably most useful to replace subcutaneous, intravenous, or inhaled treprostinil, but this use has not been studied.

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Recommended Dosing

Orenitram is supplied as extended release tablets. Individualize dosing of Orenitram according to clinical response.

The recommended starting dose of Orenitram is 0.25 mg twice daily (BID) with food, taken approximately 12 hours apart. Increase the dose as tolerated to achieve optimal clinical response. The recommended increment is 0.25 or 0.5 mg BID every 3-4 days. If 0.25 mg BID dose increments are not tolerated consider titrating slower. The total daily dose can be divided and given three times daily with food (TID; approximately 8 hours apart), titrating by increments of 0.125 mg TID.

The maximum dose is determined by tolerability. The mean dose in a controlled clinical trial at 12 weeks was 3.4 mg BID. Maximum doses studied were 12 mg BID in the 12-week blinded study and up to 21 mg BID in an open-label long-term study.

If intolerable pharmacologic effects occur, decrease the dose in increments of 0.25 mg. Avoid abrupt discontinuation [see *Warnings and Precautions* (5.1)].

**Hepatic impairment:** In patients with mild hepatic impairment (Child Pugh Class A) start at 0.125 mg BID with 0.125 mg BID dose increments every 3 to 4 days. Avoid use of Orenitram in patients with moderate hepatic impairment (Child Pugh Class B). Orenitram is contraindicated in patients with severe hepatic impairment (Child Pugh Class C) [see *Contraindications* (4), *Warnings and Precautions* (5.4), *Use in Specific Populations* (8.6), and *Clinical Pharmacology* (12.3)].

**Concomitant administration with CYP2C8 inhibitors:** When co-administered with strong CYP2C8 inhibitors (e.g., gemfibrozil) the initial dose is 0.125 mg BID with 0.125 mg BID dose increments every 3 to 4 days.

Take Orenitram with food. Swallow Orenitram intact; use only intact tablets.

#### 2.2 Interruptions and Discontinuation

If a dose of medication is missed, take the missed dose as soon as possible, with food. If a patient misses two or more doses, restart at a lower dose and re-titrate.

In the event of a planned short-term treatment interruption for patients unable to take oral medications, consider a temporary infusion of subcutaneous or intravenous treprostinil. To calculate the total daily dose (mg) of treprostinil for the parenteral route divide the oral total daily dose by 5.

When discontinuing Orenitram, reduce the dose in steps of 0.5 to 1 mg per day [see *Warnings and Precautions (5.1)*].

### 3 DOSAGE FORMS AND STRENGTHS

Orenitram (treprostinil extended-release) is available in the following four strengths:

- 0.125 mg [White tablet imprinted with UT 0.125]
- 0.25 mg [Green tablet imprinted with UT 0.25]
- 1 mg [Yellow tablet imprinted with UT 1]
- 2.5 mg [Pink tablet imprinted with UT 2.5]

### 4 CONTRAINDICATIONS

Severe hepatic impairment (Child Pugh Class C) [see *Use In Specific Populations (8.6)* and *Clinical Pharmacology (12.3)*].

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Worsening PAH Symptoms upon Abrupt Withdrawal

Abrupt discontinuation or sudden large reductions in dosage of Orenitram may result in worsening of PAH symptoms.

#### 5.2 Risk of Bleeding

Orenitram inhibits platelet aggregation and increases the risk of bleeding.

#### 5.3 Increased Exposure with Alcohol

Do not take Orenitram with alcohol as release of treprostinil from the tablet may occur at a faster rate than intended.

#### 5.4 Use in Patients with Blind-end Pouches

The tablet shell does not dissolve. In patients with diverticulosis, Orenitram tablets can lodge in a diverticulum.

### 6 ADVERSE REACTIONS

#### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

In a 12-week placebo-controlled monotherapy study (Study 1; WHO Group 1; functional class II-III), the most commonly reported adverse reactions that occurred in patients receiving Orenitram included: headache, nausea, and diarrhea. Table 1 lists the adverse reactions that occurred at a rate on Orenitram at least 5% higher than on placebo.

Orenitram patients in Table 1 for Study 1 (N = 151) had access to 0.25 mg tablets at randomization. Approximately 91% of such patients experienced an adverse reaction, but only 4% discontinued therapy for an adverse reaction (compared to 3% receiving placebo). The overall discontinuation rate for any reason was 17% for active and 14% for placebo.

**Table 1: Adverse Reactions with Rates at Least 5% Higher on Orenitram Monotherapy than on Placebo**

Reaction	Orenitram N=151	Placebo N=77
Headache	63%	19%
Diarrhea	30%	16%
Nausea	30%	18%
Flushing	15%	6%
Pain in jaw	11%	4%
Pain in extremity	14%	8%
Hypokalemia	9%	3%
Abdominal discomfort	6%	0%

Orenitram was studied in a long-term, open-label extension study in which 824 patients were dosed for a mean duration of approximately 2 years. About 70% of patients continued treatment with Orenitram for at least a year. The mean dose was 4.2 mg BID at one year. The adverse reactions were similar to those observed in the placebo-controlled trials.

## 7 DRUG INTERACTIONS

### 7.1 Antihypertensive Agents or Other Vasodilators

Concomitant administration of Orenitram with diuretics, antihypertensive agents or other vasodilators increases the risk of symptomatic hypotension.

### 7.2 Anticoagulants

Treprostinil inhibits platelet aggregation; there is increased risk of bleeding, particularly among patients receiving anticoagulants.

### 7.3 Effect of CYP2C8 Inhibitors

Co-administration of Orenitram and the CYP2C8 enzyme inhibitor gemfibrozil in healthy adult volunteers increases exposure to treprostinil. Reduce the starting dose of Orenitram to 0.125 mg BID and use 0.125 mg BID increments every 3 to 4 days [see *Dosage and Administration (2.1)* and *Clinical Pharmacology (12.3)*].

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

Pregnancy Category C.

Animal reproductive studies with treprostinil diolamine have shown an adverse effect on the fetus. There are no adequate and well-controlled studies in humans.

In rats, treatment with treprostinil diolamine had no effect on reproductive performance or sperm motility at doses up to 10 mg/kg/day. The exposures at this dose level are about 10- (male) to 18- (female) fold the usual human exposure at the mean dose of 3.4 mg BID.

In pregnant rats, reversible, dose-dependent decreases in body weight gain and food consumption were observed during the first four days of dosing in animals administered 10, 20 and 30 mg/kg/day treprostinil diolamine. In a dose range-finding study, there was a 17% decrease in the pregnancy rate in the animals administered 20 and 30 mg/kg/day. One dam in each of the 20 and 30 mg/kg/day had litters with no viable fetuses. In the definitive study (0, 5, 10 and 20 mg/kg/day), there were four treatment-related deaths, and a 32% decrease in the pregnancy rate for rats administered 20 mg/kg/day. There was an 8% decrease in the pregnancy rate in the animals administered 10 mg/kg/day. Across both studies, an increase in post-implantation loss was observed in animals administered 10 to 30 mg/kg/day, and a significant decrease in the mean number of live births was seen at dose levels  $\geq 10$  mg/kg/day. The no observed adverse effect level was 5 mg/kg/day (maternal, fetal viability and growth), and 20 mg/kg/day (teratogenicity), the highest dose tested in the definitive study. The exposures at 5 and 20 mg/kg/day doses represent 13 and 55 times, respectively, the human exposure.

For F<sub>1</sub> progeny, a decreased copulation index was observed at the 5 and 10 mg/kg/day treprostinil diolamine dose levels in rats. The no observed effect levels for physical development, reflex development, exploratory behavior, learning and memory, and sexual maturation was 10 mg/kg/day. The no observed effect level for F<sub>1</sub> progeny general development (based on body weight) was 10 mg/kg/day for females and  $\leq 2.5$  mg/kg/day for males; the no observed effect level for F<sub>1</sub> reproductive performance was 2.5 mg/kg/day or 6 times the human exposure.

In pregnant rabbits, the primary maternal adverse effects were gastrointestinal disturbance; dose-dependent decreases in mean body weight, body weight gain, and food consumption were observed. During the post-dose phase, the effect was reversed. In a dose range-finding study, there was a 17% decrease in the pregnancy rate for animals administered 4 mg/kg/day. A dose-dependent increase in post-implantation loss was observed. Two dams administered 4 mg/kg/day had litters with no viable fetuses; the mean fetal weight was slightly decreased in animals administered 4 mg/kg/day. In the definitive study, mean fetal weights were significantly decreased in animals administered 0.5 to 3 mg/kg/day of treprostinil diolamine. At doses of 1.5 and 3 mg/kg/day, external fetal and soft tissue malformations were observed in a few fetuses, and the total fetal skeletal malformations were significantly increased. The no observed adverse effect level was less than 0.5 mg/kg/day (maternal), 1.5 mg/kg/day (fetal viability and growth), and 0.5 mg/kg/day (teratogenicity). The 0.5 mg/kg/day dose represents about 5 times the human exposure.

## **8.2 Labor and Delivery**

The effect of Orenitram on labor and delivery in humans is unknown. No treprostinil treatment-related effects on labor and delivery were seen in animal studies.

## **8.3 Nursing Mothers**

It is not known whether treprostinil is excreted in human milk or absorbed systemically after ingestion. Because many drugs are excreted in human milk, choose Orenitram or breastfeeding.

## **8.4 Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

## **8.5 Geriatric Use**

Clinical studies of Orenitram did not include sufficient numbers of patients aged 65 years and over to determine whether they respond differently from younger patients. In general, dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic or cardiac function, and of concomitant disease or other drug therapy.

### 8.6 Patients with Hepatic Impairment

There is a marked increase in the systemic exposure to treprostinil in hepatically impaired patients [see *Dosage and Administration* (2.1), *Contraindications* (4), and *Clinical Pharmacology* (12.3)].

### 8.7 Patients with Renal Impairment


No dose adjustments are required in patients with renal impairment. Orenitram is not removed by dialysis [see *Clinical Pharmacology* (12.3)].

## 10 OVERDOSAGE

Signs and symptoms of overdose with Orenitram during clinical trials reflect its dose-limiting pharmacologic effects and include severe headache, nausea, vomiting, diarrhea, and hypotension. Treat supportively.

## 11 DESCRIPTION

Orenitram is an extended release osmotic tablet for oral administration. Orenitram is formulated as the diolamine salt of treprostinil, a tricyclic benzindene analogue of prostacyclin. The chemical name is Acetic acid, 2-[[[(1R,2R,3aS,9aS)-2,3,3a,4,9,9a-hexahydro-2-hydroxy-1-[(3S)-3-hydroxyoctyl]-1H-benz[f]inden-5-yl]oxy]-, complexed with 2,2'-iminobis[ethanol] (1:1). The molecular formula is  $C_{23}H_{34}O_5 \cdot C_4H_{11}NO_2$ , the molecular weight is 495.65, and it has the following structural formula:



Orenitram tablets are formulated in four strengths, which contain 0.125 mg of treprostinil (equivalent to 0.159 mg treprostinil diolamine), 0.25 mg of treprostinil (equivalent to 0.317 mg treprostinil diolamine), 1 mg of treprostinil (equivalent to 1.27 mg treprostinil diolamine), or 2.5 mg of treprostinil (equivalent to 3.17 mg treprostinil diolamine). The formulations also contain xylitol, maltodextrin, sodium lauryl sulfate, magnesium stearate, cellulose acetate, triethyl citrate, polyvinyl alcohol, titanium dioxide, polyethylene glycol, and talc. In addition tablets may contain colorants FD&C Blue #2, iron oxide yellow, and iron oxide red. The imprinting ink contains shellac glaze, ethanol, isopropyl alcohol USP, iron oxide black, n-butyl alcohol, propylene glycol, and ammonium hydroxide.

Orenitram is designed to release treprostinil at a near zero-order rate using an osmotic tablet technology. The tablet core is coated with a semi-permeable membrane and has a laser-drilled aperture through the membrane. Upon contact with water (e.g., after ingestion), the core tablet absorbs water through the semi-permeable membrane. The water dissolves the water-soluble treprostinil diolamine and the water-soluble osmotic excipients, which creates hydrostatic pressure within the membrane, eventually forcing the drug out through the tablet at a controlled rate.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

The major pharmacologic actions of treprostinil are direct vasodilation of pulmonary and systemic arterial vascular beds, inhibition of platelet aggregation, and inhibition of smooth muscle cell proliferation.

### 12.2 Pharmacodynamics

In a clinical trial of 240 healthy adult volunteers, single doses of inhaled treprostinil 54 µg (the target clinical dose) and 84 µg (supratherapeutic inhalation dose) prolonged the corrected QTc interval by approximately 10 msec. The QTc effect dissipated rapidly as the concentration of treprostinil decreased. Orenitram has not been evaluated in a thorough QTc study.

### 12.3 Pharmacokinetics

In patients with PAH, pharmacokinetics of treprostinil is dose-proportional for systemic exposure ( $AUC_{0-t}$ ) over the dose range of 0.5 and 15 mg BID. Upon repeat administration with a BID regimen, the accumulation in the systemic exposures to treprostinil is minimal and results in a peak-to-trough ratio of approximately 7. However, a TID regimen will reduce the peak-to-trough fluctuations to approximately 2.5 for the same total daily dose.

#### Absorption

The absolute oral bioavailability of Orenitram is approximately 17%. Maximum treprostinil concentrations occur between approximately 4 and 6 hours following Orenitram administration.

The absorption of Orenitram is affected by food. The  $AUC_{inf}$  of treprostinil was increased by 49% and the  $C_{max}$  was increased by an average of 13% when Orenitram was administered following a high-fat, high-calorie meal compared to fasting conditions in healthy volunteers. The relative bioavailability of treprostinil following oral administration of Orenitram 1 mg is not significantly altered by meal types ranging from 250 to 500 calories in healthy volunteers.

#### Distribution

The treprostinil component of Orenitram is highly bound to human plasma proteins, approximately 96% over a treprostinil concentration range of 0.01-10 µg/mL.

#### Metabolism and Excretion

In a study conducted in healthy volunteers using [ $^{14}C$ ] treprostinil, treprostinil was extensively metabolized on the side chain of the molecule via oxidation, oxidative cleavage, dehydration, and glucuronic acid conjugation. Treprostinil is primarily metabolized by CYP2C8 and to a lesser extent by CYP2C9. No new metabolites are found upon oral administration compared to parenteral administration of treprostinil. Only 1.13% and 0.19% is excreted as unchanged parent drug in the feces and urine, respectively. Based on *in vitro* studies treprostinil does not inhibit or induce major CYP enzymes [see *Drug Interactions* (7.3)].

#### Special Populations


**Hepatic Impairment:** In subjects with mild (n=8) hepatic impairment, administration of a single 1 mg dose of Orenitram resulted in a mean  $C_{max}$  and an  $AUC_{0-inf}$  that were 1.6- and 2.1-fold, respectively values seen in healthy subjects. With moderate impairment (n=8), the corresponding ratios were 4.0- and 4.8-fold, and with severe impairment (n=6), they were 4.8- and 7.6-fold [see *Dosage and Administration* (2.1), *Contraindications* (4), and *Use in Specific Populations* (8.6)].

**Renal Impairment:** In patients with severe renal impairment requiring dialysis (n=8), administration of a single 1 mg dose of Orenitram pre- and post-dialysis resulted in an  $AUC_{0-inf}$  that was not significantly altered compared to healthy subjects.

## Drug Interactions

Results of drug interaction studies are shown in Figure 1. Only for the strong CYP2C8 inhibitor does the interaction affect dosing.

**Figure 1: Impact of Co-Administered Drugs on the Systemic Exposure of Treprostinil 1 mg Compared to Orenitram Administered Alone**



**Warfarin:** A drug interaction study was carried out with Remodulin co-administered with warfarin (25 mg/day) in healthy volunteers. There was no clinically significant effect of either medication on the pharmacokinetics of treprostinil. Additionally, treprostinil did not affect the pharmacokinetics or pharmacodynamics of warfarin. The pharmacokinetics of R- and S- warfarin and the international normalized ratio (INR) in healthy subjects given a single 25 mg dose of warfarin were unaffected by continuous subcutaneous infusion of treprostinil at an infusion rate of 10 ng/kg/min.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Oral administration of treprostinil diolamine to Tg.rasH2 mice at 0, 5, 10 and 20 mg/kg/day in males and 0, 3, 7.5 and 15 mg/kg/day in females daily for 26 weeks did not significantly increase the incidence of tumors. The exposures obtained at the highest dose levels used in males and females are about 8- and 17-fold, respectively, the human exposure at the mean dose of 3.4 mg BID.

*In vitro* genotoxicity studies with high doses of treprostinil did not demonstrate any mutagenic or clastogenic effects. Treprostinil diolamine was tested *in vivo* in a rat micronucleus assay and did not induce an increased incidence of micronucleated polychromatic erythrocytes.

No adverse effect doses for fertility, fetal viability / growth, fetal development (teratogenicity), and postnatal development were determined in rats. In pregnant rabbits, external fetal and soft tissue malformations and fetal skeletal malformation occurred with the no observed adverse effect level for these adverse effects of 0.5 mg/kg/day (5 times the human exposure) [see *Use in Specific Populations (8.1)*].

## 14 CLINICAL STUDIES

### 14.1 Clinical Trials in Pulmonary Arterial Hypertension (PAH)


Three multi-center, randomized, double-blind studies were conducted and compared oral Orenitram to placebo in a total of 349 (Study 1), 350 (Study 2), and 310 (Study 3) patients with PAH.

#### Study 1 (effect seen with no background vasodilator)


Study 1 was a 12-week, randomized (2:1 Orenitram to placebo), double-blind, placebo-controlled, international efficacy and safety study of Orenitram in patients with WHO Group 1 PAH not currently receiving PAH therapy. The primary efficacy endpoint was placebo-corrected change in six-minute walk distance (6MWD) from Baseline to Week 12. Study drug dose was titrated to a maximum of 12 mg BID based on clinical response and study drug tolerability. Study 1 enrolled 349 patients (overall analysis population) who were not receiving any PAH medication. At the beginning of the study, subjects were dosed with only the 1 mg tablets with 0.5 and 0.25 mg tablets introduced at sequentially later dates during the study. The primary analysis population consisted of the 228 patients who had access to the 0.25 mg tablet at the time of randomization. Patients were administered Orenitram or placebo twice daily, with the doses titrated to effect over the course of the 12-week trial. Patients were in WHO functional class II (~33%) and class III (~66%) with either idiopathic or heritable PAH (~75%), collagen vascular disease associated PAH (~19%), or PAH associated with HIV (1%) or congenital heart defect (5%) or other conditions (~6%). The patients' mean baseline 6MWD was approximately 330 meters. In the primary analysis population, 17% of patients discontinued Orenitram compared to 14% of patients on placebo.

The primary efficacy endpoint of the trial was the change in 6MWD at 12 weeks for the primary analysis population. Analysis of Study 1 results demonstrated that those patients receiving Orenitram compared to patients receiving placebo improved their median 6MWD by approximately +23 meters (Hodges-Lehmann estimate;  $p=0.013$ , non-parametric analysis of covariance in accordance with the pre-specified statistical analysis plan) as compared to patients receiving placebo as demonstrated in (Figure 2). The within group median change from baseline was +25 meters for Orenitram and -5 meters for placebo at week 12 (N=228). Mean dose ( $\pm$ SD) in the Orenitram group was  $2.3 \pm 1.3$ ,  $3.2 \pm 1.9$ , and  $3.4 \pm 1.9$  mg BID at Weeks 4, 8, and 12, respectively. The distribution of the 6MWD change from baseline at Week 12 was also plotted across the range of observed values (Figure 3).

Figure 2: Hodges-Lehmann Estimate of Treatment Effect by Visit for the Primary Analysis Population (Study 1)




**Figure 3: Plot of the Distribution of Peak 6MWD Changes at Week 12 for the Primary Analysis Population (Study 1)**



The placebo-corrected median treatment effect on 6MWD was estimated (using the Hodges Lehmann estimator) within various subpopulations defined by age, gender, disease etiology, and baseline 6MWD (Figure 4).

**Figure 4: Placebo Corrected Median Treatment Effect (Hodges-Lehmann estimate with 95% CI) on 6MWD Change from Baseline at Week 12 for Various Subgroups in the Primary Analysis Population (Study 1)**



Studies 2 and 3 (no effect on a background of ERA, PDE5 inhibitor, or both)

Studies 2 (N=350) and 3 (N=310) were 16-week, randomized, double-blind, placebo-controlled, international efficacy and safety studies of Orenitram in patients with WHO Group 1 PAH. The primary efficacy endpoint was placebo-corrected change in 6MWD from Baseline to Week 16. Patients were in WHO functional class II (~23%) and class III (~77%) with either idiopathic or heritable PAH (~66%), collagen vascular disease associated PAH (~29%), or PAH associated with HIV (1%) or congenital heart defect (4%). The patients' mean baseline 6MWD was approximately 340 meters. Approximately 40% were receiving both an ERA and a PDE5 inhibitor. The results did not demonstrate a benefit in exercise testing with median 6MWD at Week 16 (11 meters [Hodges-Lehmann estimate; p=0.072] and 10 meters [Hodges-Lehmann estimate; p=0.089], respectively).

Long-Term Treatment of Pulmonary Hypertension

Patients (N=824) from the placebo-controlled studies entered a long-term, uncontrolled, open-label extension study. The average exposure to Orenitram was approximately 2 years with a maximum exposure of approximately 6 years. The dose of Orenitram continued to increase over time with doses (mean ± SD) of 3.6 ± 2.7, 4.1 ± 3.1, and 5 ± 3.7 mg BID at 6 (n=649), 12 (n=433), and 24 months (n=238), respectively. Reasons for discontinuation from the study included adverse event (16%), progression of disease (15%), death (13%), and withdrawn consent (7%). In the 522 subjects that completed the 12-month efficacy assessment, their mean 6MWD improved by 24 meters compared to baseline (30 meters in monotherapy patients and 20 meters

when Orenitram was used in combination with an ERA and/or a PDE-5 inhibitor). Of the patients that remained in the study, overall survival was 92%, 87%, and 82% at the end of 1, 2, and 3-years, respectively, with progression-free survival (progression defined as death, discontinuation or addition of a PAH therapy) of 74%, 61%, and 47%. Without a control group, these data must be interpreted cautiously.

## 16 HOW SUPPLIED / STORAGE AND HANDLING

### 16.1 How Supplied

Orenitram is a 7 mm round biconvex tablet with strength identifying color and printing and supplied as follows:

Strength	Color	Printing on Tablets	NDC # Bottle/100
0.125 mg	White	UT 0.125	66302-300-01
0.25 mg	Green	UT 0.25	66302-302-01
1 mg	Yellow	UT 1	66302-310-01
2.5 mg	Pink	UT 2.5	66302-325-01

### 16.2 Storage

Store at 25°C (77°F); excursions 15°C to 30°C (59°F to 86°F) [See USP controlled room temperature]. Keep out of reach of children.

## 17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (*Patient Package Insert*).

Tell patients:

- Abrupt discontinuation of therapy could result in worsening of PAH symptoms.
- Take Orenitram with food.
- Swallow Orenitram tablets whole. Do not split, chew, crush, or break. Do not take a tablet that is damaged or broken.
- The biologically inert components of the tablet remain intact during gastrointestinal transit and are eliminated in the feces as an insoluble shell.
- Do not take Orenitram with alcohol.

©Copyright 2013 United Therapeutics Corp. All rights reserved.

Orenitram manufactured for:

United Therapeutics Corp.  
Research Triangle Park, NC 27709

**Patient Information**  
**Orenitram (oh-REN-i-tram)**  
**(treprostinil)**  
**Extended Release Tablets**

**What is Orenitram?**

Orenitram is a prescription medicine used to treat pulmonary arterial hypertension (PAH) which is high blood pressure in the arteries of your lungs. Orenitram may improve your ability to exercise.

It is not known if Orenitram is safe and effective in children under 18 years of age.

**Who should not take Orenitram?**

Do not take Orenitram if you have severe liver problems.

**What should I tell my healthcare provider before taking Orenitram?**

**Before you take Orenitram, tell your healthcare provider if you:**

- have liver problems
- have low blood pressure or high blood pressure
- have had a stroke
- have stomach ulcers
- have diverticulosis
- have any other medical conditions
- are pregnant or plan to become pregnant. It is not known if Orenitram will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if Orenitram passes into your breast milk. You and your healthcare provider should decide if you will take Orenitram or breastfeed. You should not do both.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Orenitram and other medicines may affect each other causing side effects. Do not start any new medicine until you check with your healthcare provider.

Especially tell your healthcare provider if you take another medicine that contains treprostinil, such as Remodulin or Tyvaso.

Know the medicines you take. Keep a list of them to show to your healthcare provider and pharmacist when you get a new medicine.

**How should I take Orenitram?**

- Take Orenitram exactly as your healthcare provider tells you to take it.
- Your healthcare provider will slowly increase your dose to find the dose of Orenitram that is right for you.
- **Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider. Stopping Orenitram suddenly may cause worsening of your PAH symptoms.**
- Orenitram is usually taken every 12 hours, but may be taken three times daily (approximately every 8 hours). If you have side effects, your healthcare provider may tell you to change your dose or when you take Orenitram.

- Take Orenitram with food.
- Swallow Orenitram tablets whole. Do not split, chew, crush, or break your Orenitram tablets. Do not take Orenitram tablets that are damaged or broken. **If Orenitram tablets are not taken whole, they may release too much medicine at one time. This can lead to side effects.**
- You may see the tablet shell in your stools (bowel movements). This is usually normal. The tablet shell is not digested. If you have diverticulosis, the tablet shell may get stuck in a blind pouch or diverticulum in your intestine.
- If you miss your dose of Orenitram, take the dose as soon as possible with food.
- If you miss two or more doses of Orenitram, call your healthcare provider to see if you need to change your dose.
- If you take too much Orenitram, call your healthcare provider or go to the nearest hospital emergency room right away.

### **What should I avoid while taking Orenitram?**

- Do not take Orenitram with alcohol.

### **What are the possible side effects of Orenitram?**

**Orenitram can cause serious side effects, including an increased risk of bleeding.**

The most common side effects of Orenitram include:

- headache
- nausea
- diarrhea
- 

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of Orenitram. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store Orenitram?**

- Store Orenitram at room temperature, between 68°F to 77°F (20°C to 25°C).

**Keep Orenitram and all medicines out of the reach of children.**

### **General information about the safe and effective use of Orenitram.**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet.

Use Orenitram only for the condition for which it was prescribed. Do not give Orenitram to other people, even if they have the same symptoms you have. It may harm them.

If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about Orenitram that is written for health professionals. For more information, go to [www.ORENITRAM.com](http://www.ORENITRAM.com) or call 1-877-864-8437.

### **What are the ingredients in Orenitram?**

**Active ingredient:** treprostinil


**Inactive ingredients:** xylitol, maltodextrin, sodium lauryl sulfate, magnesium stearate, cellulose acetate, triethyl citrate, polyvinyl alcohol, titanium dioxide, polyethylene glycol, and talc. In addition tablets may contain colorants FD&C Blue #2, iron oxide yellow, and iron oxide red. The imprinting ink contains shellac glaze, ethanol, isopropyl alcohol, iron oxide black, n-butyl alcohol, propylene glycol, and ammonium hydroxide.

This Patient Information has been approved by the U.S. Food and Drug Administration.


United Therapeutics Corp.  
Research Triangle Park, NC 27709 USA

Issued: December 2013


©Copyright 2013, United Therapeutics Corp. All rights reserved.  
ORENITRAM™ is a trademark of United Therapeutics Corp.




(b) (4)



(b) (4)



(b) (4)



(b) (4)

---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/

---

NORMAN L STOCKBRIDGE  
12/20/2013