


Drug Facts Label	Size
Drug Facts Title	11 pt
Drug Facts Continued	8 pt
Headings	8 pt
Subheadings	6 pt
Body Text	6 pt
Leading	6.5 pt
Characters Per Inch	<39
Bullets	5 pt
Space Before Bullet	2 ems
Barlines	1.5 pt
Hairlines	.33 pt



Drug Facts Label	Size
Drug Facts Title	11 pt
Drug Facts Continued	8 pt
Headings	8 pt
Subheadings	6 pt
Body Text	6 pt
Leading	6.5 pt
Characters Per Inch	<39
Bullets	5 pt
Space Before Bullet	2 ems
Barlines	1.5 pt
Hairlines	.33 pt



Drug Facts Label	Size
Drug Facts Title	11 pt
Drug Facts Continued	8 pt
Headings	8 pt
Subheadings	6 pt
Body Text	6 pt
Leading	6.5 pt
Characters Per Inch	<39
Bullets	5 pt
Space Before Bullet	2 ems
Barlines	1.5 pt
Hairlines	.33 pt



Consult your healthcare provider before using this product for your heart.

ASPIRIN THERAPY ™

VAZALORE

Aspirin Capsules, 81 mg Pain Reliever/Fever Reducer (NSAID)

30 Liquid-Filled Capsules

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL WITH REPEATED TEXT "SEALED FOR YOUR PROTECTION" UNDER CAP IS MISSING OR BROKEN.

NDC 73089-081-21 Consult your healthcare provider before using this product for your heart.

New!

ASPIRIN THERAPY ™

VAZALORE

Aspirin Capsules, 81 mg Pain Reliever/Fever Reducer (NSAID)

81 mg LOW DOSE

FIRST LIQUID-FILLED ASPIRIN CAPSULES™

30 Liquid-Filled Capsules

1 Bottle Inside



BOTTLE SHOWN ACTUAL SIZE

Drug Facts (continued)

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 4 to 8 capsules every 4 hours while symptoms persist
 - do not exceed 48 capsules in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at 15-30°C (59-86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, lecithin, oleic acid, pharmaceutical ink, potassium chloride, soybean oil, titanium dioxide

Questions or comments?
Call 1-888-842-3567 9:00 AM – 7:00 PM ET, Monday – Friday

PLX PHARMA INC. Distributed by: **PLx Pharma Inc.**
9 Fishers Lane, Sparta, NJ 07871
Visit us at: www.vazalore.com
Patents: www.vazalore.com/patents

RETAIN CARTON FOR COMPLETE WARNINGS AND INFORMATION

072221 XXXXXXX



3 73089 08121 6

FPO



Drug Facts Active ingredient (in each capsule) Purposes Aspirin (NSAID)* 81 mg.....Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug Uses ■ for the temporary relief of minor aches and pains associated with: ■ headache ■ backache ■ muscular aches ■ a cold ■ toothache ■ minor pain of arthritis ■ premenstrual and menstrual cramps ■ temporarily reduces fever ■ ask your doctor about other uses for aspirin capsules Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) This product contains soy. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed	Drug Facts (continued) Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer. Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes, gout or arthritis Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts for more than 3 days ■ ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ any new symptoms occur. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
---	---



Drug Facts Label	Size
Drug Facts Title	11 pt
Drug Facts Continued	8 pt
Headings	8 pt
Subheadings	6 pt
Body Text	6 pt
Leading	6.5 pt
Characters Per Inch	<39
Bullets	5 pt
Space Before Bullet	2 ems
Barlines	1.5 pt
Hairlines	.33 pt



94.01*

89.00*

49.20*

Drug Facts Label	Size
Drug Facts Title	11 pt
Drug Facts Continued	8 pt
Headings	8 pt
Subheadings	6 pt
Body Text	6 pt
Leading	6.5 pt
Characters Per Inch	<39
Bullets	5 pt
Space Before Bullet	2 ems
Barlines	1.5 pt
Hairlines	.33 pt



Drug Facts (continued)

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
 - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
 - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at 15-30°C (59-86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

Inactive ingredients anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, lecithin, oleic acid, pharmaceutical ink, potassium chloride, soybean oil, titanium dioxide

Questions or comments?

Call 1-888-842-3567 9:00 AM – 7:00 PM ET,
Monday – Friday



Distributed by: **PLx Pharma Inc.**
9 Fishers Lane, Sparta, NJ 07871
Visit us at: www.vazalore.com
Patents: www.vazalore.com/patents

**RETAIN CARTON FOR COMPLETE
WARNINGS AND INFORMATION**

Consult your healthcare provider before using this product for your heart.



Aspirin Capsules, 325 mg Pain Reliever/Fever Reducer (NSAID)

30 Liquid-Filled Capsules

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL WITH REPEATED TEXT "SEALED FOR YOUR
PROTECTION" UNDER CAP IS MISSING OR BROKEN.**

NDC 73089-325-21

Consult your healthcare provider before using this product for your heart.

New!

ASPIRIN THERAPY

VAZALORE

Aspirin Capsules, 325 mg Pain Reliever/Fever Reducer (NSAID)

325 mg

FIRST LIQUID-FILLED
ASPIRIN CAPSULES™

30 Liquid-Filled Capsules

072221 XXXXXXX



1 Bottle Inside



BOTTLE SHOWN ACTUAL SIZE



Drug Facts

Active ingredient (in each capsule) Purposes

Aspirin (NSAID)* 325 mg Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of minor aches and pains associated with:
 - headache ■ backache ■ muscular aches
 - a cold ■ toothache ■ minor pain of arthritis
 - premenstrual and menstrual cramps
- temporarily reduces fever
- ask your doctor about other uses for aspirin capsules

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling
- shock ■ asthma (wheezing)

This product contains soy.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.



This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
11/16/2021 03:03:27 PM