

#### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Ondansetron Injection USP safely and effectively. See full prescribing information for Ondansetron Injection USP. **ONDANSETRON Injection USP for intravenous use**  
**Initial U.S. Approval 1991**

#### INDICATIONS AND USAGE

Ondansetron Injection USP is a 5-HT<sub>3</sub> receptor antagonist indicated:

- Prevention of nausea and vomiting associated with initial and repeat courses of emetogenic cancer chemotherapy. (1.1)
- Prevention of postoperative nausea and/or vomiting. (1.2)

#### DOSE AND ADMINISTRATION

Prevention of nausea and vomiting associated with initial and repeat courses of emetogenic cancer chemotherapy (2.1):

- Adults and Pediatric patients (6 months to 18 years): Three 0.15 mg/kg doses, up to a maximum of 16 mg per dose, infused intravenously over 15 minutes. The first dose should be administered 30 minutes before the start of chemotherapy. Subsequent doses are administered 4 and 8 hours after the first dose.

Prevention of postoperative nausea and/or vomiting (2.2):

Population	Age	Ondansetron Injection USP Dosage	Intravenous Infusion Rate
Adults	>12 yrs	4 mg x 1	over 2 to 5 min
Pediatrics (> 40 kg)	1 mo. to 12 yrs	4 mg x 1	over 2 to 5 min
Pediatrics (≤40 kg)	1 mo. to 12 yrs	0.1 mg/kg x 1	over 2 to 5 min

- In patients with severe hepatic impairment, a total daily dose of 8 mg should not be exceeded. (2.4)

#### DOSE FORMS AND STRENGTHS

Ondansetron Injection USP (2 mg/mL), 2 mL single-use vials. (3)

#### CONTRAINDICATIONS

- Patients known to have hypersensitivity (e.g., anaphylaxis) to this product or any of its components. (4)
- Concomitant use of apomorphine. (4)

#### WARNINGS AND PRECAUTIONS

- Hypersensitivity reactions including anaphylaxis and bronchospasm, have been reported in patients who have exhibited hypersensitivity to other selective 5-HT<sub>3</sub> receptor antagonists. (5.1)
- QT prolongation occurs in a dose-dependent manner. Cases of Torsade de Pointes have been reported. Avoid Ondansetron Injection USP in patients with congenital long QT syndrome. (5.2)
- Use in patients following abdominal surgery or in patients with chemotherapy-induced nausea and vomiting may mask a progressive ileus and/or gastric distention. (5.3)(5.4)

#### ADVERSE REACTIONS

Chemotherapy-Induced Nausea and Vomiting –

- The most common adverse reactions (≥7%) in adults are diarrhea, headache, and fever. (6.1)
- Postoperative Nausea and Vomiting –
- The most common adverse reaction (≥10%) which occurs at a higher frequency compared to placebo in adults is headache. (6.1)
- The most common adverse reaction (≥2%) which occurs at a higher frequency compared to placebo in pediatric patients 1 to 24 months of age is diarrhea. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Qilu pharmaceutical Co. Ltd. at 86-531-8312-6000 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

#### DRUG INTERACTIONS

- Apomorphine – profound hypotension and loss of consciousness. Concomitant use with ondansetron is contraindicated. (7.2)

See 17 for PATIENT COUNSELING INFORMATION

Revised 2/2014

#### FULL PRESCRIBING INFORMATION CONTENTS \*

##### 1 INDICATIONS AND USAGE

- Prevention of Nausea and Vomiting Associated with Initial and Repeat Courses of Emetogenic Cancer Chemotherapy
- Prevention of Postoperative Nausea and/or Vomiting

##### 2 DOSAGE AND ADMINISTRATION

- Prevention of Nausea and Vomiting Associated with Initial and Repeat Courses of Emetogenic Chemotherapy
- Prevention of Postoperative Nausea and Vomiting
- Stability and Handling
- Dosage Adjustment for Patients with Impaired Hepatic Function

##### 3 DOSAGE FORMS AND STRENGTHS

##### 4 CONTRAINDICATIONS

##### 5 WARNINGS AND PRECAUTIONS

- Hypersensitivity Reactions
- QT Prolongation
- Masking of Progressive Ileus and Gastric Distention
- Effect on Peristalsis

##### 6 ADVERSE REACTIONS

- Clinical Trials Experience
- Postmarketing Experience

##### 7 DRUG INTERACTIONS

- Drugs Affecting Cytochrome P-450 Enzymes
- Apomorphine
- Phenytoin, Carbamazepine, and Rifampin
- Tramadol
- Chemotherapy
- Temazepam
- Alfentanil and Atracurium

##### 8 USE IN SPECIFIC POPULATIONS

- Pregnancy
- Nursing Mothers
- Pediatric Use
- Geriatric Use
- Hepatic Impairment
- Renal Impairment

##### 9 DRUG ABUSE AND DEPENDENCE

##### 10 OVERDOSAGE

##### 11 DESCRIPTION

##### 12 CLINICAL PHARMACOLOGY

- Mechanism of Action
- Pharmacodynamics
- Pharmacokinetics

##### 13 NONCLINICAL TOXICOLOGY

- Carcinogenesis, Mutagenesis, Impairment of Fertility
- CLINICAL STUDIES
- Chemotherapy-Induced Nausea and Vomiting
- Prevention of Postoperative Nausea and/or Vomiting
- Prevention of Further Postoperative Nausea and Vomiting

##### 16 HOW SUPPLIED/STORAGE AND HANDLING

##### 17 PATIENT COUNSELING INFORMATION

\* Sections or subsections omitted from the full prescribing information are not listed

#### FULL PRESCRIBING INFORMATION

##### 1 INDICATIONS AND USAGE

##### 1.1 Prevention of Nausea and Vomiting Associated with Initial and Repeat Courses of Emetogenic Cancer Chemotherapy

Ondansetron Injection USP is indicated for the prevention of nausea and vomiting associated with initial and repeat courses of emetogenic cancer chemotherapy, including high-dose cisplatin [see *Clinical Studies* (14.1)].

Ondansetron Injection USP is approved for patients aged 6 months and older.

##### 1.2 Prevention of Postoperative Nausea and/or Vomiting

Ondansetron Injection USP is indicated for the prevention of postoperative nausea and/or

vomiting. As with other antiemetics, routine prophylaxis is not recommended for patients in whom there is little expectation that nausea and/or vomiting will occur postoperatively. In patients in whom nausea and/or vomiting must be avoided postoperatively, Ondansetron Injection USP is recommended even when the incidence of postoperative nausea and/or vomiting is low. For patients who do not receive prophylactic Ondansetron Injection USP and experience nausea and/or vomiting postoperatively, Ondansetron Injection USP may be given to prevent further episodes [see *Clinical Studies* (14.3)].

Ondansetron Injection USP is approved for patients aged 1 month and older.

##### 2 DOSAGE AND ADMINISTRATION

##### 2.1 Prevention of Nausea and Vomiting Associated with Initial and Repeat Courses of Emetogenic Chemotherapy

Ondansetron Injection USP should be diluted in 50 mL of 5% Dextrose Injection or 0.9% Sodium Chloride Injection before administration.

##### Adults

The recommended adult intravenous dosage of Ondansetron Injection USP is three 0.15-mg/kg doses up to a maximum of 16 mg per dose [see *Clinical Pharmacology* (12.2)]. The first dose is infused over 15 minutes beginning 30 minutes before the start of emetogenic chemotherapy. Subsequent doses (0.15 mg/kg up to a maximum of 16 mg per dose) are administered 4 and 8 hours after the first dose of Ondansetron Injection USP.

##### Pediatrics

For pediatric patients 6 months through 18 years of age, the intravenous dosage of Ondansetron Injection USP is three 0.15-mg/kg doses up to a maximum of 16 mg per dose [see *Clinical Studies* (14.1) and *Clinical Pharmacology* (12.2 and 12.3)]. The first dose is to be administered 30 minutes before the start of moderately to highly emetogenic chemotherapy. Subsequent doses (0.15 mg/kg up to a maximum of 16 mg per dose) are administered 4 and 8 hours after the first dose of Ondansetron Injection USP. The drug should be infused intravenously over 15 minutes.

##### 2.2 Prevention of Postoperative Nausea and Vomiting

Ondansetron Injection USP should not be mixed with solutions for which physical and chemical compatibility has not been established. In particular, this applies to alkaline solutions as a precipitate may form.

##### Adults

The recommended adult intravenous dosage of Ondansetron Injection USP is 4 mg *undiluted* administered intravenously in not less than 30 seconds, preferably over 2 to 5 minutes, immediately before induction of anesthesia, or postoperatively if the patient did not receive prophylactic antiemetics and experiences nausea and/or vomiting occurring within 2 hours after surgery. Alternatively, 4 mg *undiluted* may be administered intramuscularly as a single injection for adults. While recommended as a fixed dose for patients weighing more than 40 kg, few patients above 80 kg have been studied. In patients who do not achieve adequate control of postoperative nausea and vomiting following a single, prophylactic, preinduction, intravenous dose of ondansetron 4 mg, administration of a second intravenous dose of 4 mg ondansetron postoperatively does not provide additional control of nausea and vomiting.

##### Pediatrics

For pediatric patients 1 month through 12 years of age, the dosage is a single 0.1-mg/kg dose for patients weighing 40 kg or less, or a single 4-mg dose for patients weighing more than 40 kg. The rate of administration should not be less than 30 seconds, preferably over 2 to 5 minutes immediately prior to or following anesthesia induction, or postoperatively if the patient did not receive prophylactic antiemetics and experiences nausea and/or vomiting occurring shortly after surgery. Prevention of further nausea and vomiting was only studied in patients who had not received prophylactic Ondansetron Injection USP.

##### 2.3 Stability and Handling

After dilution, do not use beyond 24 hours. Although Ondansetron Injection USP is chemically and physically stable when diluted as recommended, sterile precautions should be observed because diluents generally do not contain preservative.

Ondansetron Injection USP is stable at room temperature under normal lighting conditions for 48 hours after dilution with the following intravenous fluids: 0.9% Sodium Chloride Injection, 5% Dextrose Injection, 5% Dextrose and 0.9% Sodium Chloride Injection, 5% Dextrose and 0.45% Sodium Chloride Injection, and 3% Sodium Chloride Injection.

**Note:** Parenteral drug products should be inspected visually for particulate matter and discoloration before administration whenever solution and container permit.

**Precaution:** Occasionally, ondansetron precipitates at the stopper/vial interface in vials stored upright. Potency and safety are not affected. If a precipitate is observed, resubstitute by shaking the vial vigorously.

##### 2.4 Dosage Adjustment for Patients with Impaired Hepatic Function

In patients with severe hepatic impairment (Child-Pugh score of 10 or greater), a single maximal daily dose of 8 mg infused over 15 minutes beginning 30 minutes before the start of the emetogenic chemotherapy is recommended. There is no experience beyond first-day administration of ondansetron in these patients [see *Clinical Pharmacology* (12.3)].

##### 3 DOSAGE FORMS AND STRENGTHS

Ondansetron Injection USP, 2 mg/mL is a clear, colorless, nonpyrogenic, sterile solution available as 2 mL single-use vials.

##### 4 CONTRAINDICATIONS

Ondansetron Injection USP is contraindicated for patients known to have hypersensitivity (e.g., anaphylaxis) to this product or any of its components. Anaphylactic reactions have been reported in patients taking ondansetron. [See *Adverse Reactions* (6.2)].

The concomitant use of apomorphine with ondansetron is contraindicated based on reports of profound hypotension and loss of consciousness when apomorphine was administered with ondansetron.

##### 5 WARNINGS AND PRECAUTIONS

**5.1 Hypersensitivity Reactions**  
Hypersensitivity reactions, including anaphylaxis and bronchospasm, have been reported in patients who have exhibited hypersensitivity to other selective 5-HT<sub>3</sub> receptor antagonists.

##### 5.2 QT Prolongation

Ondansetron prolongs the QT interval in a dose-dependent manner [see *Clinical Pharmacology* (12.2)]. In addition, post-marketing cases of Torsade de Pointes have been reported in patients using ondansetron. Avoid Ondansetron Injection USP in patients with congenital long QT syndrome. ECG monitoring is recommended in patients with electrolyte abnormalities (e.g., hypokalemia or hypomagnesemia), congestive heart failure, bradyarrhythmias, or patients taking other medicinal products that lead to QT prolongation.

##### 5.3 Masking of Progressive Ileus and Gastric Distension

The use of Ondansetron Injection USP in patients following abdominal surgery or in patients with chemotherapy-induced nausea and vomiting may mask a progressive ileus and gastric distention.

##### 5.4 Effect on Peristalsis

Ondansetron Injection USP is not a drug that stimulates gastric or intestinal peristalsis. It should not be used instead of nasogastric suction.

##### 6 ADVERSE REACTIONS

##### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The following adverse reactions have been reported in clinical trials of adult patients treated with ondansetron, the active ingredient of intravenous Ondansetron Injection USP across a range of dosages. A causal relationship to therapy with Ondansetron Injection USP (ondansetron) was unclear in many cases.

##### Chemotherapy-Induced Nausea and Vomiting

Table 1. Adverse Reactions Reported in > 5% of Adult Patients Who Received Ondansetron at a Dosage of Three 0.15-mg/kg Doses

Adverse Reaction	Number of Adult Patients With Reaction		
	Ondansetron Injection USP 0.15 mg/kg x 3 n = 419	Metoclopramide n = 156	Placebo n = 34
Diarrhea	16%	44%	18%
Headache	17%	7%	15%
Fever	8%	5%	3%

##### Cardiovascular

Rare cases of angina (chest pain), electrocardiographic alterations, hypotension, and tachycardia have been reported.

##### Gastrointestinal

Constipation has been reported in 11% of chemotherapy patients receiving multiday ondansetron.

##### Hepatic

In comparative trials in cisplatin chemotherapy patients with normal baseline values of aspartate transaminase (AST) and alanine transaminase (ALT), these enzymes have been reported to exceed twice the upper limit of normal in approximately 5% of patients. The increases were transient and did not appear to be related to dose or duration of therapy. On repeat exposure, similar transient elevations in transaminase values occurred in some courses, but symptomatic hepatic disease did not occur.

##### Integumentary

Rash has occurred in approximately 1% of patients receiving ondansetron.

##### Neurological

There have been rare reports consistent with, but not diagnostic of, extrapyramidal reactions in patients receiving Ondansetron Injection USP, and rare cases of grand mal seizure.

##### Other

Rare cases of hypokalemia have been reported.

##### Postoperative Nausea and Vomiting

The adverse reactions in Table 2 have been reported in ≥ 2% of adults receiving ondansetron at a dosage of 4 mg intravenous over 2 to 5 minutes in clinical trials.

Table 2. Adverse Reactions Reported in ≥ 2% (and With Greater Frequency than the Placebo Group) of Adult Patients Receiving Ondansetron at a Dosage of 4 mg Intravenous over 2 to 5 Minutes

Adverse Reaction <sup>a, b</sup>	Ondansetron Injection USP 4 mg Intravenous n = 547 patients	Placebo n = 547 patients
Headache	92 (17%)	77 (14%)
Drowsiness/sedation	44 (8%)	37 (7%)
Injection site reaction	21 (4%)	18 (3%)
Fever	10 (2%)	6 (1%)
Cold sensation	9 (2%)	8 (1%)
Pruritus	9 (2%)	3 (< 1%)
Paresthesia	9 (2%)	2 (< 1%)

<sup>a</sup> Adverse Reactions: Rates of these reactions were not significantly different in the ondansetron and placebo groups

<sup>b</sup> Patients were receiving multiple concomitant perioperative and postoperative medications

##### Pediatric Use

Rates of adverse reactions were similar in both the ondansetron and placebo groups in pediatric patients receiving ondansetron (a single 0.1-mg/kg dose for pediatric patients weighing 40 kg or less, or 4 mg for pediatric patients weighing more than 40 kg) administered intravenously over at least 30 seconds. Diarrhea was seen more frequently in patients taking Ondansetron Injection USP (2%) compared to placebo (<1%) in the 1 month to 24 month age group. These patients were receiving multiple concomitant perioperative and postoperative medications.

##### 6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of ondansetron. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. The reactions have been chosen for inclusion due to a combination of their seriousness, frequency of reporting, or potential causal connection to ondansetron.

##### Cardiovascular

Arrhythmias (including ventricular and supraventricular tachycardia, premature ventricular contractions, and atrial fibrillation), bradycardia, electrocardiographic alterations (including second-degree heart block, QT/QTc interval prolongation, and ST segment depression), palpitations, and syncope. Rarely and predominantly with intravenous ondansetron, transient ECG changes including QT/QTc interval prolongation have been reported [see *Warnings and Precautions* (5.2)].

##### General

Flushing. Rare cases of hypersensitivity reactions, sometimes severe (e.g., aphalytic reactions, angioedema, bronchospasm, cardiopulmonary arrest, hypotension, laryngeal edema, laryngospasm, shock, shortness of breath, stridor) have also been reported. A positive lymphocyte transformation test to ondansetron has been reported, which suggests immunologic sensitivity to ondansetron.

##### Hepatobiliary

Liver enzyme abnormalities have been reported. Liver failure and death have been reported in patients with cancer receiving concurrent medications including potentially hepatotoxic cytotoxic chemotherapy and antibiotics.

##### Local Reactions

Pain, redness, and burning at site of injection.

##### Lower Respiratory

##### Hiccups

##### Neurological

Oculogyric crisis, appearing alone, as well as with other dystonic reactions. Transient dizziness during or shortly after intravenous infusion.

##### Skin

Urticaria, Stevens-Johnson syndrome, and toxic epidermal necrolysis.

##### Eye Disorders

Cases of transient blindness, predominantly during intravenous administration, have been reported. These cases of transient blindness were reported to resolve within a few minutes up to 48 hours. Transient blurred vision, in some cases associated with abnormalities of accommodation, has also been reported.

##### 7 DRUG INTERACTIONS

##### 7.1 Drugs Affecting Cytochrome P-450 Enzymes

Ondansetron does not appear to induce or inhibit the cytochrome P-450 drug-metabolizing enzyme system of the liver. Because ondansetron is metabolized by hepatic cytochrome P-450 drug-metabolizing enzymes (CYP3A4, CYP2D6, CYP1A2), inducers or inhibitors of these enzymes may change the clearance and, hence, the half-life of ondansetron [see *Clinical Pharmacology* (12.3)]. On the basis of limited available data, no dosage adjustment is recommended for patients on these drugs.

##### 7.2 Apomorphine

Based on reports of profound hypotension and loss of consciousness when apomorphine was administered with ondansetron, the concomitant use of apomorphine with ondansetron is contraindicated [see *Contraindications* (4)].

##### 7.3 Phenytoin, Carbamazepine, and Rifampin

In patients treated with potent inducers of CYP3A4 (i.e., phenytoin, carbamazepine, and rifampin), the clearance of ondansetron was significantly increased and ondansetron blood concentrations were decreased. However, on the basis of available data, no dosage adjustment for ondansetron is recommended for patients on these drugs [see *Clinical Pharmacology* (12.3)].

##### 7.4 Tramadol

Although there are no data on pharmacokinetic drug interactions between ondansetron and tramadol, data from two small studies indicate that concomitant use of ondansetron may result in reduced analgesic activity of tramadol. Patients on concomitant ondansetron self administered tramadol more frequently in these studies, leading to an increased cumulative dose in patient controlled administration (PCA) of tramadol.

##### 7.5 Chemotherapy

In humans, carmustine, etoposide, and cisplatin do not affect the pharmacokinetics of ondansetron.

In a crossover study in 76 pediatric patients, intravenous ondansetron did not increase blood levels of high-dose methotrexate.

##### 7.6 Temazepam

The coadministration of ondansetron had no effect on the pharmacokinetics and

pharmacodynamics of temazepam.

##### 7.7 Alfentanil and Atracurium

Ondansetron does not alter the respiratory depressant effects produced by alfentanil or the degree of neuromuscular blockade produced by atracurium. Interactions with general or local anesthetics have not been studied.

##### 8 USE IN SPECIFIC POPULATIONS

##### 8.1 Pregnancy

Pregnancy Category B. Reproduction studies have been performed in pregnant rats and rabbits at intravenous doses up to 4 mg/kg per day (approximately 1.4 and 2.9 times the recommended human intravenous dose of 0.15 mg/kg given three times a day, respectively, based on body surface area) and have revealed no evidence of impaired fertility or harm to the fetus due to ondansetron. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

##### 8.3 Nursing Mothers

Ondansetron is excreted in the breast milk of rats. It is not known whether ondansetron is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when ondansetron is administered to a nursing woman.

##### 8.4 Pediatric Use

Little information is available about the use of ondansetron in pediatric surgical patients younger than 1 month of age. [See *Clinical Studies* (14.2)]. Little information is available about the use of ondansetron in pediatric cancer patients younger than 6 months of age. [See *Clinical Studies* (14.1) and *Dosage and Administration* (2)].

The clearance of ondansetron in pediatric patients 1 month to 4 months of age is slower and the half-life is ~2.5 fold longer than patients who are > 4 to 24 months of age. As a precaution, it is recommended that patients less than 4 months of age receiving this drug be closely monitored. [See *Clinical Pharmacology* (12.3)].

##### 8.5 Geriatric Use

Of the total number of subjects enrolled in cancer chemotherapy-induced and postoperative nausea and vomiting in US- and foreign-controlled clinical trials, 862 were 65 years of age and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out. Dosage adjustment is not needed in patients over the age of 65 [see *Clinical Pharmacology* (12.3)].

##### 8.6 Hepatic Impairment

In patients with severe hepatic impairment (Child-Pugh score of 10 or greater), clearance is reduced and apparent volume of distribution is increased with a resultant increase in plasma half-life [see *Clinical Pharmacology* (12.3)]. In such patients, a total daily dose of 8 mg should not be exceeded [see *Dosage and Administration* (2.3)].

##### 8.7 Renal Impairment

Although plasma clearance is reduced in patients with severe renal impairment (creatinine clearance < 30 mL/min), no dosage adjustment is recommended [see *Clinical Pharmacology* (12.3)].

##### 9 DRUG ABUSE AND DEPENDENCE

Animal studies have shown that ondansetron is not discriminated as a benzodiazepine nor does it substitute for benzodiazepines in direct addiction studies.

##### 10 OVERDOSAGE

There is no specific antidote for ondansetron overdose. Patients should be managed with appropriate supportive therapy. Individual intravenous doses as large as 150 mg and total daily intravenous doses as large as 252 mg have been inadvertently administered without significant adverse events. These doses are more than 10 times the recommended daily dose.

In addition to the adverse reactions listed above, the following events have been described in the setting of ondansetron overdose:

injection. Systemic exposure as measured by mean AUC were equivalent, with values of 156 [95% CI 136, 180] and 161 [95% CI 137, 190] ng•h/mL for intravenous and intramuscular groups, respectively. Mean peak plasma concentrations were 42.9 [95% CI 33.8, 54.4] ng/mL at 10 minutes after intravenous infusion and 31.9 [95% CI 26.3, 38.6] ng/mL at 41 minutes after intramuscular injection.

**Distribution**  
 Plasma protein binding of ondansetron as measured in vitro was 70% to 76%, over the pharmacologic concentration range of 10 to 500 ng/mL. Circulating drug also distributes into erythrocytes.

**Metabolism**  
 Ondansetron is extensively metabolized in humans, with approximately 5% of a radiolabeled dose recovered as the parent compound from the urine. The primary metabolic pathway is hydroxylation on the indole ring followed by subsequent glucuronide or sulfate conjugation. Although some nonconjugated metabolites have pharmacologic activity, these are not found in plasma at concentrations likely to significantly contribute to the biological activity of ondansetron. The metabolites are observed in the urine.

In vitro metabolism studies have shown that ondansetron is a substrate for multiple human hepatic cytochrome P-450 enzymes, including CYP1A2, CYP2D6, and CYP3A4. In terms of overall ondansetron turnover, CYP3A4 plays a predominant role while formation of the major in vivo metabolites is apparently mediated by CYP1A2. The role of CYP2D6 in ondansetron in vivo metabolism is relatively minor.

The pharmacokinetics of intravenous ondansetron did not differ between subjects who were poor metabolizers of CYP2D6 and those who were extensive metabolizers of CYP2D6, further supporting the limited role of CYP2D6 in ondansetron disposition in vivo.

**Elimination**  
 In adult cancer patients, the mean ondansetron elimination half-life was 4 hours, and there was no difference in the multidose pharmacokinetics over a 4-day period. In a dose proportionality study, systemic exposure to 32 mg of ondansetron was not proportional to dose as measured by comparing dose-normalized AUC values to an 8-mg dose. This is consistent with a small decrease in systemic clearance with increasing plasma concentrations.

**Geriatrics**  
 A reduction in clearance and increase in elimination half-life are seen in patients over 75 years of age. In clinical trials with cancer patients, safety and efficacy were similar in patients over 65 years of age and those under 65 years of age; there was an insufficient number of patients over 75 years of age to permit conclusions in that age-group. No dosage adjustment is recommended in the elderly.

**Pediatrics**  
 Pharmacokinetic samples were collected from 74 cancer patients 6 to 48 months of age, who received a dose of 0.15 mg/kg of intravenous ondansetron every 4 hours for 3 doses during a safety and efficacy trial. These data were combined with sequential pharmacokinetics data from 41 surgery patients 1 month to 24 months of age, who received a single dose of 0.1 mg/kg of intravenous ondansetron prior to surgery with general anesthesia, and a population pharmacokinetic analysis was performed on the combined data set. The results of this analysis are included in Table 4 and are compared to the pharmacokinetic results in cancer patients 4 to 18 years of age.

**Table 4. Pharmacokinetics in Pediatric Cancer Patients 1 Month to 18 Years of Age**

Subjects and Age Group	N	CL	V <sub>d<sub>ss</sub></sub>	T <sub>1/2</sub>
		(L/h/kg)	(L/kg)	(h)
		Geometric Mean		Mean
Pediatric Cancer Patients 4 to 18 years of age	N = 21	0.599	1.9	2.8
Population PK Patients <sup>a</sup> 1 month to 48 months of age	N = 115	0.582	3.65	4.9

<sup>a</sup> Population PK (Pharmacokinetic) Patients: 64% cancer patients and 36% surgery patients. Based on the population pharmacokinetic analysis, cancer patients 6 to 48 months of age who receive a dose of 0.15 mg/kg of intravenous ondansetron every 4 hours for 3 doses would be expected to achieve a systemic exposure (AUC) consistent with the exposure achieved in previous pediatric studies in cancer patients (4 to 18 years of age) at similar doses. In a study of 21 pediatric patients (3 to 12 years of age) who were undergoing surgery requiring anesthesia for a duration of 45 minutes to 2 hours, a single intravenous dose of ondansetron, 2 mg (3 to 7 years) or 4 mg (8 to 12 years), was administered immediately prior to anesthesia induction. Mean weight-normalized clearance and volume of distribution values in these pediatric surgical patients were similar to those previously reported for young adults. Mean terminal half-life was slightly reduced in pediatric patients (range, 2.5 to 3 hours) in comparison with adults (range, 3 to 3.5 hours).

In a study of 51 pediatric patients (1 month to 24 months of age) who were undergoing surgery requiring general anesthesia, a single intravenous dose of ondansetron, 0.1 or 0.2 mg/kg, was administered prior to surgery. As shown in Table 5, the 41 patients with pharmacokinetic data were divided into 2 groups, patients 1 month to 4 months of age and patients 5 to 24 months of age, and are compared to pediatric patients 3 to 12 years of age.

**Table 5. Pharmacokinetics in Pediatric Surgery Patients 1 Month to 12 Years of Age**

Subjects and Age Group	N	CL	V <sub>d<sub>ss</sub></sub>	T <sub>1/2</sub>
		(L/h/kg)	(L/kg)	(h)
		Geometric Mean		Mean
Pediatric Surgery Patients 3 to 12 years of age	N = 21	0.439	1.65	2.9
Pediatric Surgery Patients 5 to 24 months of age	N = 22	0.581	2.3	2.9
Pediatric Surgery Patients 1 month to 4 months of age	N = 19	0.401	3.5	6.7

In general, surgical and cancer pediatric patients younger than 18 years tend to have a higher ondansetron clearance compared to adults leading to a shorter half-life in most pediatric patients. In patients 1 month to 4 months of age, a longer half-life was observed due to the higher volume of distribution in this age group.

In a study of 21 pediatric cancer patients (4 to 18 years of age) who received three intravenous doses of 0.15 mg/kg of ondansetron at 4-hour intervals, patients older than 15 years of age exhibited ondansetron pharmacokinetic parameters similar to those of adults.

**Renal Impairment**  
 Due to the very small contribution (5%) of renal clearance to the overall clearance, renal impairment was not expected to significantly influence the total clearance of ondansetron. However, ondansetron mean plasma clearance was reduced by about 41% in patients with severe renal impairment (creatinine clearance < 30 mL/min). This reduction in clearance is variable and was not consistent with an increase in half-life. No reduction in dose or dosing frequency in these patients is warranted.

**Hepatic Impairment**  
 In patients with mild-to-moderate hepatic impairment, clearance is reduced 2-fold and mean half-life is increased to 11.6 hours compared to 5.7 hours in those without hepatic impairment. In patients with severe hepatic impairment (Child-Pugh score of 10 or greater), clearance is reduced 2-fold to 3-fold and apparent volume of distribution is increased with a resultant increase in half-life to 20 hours. In patients with severe hepatic impairment, a total daily dose of 8 mg should not be exceeded.

**13 NONCLINICAL TOXICOLOGY**

**13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**  
 Carcinogenic effects were not seen in 2-year studies in rats and mice with oral ondansetron doses up to 10 and 30 mg/kg per day, respectively (approximately 3.6 and 5.4 times the recommended human intravenous dose of 0.15 mg/kg given three times a day, based on body surface area). Ondansetron was not mutagenic in standard tests for mutagenicity. Oral administration of ondansetron up to 15 mg/kg per day (approximately 3.8 times the recommended human intravenous dose, based on body surface area) did not affect fertility or general reproductive performance of male and female rats.

**14 CLINICAL STUDIES**

The clinical efficacy of ondansetron hydrochloride, the active ingredient of Ondansetron Injection USP, was assessed in clinical trials as described below.

**14.1 Chemotherapy-Induced Nausea and Vomiting Adults**  
 In a double-blind study of three different dosing regimens of Ondansetron Injection USP, 0.015 mg/kg, 0.15 mg/kg, and 0.30 mg/kg, each given three times during the course of cancer chemotherapy, the 0.15-mg/kg dosing regimen was more effective than the 0.015-mg/kg dosing regimen. The 0.30-mg/kg dosing regimen was not shown to be more effective than the 0.15-mg/kg dosing regimen.

**Cisplatin-Based Chemotherapy**  
 In a double-blind study in 28 patients, Ondansetron Injection USP (three 0.15-mg/kg doses) was significantly more effective than placebo in preventing nausea and vomiting induced by cisplatin-based chemotherapy. Therapeutic response was as shown in Table 6.

**Table 6. Therapeutic Response in Prevention of Chemotherapy-Induced Nausea and Vomiting in Single-Day Cisplatin Therapy<sup>a</sup> in Adults**

	Ondansetron Injection USP (0.15 mg/kg x 3)	Placebo	P Value <sup>b</sup>
Number of patients	14	14	
Treatment response			
0 Emetic episodes	2 (14%)	0 (0%)	
1 to 2 Emetic episodes	8 (57%)	0 (0%)	
3 to 5 Emetic episodes	2 (14%)	1 (7%)	
More than 5 emetic episodes/rescued	2 (14%)	13 (93%)	0.001
Median number of emetic episodes	1.5	Undefined <sup>c</sup>	
Median time to first emetic episode (h)	11.6	2.8	0.001
Median nausea scores (0 to 100) <sup>d</sup>	3	59	0.034
Global satisfaction with control of nausea and vomiting (0 to 100) <sup>e</sup>	96	10.5	0.009

<sup>a</sup> Chemotherapy was high dose (100 and 120 mg/m<sup>2</sup>; Ondansetron Injection USP n = 6, placebo n = 5) or moderate dose (50 and 80 mg/m<sup>2</sup>; Ondansetron Injection USP n = 8, placebo n = 9). Other chemotherapeutic agents included fluorouracil, doxorubicin, and cyclophosphamide. There was no difference between treatments in the types of chemotherapy that would account for differences in response.

<sup>b</sup> Efficacy based on "all patients treated" analysis.  
<sup>c</sup> Median undefined since at least 50% of the patients were rescued or had more than five emetic episodes.

<sup>d</sup> Visual analog scale assessment of nausea: 0 = no nausea, 100 = nausea as bad as it can be.  
<sup>e</sup> Visual analog scale assessment of satisfaction: 0 = not at all satisfied, 100 = totally satisfied. Ondansetron Injection USP (0.15-mg/kg x 3 doses) was compared with metoclopramide (2 mg/kg x 6 doses) in a single-blind trial in 307 patients receiving cisplatin ≥100 mg/m<sup>2</sup> with or without other chemotherapeutic agents. Patients received the first dose of ondansetron or metoclopramide 30 minutes before cisplatin. Two additional ondansetron doses were administered 4 and 8 hours later, or five additional metoclopramide doses were administered 2, 4, 7, 10, and 13 hours later. Cisplatin was administered over a period of 3 hours or less. Episodes of vomiting andretching were tabulated over the period of 24 hours after cisplatin. The results of this study are summarized in Table 7.

**Table 7. Therapeutic Response in Prevention of Vomiting Induced by Cisplatin (≥ 100 mg/m<sup>2</sup>) Single-Day Therapy<sup>a</sup> in Adults**

	Ondansetron Injection USP (0.15 mg/kg x 3)	Metoclopramide 2 mg/kg x 6	P Value
Dose	0.15 mg/kg x 3	2 mg/kg x 6	
Number of patients in efficacy population	136	138	
Treatment response			
0 Emetic episodes	54 (40%)	41 (30%)	
1 to 2 Emetic episodes	34 (25%)	30 (22%)	
3 to 5 Emetic episodes	19 (14%)	18 (13%)	
More than 5 emetic episodes/rescued	29 (21%)	49 (36%)	
Comparison of treatments with respect to			
0 Emetic episodes	54/136	41/138	0.083
More than 5 emetic episodes/rescued	29/136	49/138	0.009
Median number of emetic episodes	1	2	0.005
Median time to first emetic episode (h)	20.5	4.3	< 0.001
Global satisfaction with control of nausea and vomiting (0 to 100) <sup>b</sup>	85	63	0.001
Acute dystonic reactions	0	8	0.005
Akathisia	0	10	0.002

<sup>a</sup> In addition to cisplatin, 68% of patients received other chemotherapeutic agents, including cyclophosphamide, etoposide, and fluorouracil. There was no difference between treatments in the types of chemotherapy that would account for differences in response.

<sup>b</sup> Visual analog scale assessment: 0 = not at all satisfied, 100 = totally satisfied.  
**Cyclophosphamide-Based Chemotherapy**  
 In a double-blind, placebo-controlled study of Ondansetron Injection USP (three 0.15-mg/kg doses) in 20 patients receiving cyclophosphamide (500 to 600 mg/m<sup>2</sup>) chemotherapy, Ondansetron Injection USP was significantly more effective than placebo in preventing nausea and vomiting. The results are summarized in Table 8.

**Table 8. Therapeutic Response in Prevention of Chemotherapy-Induced Nausea and Vomiting in Single-Day Cyclophosphamide Therapy<sup>a</sup> in Adults**

	Ondansetron Injection USP (0.15 mg/kg x 3)	Placebo	P Value <sup>b</sup>
Number of patients	10	10	
Treatment response			
0 Emetic episodes	7 (70%)	0 (0%)	0.001
1 to 2 Emetic episodes	0 (0%)	2 (20%)	
3 to 5 Emetic episodes	2 (20%)	4 (40%)	
More than 5 emetic episodes/rescued	1 (10%)	4 (40%)	0.131
Median number of emetic episodes	0	4	0.008
Median time to first emetic episode (h)	Undefined <sup>c</sup>	8.79	
Median nausea scores (0 to 100) <sup>d</sup>	0	60	0.001
Global satisfaction with control of nausea and vomiting (0 to 100) <sup>e</sup>	100	52	0.008

<sup>a</sup> Chemotherapy consisted of cyclophosphamide in all patients, plus other agents, including fluorouracil, doxorubicin, methotrexate, and vincristine. There was no difference between treatments in the type of chemotherapy that would account for differences in response.

<sup>b</sup> Efficacy based on "all patients treated" analysis.  
<sup>c</sup> Median undefined since at least 50% of patients did not have any emetic episodes.  
<sup>d</sup> Visual analog scale assessment of nausea: 0 = no nausea, 100 = nausea as bad as it can be.  
<sup>e</sup> Visual analog scale assessment of satisfaction: 0 = not at all satisfied, 100 = totally satisfied.

**Re-treatment**  
 In uncontrolled trials, 127 patients receiving cisplatin (median dose, 100 mg/m<sup>2</sup>) and ondansetron who had two or fewer emetic episodes were re-treated with ondansetron and chemotherapy, mainly cisplatin, for a total of 269 re-treatment courses (median, 2; range, 1 to 10). No emetic episodes occurred in 160 (59%), and two or fewer emetic episodes occurred in 217 (81%) re-treatment courses.

**Pediatrics**  
 Four open-label, noncomparative (one US, three foreign) trials have been performed with

209 pediatric cancer patients 4 to 18 years of age given a variety of cisplatin or noncisplatin regimens. In the three foreign trials, the initial Ondansetron Injection USP dose ranged from 0.04 to 0.87 mg/kg for a total dose of 2.16 to 12 mg. This was followed by the oral administration of ondansetron ranging from 4 to 24 mg daily for 3 days. In the US trial, Ondansetron Injection USP was administered intravenously (only) in three doses of 0.15 mg/kg each for a total daily dose of 7.2 to 39 mg. In these studies, 58% of the 196 evaluable patients had a complete response (no emetic episodes) on day 1. Thus, prevention of vomiting in these pediatric patients was essentially the same as for patients older than 18 years of age.

An open-label, multicenter, noncomparative trial has been performed in 75 pediatric cancer patients 6 to 48 months of age receiving at least one moderately or highly emetogenic chemotherapeutic agent. Fifty-seven percent (57%) were females; 67% were white, 18% were American Hispanic, and 15% were black patients. Ondansetron Injection USP was administered intravenously over 15 minutes in three doses of 0.15 mg/kg. The first dose was administered 30 minutes before the start of chemotherapy, the second and third doses were administered 4 and 8 hours after the first dose, respectively. Eighteen patients (25%) received routine prophylactic dexamethasone (i.e., not given as rescue). Of the 75 evaluable patients, 56% had a complete response (no emetic episodes) on day 1. Thus, prevention of vomiting in these pediatric patients was comparable to the prevention of vomiting in patients 4 years of age and older.

**14.2 Prevention of Postoperative Nausea and/or Vomiting Adults**  
 Adult surgical patients who received ondansetron immediately before the induction of general balanced anesthesia (barbiturate: thiopental, methohexital, or thiamylal; opioid: alfentanil or fentanyl; nitrous oxide; neuromuscular blockade: succinylcholine/curare and/or vecuronium or atracurium; and supplemental isoflurane) who received no prophylactic antiemetics and who experienced nausea and/or vomiting within 2 hours postoperatively were evaluated in two double-blind US studies involving 441 patients. Patients who experienced an episode of postoperative nausea and/or vomiting were given Ondansetron Injection USP (4 mg) intravenously over 2 to 5 minutes, and this was significantly more effective than placebo. The results of these studies are summarized in Table 9.

**Table 9. Therapeutic Response in Prevention of Postoperative Nausea and Vomiting in Adult Patients**

	Ondansetron 4 mg Intravenous	Placebo	P Value
Study 1			
Emetic episodes:			
Number of patients	136	139	
Treatment response over 24-h postoperative period			
0 Emetic episodes	103 (76%)	64 (46%)	< 0.001
1 Emetic episode	13 (10%)	17 (12%)	
More than 1 emetic episode/rescued	20 (15%)	58 (42%)	
Nausea assessments:			
Number of patients	134	136	
No nausea over 24-h postoperative period	56 (42%)	39 (29%)	
Study 2			
Emetic episodes:			
Number of patients	136	143	
Treatment response over 24-h postoperative period			
0 Emetic episodes	85 (63%)	63 (44%)	0.002
1 Emetic episode	16 (12%)	29 (20%)	
More than 1 emetic episode/rescued	35 (26%)	51 (36%)	
Nausea assessments:			
Number of patients	125	133	
No nausea over 24-h postoperative period	48 (38%)	42 (32%)	

The study populations in Table 9 consisted mainly of females undergoing laparoscopic procedures. In a placebo-controlled study conducted in 468 males undergoing outpatient procedures, a single 4-mg intravenous ondansetron dose prevented postoperative vomiting over a 24-hour study period in 79% of males receiving drug compared to 63% of males receiving placebo (P < 0.001).

Two other placebo-controlled studies were conducted in 2,792 patients undergoing major abdominal or gynecological surgeries to evaluate a single 4-mg or 8-mg intravenous ondansetron dose for prevention of postoperative nausea and vomiting over a 24-hour study period. At the 4-mg dosage, 59% of patients receiving ondansetron versus 45% receiving placebo in the first study (P < 0.001) and 41% of patients receiving ondansetron versus 30% receiving placebo in the second study (P = 0.001) experienced no emetic episodes. No additional benefit was observed in patients who received intravenous ondansetron 8 mg compared to patients who received intravenous ondansetron 4 mg.

**Pediatrics**  
 Three double-blind, placebo-controlled studies have been performed (one US, two foreign) in 1,049 male and female patients (2 to 12 years of age) undergoing general anesthesia with nitrous oxide. The surgical procedures included tonsillectomy with or without adenoidectomy, strabismus surgery, herniorrhaphy, and orchidopexy. Patients were randomized to either single intravenous doses of ondansetron (0.1 mg/kg for pediatric patients weighing 40 kg or less, 4 mg for pediatric patients weighing more than 40 kg) or placebo. Study drug was administered over at least 30 seconds, immediately prior to or following anesthesia induction. Ondansetron was significantly more effective than placebo in preventing nausea and vomiting. The results of these studies are summarized in Table 10.

**Table 10. Therapeutic Response in Prevention of Postoperative Nausea and Vomiting in Pediatric Patients 2 to 12 Years of Age**

	Ondansetron n (%)	Placebo n (%)	P Value
Treatment Response Over 24 Hours			
Study 1			
Number of patients	205	210	
0 Emetic episodes	140 (68%)	82 (39%)	≤ 0.001
Failure <sup>a</sup>	65 (32%)	128 (61%)	
Study 2			
Number of patients	112	110	
0 Emetic episodes	68 (61%)	38 (35%)	≤ 0.001
Failure <sup>a</sup>	44 (39%)	72 (65%)	
Study 3			
Number of patients	206	206	
0 Emetic episodes	123 (60%)	96 (47%)	≤ 0.01
Failure <sup>a</sup>	83 (40%)	110 (53%)	
Nausea assessments <sup>b</sup> :			
Number of patients	185	191	
None	119 (64%)	99 (52%)	≤ 0.01

<sup>a</sup> Failure was one or more emetic episodes, rescued, or withdrawn.  
<sup>b</sup> Nausea measured as none, mild, or severe.

A double-blind, multicenter, placebo-controlled study was conducted in 670 pediatric patients 1 month to 24 months of age who were undergoing routine surgery under general anesthesia. Seventy-five percent (75%) were males; 64% were white, 15% were black, 13% were American Hispanic, 2% were Asian, and 6% were "other race" patients. A single 0.1-mg/kg intravenous dose of ondansetron administered within 5 minutes following induction of anesthesia was statistically significantly more effective than placebo in preventing vomiting. In the placebo group, 28% of patients experienced vomiting compared to 11% of subjects who received ondansetron (P ≤ 0.01). Overall, 32 (10%) of placebo patients and 18 (5%) of patients who received ondansetron received antiemetic rescue medication(s) or prematurely withdrew from the study.

**14.3 Prevention of Further Postoperative Nausea and Vomiting**

**Adults**  
 Adult surgical patients receiving general balanced anesthesia (barbiturate: thiopental, methohexital, or thiamylal; opioid: alfentanil or fentanyl; nitrous oxide; neuromuscular blockade: succinylcholine/curare and/or vecuronium or atracurium; and supplemental isoflurane) who received no prophylactic antiemetics and who experienced nausea and/or vomiting within 2 hours postoperatively were evaluated in two double-blind US studies involving 441 patients. Patients who experienced an episode of postoperative nausea and/or vomiting were given Ondansetron Injection USP (4 mg) intravenously over 2 to 5 minutes, and this was significantly more effective than placebo. The results of these studies are summarized in Table 11.

**Table 11. Therapeutic Response in Prevention of Further Postoperative Nausea and Vomiting in Adult Patients**

	Ondansetron 4 mg Intravenous	Placebo	P Value
Study 1			
Emetic episodes:			
Number of patients	104	117	
Treatment response 24 h after study drug			
0 Emetic episodes	49 (47%)	19 (16%)	< 0.001
1 Emetic episode	12 (12%)	9 (8%)	
More than 1 emetic episode/rescued	43 (41%)	89 (76%)	
Median time to first emetic episode (min) <sup>a</sup>	55	43	
Nausea assessments:			
Number of patients	98	102	
Mean nausea score over 24-h postoperative period <sup>b</sup>	1.7	3.1	
Study 2			
Emetic episodes:			
Number of patients	112	108	
Treatment response 24 h after study drug			
0 Emetic episodes	49 (44%)	28 (26%)	0.006
1 Emetic episode	14 (13%)	3 (3%)	
More than 1 emetic episode/rescued	49 (44%)	77 (71%)	
Median time to first emetic episode (min) <sup>a</sup>	60.5	34	
Nausea assessments:			
Number of patients	105	85	
Mean nausea score over 24-h postoperative period <sup>b</sup>	1.9	2.9	

<sup>a</sup> After administration of study drug.  
<sup>b</sup> Nausea measured on a scale of 0 to 10 with 0 = no nausea, 10 = nausea as bad as it can be.

The study populations in Table 11 consisted mainly of women undergoing laparoscopic procedures.

**Repeat Dosing in Adults**  
 In patients who do not achieve adequate control of postoperative nausea and vomiting following a single, prophylactic, preinduction, intravenous dose of ondansetron 4 mg, administration of a second intravenous dose of ondansetron 4 mg postoperatively does not provide additional control of nausea and vomiting.

**Pediatrics**  
 One double-blind, placebo-controlled, US study was performed in 351 male and female outpatients (2 to 12 years of age) who received general anesthesia with nitrous oxide and no prophylactic antiemetics. Surgical procedures were unrestricted. Patients who experienced two or more emetic episodes within 2 hours following discontinuation of nitrous oxide were randomized to either single intravenous doses of ondansetron (0.1 mg/kg for pediatric patients weighing 40 kg or less, 4