

0.750

1.750

1.583

NO COATING

NC

NC

NC

NC

NC

NO COAT

NC

NO COATING

NC

NC

NC

NC

**Drug Facts (continued)**

**Other information**

- each capsule contains sodium 340 mg
- read the directions, warnings and accompanying label information before use.
- store at room temperature between 20° to 25°C (68° to 77°F)
- tamper evident: do not use if the blue band around the capsule is missing or broken. Do not use if foil inner seal imprinted with "Sealed for your protection" is missing, open or broken.
- keep product out of high heat and humidity
- avoid product from moisture

**inactive ingredients**

croscarmellose sodium, FD&C blue #1, gelatin, polyacrylate 90, sodium lauryl sulfate, sodium steryl fumarate and titanium dioxide. Printing ink contains black iron oxide, butyl alcohol, dehydrated alcohol, isopropyl alcohol, potassium hydroxide, polyethylene glycol, purified water, shellac and strong ammonia solution.

**Questions?** 1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.

\*This product is not affiliated with, manufactured by or produced by the makers or owners of Zegerid OTC®.

Manufactured for:  
Actavis Pharma, Inc.  
Parsippany, NJ 07054 USA

Made in India      DP19537  
04410216              Rev. 02/16

NDC 45963-441-40

Compare to the active ingredients in Zegerid OTC®.\*

**Treats Frequent Heartburn**

## Omeprazole and Sodium Bicarbonate Capsules

**20 mg/1100 mg**

- Omeprazole 20 mg/Acid Reducer
- Sodium Bicarbonate 1100 mg/Allows Absorption of this Omeprazole Product

**One-14-Day Course of Treatment**

**14 capsules**

**How Omeprazole and Sodium Bicarbonate Capsules Work For Your Frequent Heartburn**

Omeprazole and sodium bicarbonate capsules work differently from other OTC heartburn products, such as antacids and other acid reducers. Omeprazole and sodium bicarbonate capsules stop acid production at the source—the acid pump that produces stomach acid. Omeprazole and sodium bicarbonate capsules are to be used once a day (every 24 hours), every day for 14 days.

**Tips for Managing Heartburn**

- Do not lie flat or bend over soon after eating.
- Do not eat late at night or just before bedtime.
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some fruits and vegetables.
- Eat slowly and do not eat big meals.
- If you are overweight, lose weight.
- If you smoke, quit smoking.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.



**Drug Facts**

**Active ingredients (in each capsule)**

Omeprazole, USP 20 mg      Acid reducer

Sodium Bicarbonate, USP 1100 mg      Allows absorption of this omeprazole product

**Use**

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.

**Warnings**

**Allergy alert:** Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food
- vomiting with blood
- bloody or black stools

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- bad heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- trouble or pain swallowing food
- unexplained weight loss
- nausea or vomiting
- stomach pain
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking:

- warfarin, clopidogrel or clostazolol (blood thinning medicines)
- prescription antifungal or anti-yeast medicines
- disopyramide (antiarrhythmic medicine)
- digoxin (heart medicine)
- baclofen or mycophenolate mofetil (immune system medicines)

**Drug Facts (continued)**

- prescription antiretrovirals (medicines for HIV infection)
- metformin (diabetes medicine)
- any other prescription drugs. Sodium bicarbonate may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**

- swallow 1 capsule with a glass of water at least 1 hour before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- do not chew or crush the capsule
- do not open capsule and sprinkle on food
- do not use for more than 14 days unless directed by your doctor
- repeat 14-Day Course (if needed)
- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

**FONTS:** Myriad Pro – roman, bold, bold italic, black, black italic; Helvetica condensed medium, condensed bold, condensed bold oblique.

**Drug Facts:** 10 pt

**Headers:** 8 pt

**Body text:** 6 pt

**Drug Facts (continued):** 8 pt

**Leadings:** 6.5 pt

**Bullets:** 6 pt

**Frames:** 1.5 pt

**Bars:** 1 pt

**Rules:** 0.75 pt



## LEAFLET INSIDE PRINTING

## LEAFLET OUTSIDE PRINTING

<p><b>Drug Facts (continued)</b></p> <p><b>Use</b></p> <ul style="list-style-type: none"> <li>■ treats frequent heartburn (occurs 2 or more days a week)</li> <li>■ not intended for immediate relief of heartburn. This drug may take 1 to 4 days for full effect.</li> </ul> <p><b>Warnings</b></p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> <li>■ trouble or pain swallowing food</li> </ul>	<div style="border: 1px solid black; height: 40px;"></div>
--	--

Inside Panel  
PAGE 2

Inside Panel  
PAGE 15

<div style="border: 1px solid black; height: 40px;"></div>	<p style="font-size: 8pt;">NDC 45963 441 40</p> <p><b>Omeprazole and Sodium Bicarbonate</b> Capsules 20 mg/1100 mg</p> <p style="font-size: 8pt;">*Omeprazole 20 mg/Acid Reducer *Sodium Bicarbonate 1100 mg/Acid Reducer of this Omeprazole Product</p> <p style="background-color: #0070C0; color: white; padding: 2px; text-align: center;"><b>One 14-Day Course of Treatment</b></p> <p style="text-align: center; font-weight: bold;">14 capsules</p> <p style="font-size: 8pt;">Open to read directions and additional drug facts information</p> <p><b>Drug Facts</b></p> <table border="1" style="font-size: 8pt; border-collapse: collapse;"> <thead> <tr> <th style="text-align: left;">Active ingredients (in each capsule)</th> <th style="text-align: left;">Purpose</th> </tr> </thead> <tbody> <tr> <td>Omeprazole, USP 20 mg</td> <td>Acid reducer</td> </tr> <tr> <td>Sodium Bicarbonate, USP 1100 mg</td> <td>Acid reducer</td> </tr> <tr> <td colspan="2" style="font-size: 7pt;">Allows absorption of this omeprazole product</td> </tr> </tbody> </table>	Active ingredients (in each capsule)	Purpose	Omeprazole, USP 20 mg	Acid reducer	Sodium Bicarbonate, USP 1100 mg	Acid reducer	Allows absorption of this omeprazole product	
Active ingredients (in each capsule)	Purpose								
Omeprazole, USP 20 mg	Acid reducer								
Sodium Bicarbonate, USP 1100 mg	Acid reducer								
Allows absorption of this omeprazole product									

Outside Panel  
PAGE 16

Outside Panel  
PAGE 1

<p>How are Omeprazole and Sodium Bicarbonate Capsules Sold</p> <p>Omeprazole and sodium bicarbonate capsules are available in 14, 28 and 42 count sizes. These sizes contain one, two and three 14-day courses of treatment, respectively. Do not use for more than 14 days in a row unless directed by your doctor. For the 28 (two 14-day courses) and 42 count (three 14-day courses), you may repeat a 14-day course every 4 months.</p>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ vomiting with blood</li> <li>■ bloody or black stools</li> </ul> <p>These may be signs of a serious condition. See your doctor.</p> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>■ had heartburn over 3 months. This may be a sign of a more serious condition.</li> <li>■ heartburn with lightheadedness, sweating or dizziness</li> </ul>
--	---

Inside Panel  
PAGE 14

Inside Panel  
PAGE 3

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness</li> <li>■ frequent chest pain</li> <li>■ frequent wheezing, particularly with heartburn</li> <li>■ unexplained weight loss</li> <li>■ nausea or vomiting</li> <li>■ stomach pain</li> <li>■ a sodium-restricted diet</li> </ul>	<p style="text-align: center; font-size: 8pt;"><b>Tips for Managing Heartburn</b></p> <ul style="list-style-type: none"> <li>■ Do not lie flat or bend over soon after eating.</li> <li>■ Do not eat late at night or just before bedtime.</li> <li>■ Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some fruits and vegetables.</li> <li>■ Eat slowly and do not eat big meals.</li> <li>■ If you are overweight, lose weight.</li> <li>■ If you smoke, quit smoking.</li> <li>■ Raise the head of your bed.</li> <li>■ Wear loose-fitting clothing around your stomach.</li> </ul>
--	---

Outside Panel  
PAGE 4

Outside Panel  
PAGE 13

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ any other prescription drugs. Sodium bicarbonate may interact with certain prescription drugs.</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ your heartburn continues or worsens</li> <li>■ you need to take more than 14 days</li> <li>■ you need to take more than 1 course of treatment every 4 months</li> <li>■ you get diarrhea</li> </ul> <p><b>If pregnant or breast-feeding, ask a health professional before use.</b></p>	<p><b>Drug Facts (continued)</b></p> <p><b>Inactive ingredients</b></p> <p>croscarmellose sodium, FD&amp;C blue #1, gelatin, polyacrylate 80, sodium lauryl sulfate, sodium stearoyl fumarate and titanium dioxide. Printing ink contains black iron oxide, butyl alcohol, dehydrated alcohol, isopropyl alcohol, potassium hydroxide, propylene glycol, purified water, shellac and strong ammonia solution.</p> <p><b>Questions? 1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.</b></p>
--	---

Inside Panel  
PAGE 6

Inside Panel  
PAGE 11

<p>How Omeprazole and Sodium Bicarbonate Capsules Work For Your Frequent Heartburn</p> <p>Omeprazole and sodium bicarbonate capsules work differently from other OTC heartburn products, such as antacids and other acid reducers. Omeprazole and sodium bicarbonate capsules stop acid production at the source—the acid pump that produces stomach acid. Omeprazole and sodium bicarbonate capsules are to be used once a day (every 24 hours), every day for 14 days.</p> <p><b>What to Expect When Using Omeprazole and Sodium Bicarbonate Capsules</b></p> <p>Omeprazole and sodium bicarbonate capsules are a different type of medicine from antacids and other acid reducers. Omeprazole and sodium bicarbonate capsules may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours. Make sure you take the entire 14 days of dosing to treat your frequent heartburn.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor or pharmacist before use if you are taking</b></p> <ul style="list-style-type: none"> <li>■ warfarin, clopidogrel or cloacizol (blood-thinning medicines)</li> <li>■ prescription antifungal or anti-yeast medicines</li> <li>■ diazepam (anxiety medicine)</li> <li>■ digoxin (heart medicine)</li> <li>■ tacrolimus or mycophenolate mofetil (immune system medicines)</li> <li>■ prescription antiretrovirals (medicines for HIV infection)</li> <li>■ methotrexate (arthritis medicine)</li> </ul>
--	--

Outside Panel  
PAGE 12

Outside Panel  
PAGE 5

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ store at room temperature between 20° to 25°C (68° to 77°F)</li> <li>■ tamper evident: do not use if the blue band around the capsule is missing or broken. Do not use if foil inner seal imprinted with "Sealed for your protection" is missing, open or broken.</li> <li>■ keep product out of high heat and humidity</li> <li>■ protect product from moisture</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ for adults 18 years of age and older</li> <li>■ this product is to be used once a day (every 24 hours), every day for 14 days</li> <li>■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours</li> </ul> <p><b>14-Day Course of Treatment</b></p>
---	---

Inside Panel  
PAGE 10

Inside Panel  
PAGE 7

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ swallow 1 capsule with a glass of water at least 1 hour before eating in the morning</li> <li>■ take every day for 14 days</li> <li>■ do not take more than 1 capsule a day</li> <li>■ do not chew or crush the capsule</li> <li>■ do not open capsule and sprinkle on food</li> <li>■ do not use for more than 14 days unless directed by your doctor</li> </ul> <p><b>Repeated 14-Day Courses (if needed)</b></p> <ul style="list-style-type: none"> <li>■ you may repeat a 14-day course every 4 months</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</li> <li>■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ each capsule contains: sodium 303 mg</li> <li>■ read the directions, warnings and accompanying label information before use.</li> </ul>
---	--

Outside Panel  
PAGE 8

Outside Panel  
PAGE 9

<p><b>Treats Frequent Heartburn</b></p> <p><b>Omeprazole and Sodium Bicarbonate Capsules</b></p> <p style="background-color: #0070C0; color: white; padding: 2px; text-align: center;"><b>20 mg/1100 mg</b></p> <p style="font-size: 8pt;">*Omeprazole 20 mg/Acid Reducer *Sodium Bicarbonate 1100 mg/ Allows Absorption of this Omeprazole Product</p>	<p style="font-size: 8pt;">NDC 45963-441-40</p> <p style="text-align: center;"><b>One-14-Day Course of Treatment</b></p> <p style="text-align: center;">Manufactured for: Actavis Pharma, Inc. Parsippany, NJ 07054 USA</p> <p style="text-align: center;">Made in India</p> <p style="font-size: 8pt;">Rev. 02/16 DPI19540</p> <p style="text-align: center;"><b>14 capsules</b>      04410216</p>	<div style="border: 1px solid black; height: 40px;"></div>
---	---	--

BASE LABEL With Liner

<b>FONTS:</b> Myriad Pro – roman, semibold, semibold italic, bold italic, black, black italic, black condensed; Helvetica condensed medium, condensed bold, condensed bold oblique.	
<b>Drug Facts:</b>	10 pt
<b>Headers:</b>	8 pt
<b>Body text:</b>	6 pt
<b>Drug Facts (continued):</b>	8 pt
<b>Leading:</b>	7 pt
<b>Bullets:</b>	6 pt
<b>Frames:</b>	1 pt
<b>Bars:</b>	0.5 pt
<b>Rules:</b>	0.5 pt