

090177e18954c2ec\Final\Final On: 01-Aug-2016 03:21

PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	N/A
• SUBHEADINGS/BODY TEXT	7 pt
• LEADING	7.5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	N/A




ea\Final\Final On: 01-Aug-2016 03:22




PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces





8954bfc2\Final\Final On: 01-Aug-2016 03:17


PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	(b) (4)



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	.5 pt



090177e18954bfbf\Final\On: 01-Aug-2016 03:14

PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

090177e18954c382\Final\Final On: 01-Aug-2016 03:18

UNCOATED
NO COPY
NO PRINT

128 Code = 075009

128 CODE AREA

128 Code = 075009

128 CODE AREA

128 Code = 075009

128 CODE AREA

MICRO PRINT LOCATION

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts
Active ingredient (in each capsule) Purpose
Esomeprazole 20 mg Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain
These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting ■ stomach pain
Ask a doctor or pharmacist before use if you are taking
■ warfarin, clopidogrel or clobazam (blood-thinning medicines)
■ prescription antifungal or anti-yeast medicines
■ digoxin (heart medicine)
■ diazepam (anxiety medicine)
■ tacrolimus or mycophenolate mofetil (immune system medicines)
■ prescription antiretrovirals (medicines for HIV infection)
■ methotrexate (arthritis medicine)

Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

Drug Facts (continued)
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea
If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ may take 1 to 4 days for full effect
14-Day Course of Treatment
■ swallow 1 capsule with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 capsule a day
■ swallow whole. Do not crush or chew capsules.
■ do not use for more than 14 days unless directed by your doctor
Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F)

Inactive ingredients
corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments?
call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600

Tips for Managing Heartburn
• Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
• Eat slowly and do not eat big meals.
• Do not eat late at night or just before bedtime.
• Do not lie flat or bend over soon after eating.
• Raise the head of your bed.
• Wear loose-fitting clothing around your stomach.
• If you are overweight, lose weight.
• If you smoke, quit smoking.

Nexium 24HR
esomeprazole magnesium delayed-release capsules 20 mg/acid reducer

Marketed by:
Pfizer, Madison, NJ 07940 USA
© 2016 Pfizer Inc.
Made in France
For most recent product information, visit www.Nexium24HR.com
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

Nexium 24HR
esomeprazole magnesium delayed-release capsules 20 mg/acid reducer

28 CAPSULES
Two 14-day courses of treatment

May take 1 to 4 days for full effect

2 BOTTLES INSIDE

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

ACTUAL SIZE

3 0573 2450 28 5

128 CODE AREA

MICRO PRINT LOCATION

128 Code = 075009

LOT & EXP AREA. DARK COLOR FOR LASER PRINTING.

3/16" DATA MATRIX CODE WITH 1/8" QUIET ZONE ON ALL SIDES
DATA MATRIX CODE = 075009

128 CODE AREA

128 Code = 075009

128 CODE AREA

128 Code = 075009

128 CODE AREA - SUBSET C 10 MIL DENSITY WITH 4 OR 6 DIGITS (CODE TO EXTEND TO EDGE OF FLAP - UNVARNISHED) PRINTED SIDE OF CARTON
128 Code = 075009

PRODUCTION (NOT COLOR ACCURATE)

Job Information	Color Legend

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



090177e18954c082\Final\Final On: 01-Aug-2016-03:17

128 CODE AREA - SUBSET C 10 MIL DENSITY WITH 4 OR 6 DIGITS (CODE TO EXTEND TO EDGE OF FLAP - UNVARNISHED) PRINTED SIDE OF CARTON
128 Code = 072186

128 Code = 072186


LOT & EXP AREA. DARK COLOR FOR LASER PRINTING.
3/16" DATA MATRIX CODE WITH 1/8" QUIET ZONE ON ALL SIDES
DATA MATRIX CODE = 072186

128 Code = 072186

PRODUCTION (NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



NDC 0573-2450-15

Treats Frequent Heartburn

Nexium[®]
esomeprazole magnesium
delayed-release capsules
20 mg/acid reducer

May take 1 to 4 days for full effect

14 CAPSULES
One 14-day course of treatment

Capsules

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Marketed by: Pfizer, Madison, NJ 07940 USA © 2016 Pfizer Inc. Made in France
For most recent product information, visit www.Nexium24hr.com

Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts

Active ingredient (in each capsule) ... Acid reducer
Esomeprazole 20 mg ...
(each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain
These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have
■ heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss ■ nausea or vomiting
■ stomach pain

Ask a doctor or pharmacist before use if you are taking
■ warfarin, clopidogrel or cizoset (blood-thinning medicines)
■ prescription antifungal or anti-yeast medicines
■ digoxin (heart medicine)
■ diazepam (anxiety medicine)
■ tacrolimus or mycophenolate mofetil (immune system medicines)
■ prescription antiviral (medicines for HIV infection)
■ methotrexate (arthritis medicine)
Stop use and ask a doctor if
■ your heartburn continues or worsens
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea

Other information
■ read the directions and warnings before use
■ keep the card. It contains important information.
■ store at 20-25°C (68-77°F)
Inactive ingredients corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polyethylene glycol 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600

LOT/EXP
WHITE HOLDOUT

REMOVE VARNISH IN THIS AREA ONLY

DATA MATRIX CODE = 054223

DATA MATRIX CODE

WHITE HOLDOUT QUIET AREA

3 0573 2450 15 5
FPO

090177e18954c37f\Final\Final On: 01-Aug-2016 03:15

PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

NDC 0573-2450-29

Treats Frequent Heartburn

Nexium[®]

esomeprazole magnesium
delayed-release capsules
20 mg/acid reducer

24HR

May take 1 to 4 days for full effect


2 Pack

Capsules



28
CAPSULES

Two 14-day courses
of treatment



00000054539 FDA02

WHITE HOLDOUT
QUIET AREA
DATA MATRIX CODE
054539

REMOVE VARNISH
WHITE HOLDOUT
LOT/EXP

Marketed by: Pfizer, Madison, NJ 07940 USA
© 2016 Pfizer Inc.
Made in France
For most recent product information,
visit www.Nexium24HR.com
Nexium and the colors purple and gold as
applied to the capsule are registered trademarks
of AstraZeneca AB and are used under license.



Drug Facts
Active ingredient (in each capsule) Esomeprazole magnesium (esomeprazole magnesium trihydrate) 20 mg. (Each delayed-release capsule corresponds to 2.3 mg esomeprazole magnesium trihydrate)
Purpose Acid reducer

Uses
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
■ **Do not use if you have:**
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with **lightheadedness, sweating or dizziness**
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent **chest pain**
These may be signs of a serious condition. See your doctor.
■ **Ask a doctor before use if you have**
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Ask a doctor or pharmacist before use if you are taking
■ warfarin, clopidogrel or clostazolol (blood-thinning medicines)
■ prescription antifungal or anti-yeast medicines
■ digoxin (heart medicine)
■ dexamethasone (steroid medicine)
■ tacrolimus or mycophenolate mofetil (immune system medicines)
■ prescription antiviral (medicines for HIV infection)
■ methotrexate (arthritis medicine)

Stop and ask a doctor if
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

Questions or comments?
call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600

Drug Facts (continued)
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ may take 1 to 4 days for full effect

14-Day Course of Treatment
■ swallow 1 capsule with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 capsule a day
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain
These may be signs of a serious condition. See your doctor.
■ **Ask a doctor before use if you have**
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Other Information
■ read the directions and warnings before use
■ keep the card. It contains important information.
■ store at 20-25°C (68-77°F)

Inactive ingredients
corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea

KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Nexium[®]

esomeprazole magnesium
delayed-release capsules
20 mg/acid reducer

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

Tips for Managing Heartburn
• Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
• Eat slowly and do not eat big meals.
• Do not eat late at night or just before bedtime.
• Do not lie flat or bend over soon after eating.
• Raise the head of your bed.
• Wear loose-fitting clothing around your stomach.
• If you are overweight, lose weight.
• If you smoke, quit smoking.

PRODUCTION


(NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	10 pt
• DRUG FACTS CONTINUED	9 pt
• HEADINGS	9 pt
• SUBHEADINGS/BODY TEXT	7 pt
• LEADING	7.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	6 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

090177e18954cab4\Final\Final On: 01-Aug-2016 03:18

FRONT



NDC 0573-2450-43

Treats Frequent Heartburn

Nexium[®]

esomeprazole magnesium
delayed-release capsules
20 mg/acid reducer

24HR

May take 1 to 4 days for full effect

3 Pack Capsules

42 CAPSULES
Three 14-day courses of treatment

REMOVE VARNISH
IN THIS AREA ONLY

WHITE HOLDOUT
QUIET AREA

DATA MATRIX CODE
054222

REMOVE VARNISH
WHITE HOLDOUT
LOT/EXP

Marketed by: Pfizer, Madison, NJ 07940 USA
© 2016 Pfizer Inc.
Made in France
For most recent product information,
visit www.Nexium24HR.com
Nexium and the colors purple and gold as
applied to the capsule are registered trademarks
of AstraZeneca AB and are used under license.



KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts
Active ingredient (in each capsule) Esomeprazole 20 mg (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)
Purpose Acid reducer

Uses
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
■ frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Ask a doctor before use if you are taking:
■ warfarin, clopidogrel or clostazol (blood-thinning medicines)
■ prescription antifungal or anti-yeast medicines
■ disease (heart medicine)
■ tacrolimus or mycophenolate mofetil (immune system medicines)
■ prescription antiretrovirals (medicines for HIV infection)
■ methotrexate (arthritis medicine)

Stop use and ask a doctor if:
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

Questions or comments?
call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600

Other information
■ children may sometimes be caused by a serious condition.
■ 4 months unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Inactive ingredients
■ store at 20-25°C (68-77°F)
■ read the directions and warnings before use
■ keep the card. It contains important information.

Directions
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ may take 1 to 4 days for full effect
■ swallow 1 capsule with a glass of water before eating in the morning

14-Day Course of Treatment
■ take every day for 14 days
■ do not take more than 1 capsule a day
■ swallow whole. Do not crush or chew capsules.
■ do not use for more than 14 days unless directed by your doctor

Repeat 14-Day Courses (if needed)
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea

Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not lie flat or bend over soon after eating.
- Do not eat late at night or just before bedtime.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Nexium[®]
esomeprazole magnesium
delayed-release capsules
20 mg/acid reducer
24HR

PRINT SIDE

BACK



PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	10 pt
• DRUG FACTS CONTINUED	9 pt
• HEADINGS	9 pt
• SUBHEADINGS/BODY TEXT	7 pt
• LEADING	7.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	6 pt
• SPACE BEFORE BULLET	N/A
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

090177e18954c084\Final\Final On: 01-Aug-2016 03:20



8954c03d\Final\Final On: 01-Aug-2016 03:13

PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	.5 pt



090177e18954c385\Final\Final On: 01-Aug-2016 03:18

128 CODE AREA - SUBSET C 10 MIL DENSITY WITH 4 OR 6 DIGITS (CODE TO EXTEND TO EDGE OF FLAP - UNVARNISHED) PRINTED SIDE OF CARTON
128 Code = 072959

128 Code = 072959

LOT & EXP AREA. DARK COLOR FOR LASER PRINTING.
3/16" DATA MATRIX CODE WITH 1/8" QUIET ZONE ON ALL SIDES
DATA MATRIX CODE = 072959

128 Code = 072959



(b) (4)

PRODUCTION (NOT COLOR ACCURATE)

Job Information	Color Legend

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces