



090177e18c5541b0\Final\Final On: 13-Jul-2017 20:01 (GMT)



PRODUCTION

(NOT COLOR ACCURATE)






(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8.0 pt
• DRUG FACTS CONTINUED	7.0 pt
• HEADINGS	7.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.0 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt., 5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces






(b) (4)



PRODUCTION (NOT COLOR ACCURATE)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• HAIRLINES	.5 pt

090177e18c55424\Final\Final On: 13-Jul-2017 20:06 (GMT)



PRODUCTION

(NOT COLOR ACCURATE)

090177e18c596c26\Final\Final On: 17-Jul-2017 16:13 (GMT)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• HAIRLINES	.5 pt

Report for: PAA090866_419684.pdf

Page: 1



PRODUCTION

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.0 pt
• DRUG FACTS CONTINUED	8.0 pt
• HEADINGS	8.0 pt
• SUBHEADING/BODY TEXT	6.5 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.0 pt
• SPACE BEFORE BULLET	2 ems
• HORIZONTALS, HORIZONTALS	1.5 pt, 5 pt
• SPACE BETWEEN HORIZONTALS AND BOX END	2 spaces

(b) (4)

090177e18c5541b3\Final\Final On: 13-Jul-2017 20:03 (GMT)

(b) (4)

NDC 0573-2450-15
See new warning

Treats Frequent Heartburn

Nexium[®]
esomeprazole magnesium
delayed-release capsules
20 mg/acid reducer

May take 1 to 4 days
for full effect

14 CAPSULES
One 14-day course
of treatment

Capsules

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

REMOVE VARNISH IN THIS AREA ONLY

DATA MATRIX CODE = 58960

DATA MATRIX CODE

WHITE HOLDOUT QUIET AREA

LOT/EXP
WHITE HOLDOUT

Questions or comments? call toll-free
weekdays 9 AM to 5 PM EST at 1-866-226-1600

Drug Facts (continued)

Active ingredient (in each capsule) Purpose
Esomeprazole 20 mg Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses
• treats frequent heartburn (occurs 2 or more days a week)
• not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Do not use if you have:
• trouble or pain swallowing food, vomiting with blood, or bloody or black stools
• heartburn with **lightheadedness, sweating or dizziness**
• chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
• frequent **chest pain**
These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have
• had heartburn over 3 months. This may be a sign of a more serious condition.
• frequent wheezing, particularly with heartburn
• unexplained weight loss
• nausea or vomiting
• stomach pain

Ask a doctor or pharmacist before use if you are taking
• warfarin, clopidogrel or clostazolol (blood-thinning medicines)
• prescription antifungal or anti-viral medicines
• digoxin (heart medicine) ■ diazepam (anxiety medicine)
• tacrolimus or mycophenolate mofetil (immune system medicines)
• prescription antiretroviral (medicines for HIV infection)
• methotrexate (arthritis medicine)

Stop use and ask a doctor if
• your heartburn continues or worsens
• you need to take this product for more than 14 days
• you need to take more than 1 course of treatment every 4 months
• you get diarrhea
• you develop a rash or joint pain

Marketed by: Pfizer, Madison, NJ 07940 USA © 2017 Pfizer Inc. Made in France
For most recent product information, visit www.Nexium24HR.com

D000058960

Directions
• adults 18 years of age and older
• this product is to be used once a day (every 24 hours), every day for 14 days
• may take 1 to 4 days for full effect

14-Day Course of Treatment
• swallow 1 capsule with a glass of water before eating in the morning
• take every day for 14 days
• do not take more than 1 capsule a day
• do not crush or chew capsules.
• swallow whole. Do not crush or chew capsules.
• do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
• you may repeat a 14-day course every 4 months
• do not take for more than 14 days or more often than every 4 months unless directed by a doctor
• children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information
• read the directions and warnings before use
• keep the card. It contains important information.
• store at 20-25°C (68-77°F)

Inactive ingredients
hydroxypropyl cellulose, hypromellose, magnesium stearate, no. 1, FD&C red no. 40, ferric oxide gelatin, glyceryl monostearate, polyacrylic acid copolymer, pharmaceutical ink, triethyl citrate

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions
• adults 18 years of age and older
• this product is to be used once a day (every 24 hours), every day for 14 days
• may take 1 to 4 days for full effect

14-Day Course of Treatment
• swallow 1 capsule with a glass of water before eating in the morning
• take every day for 14 days
• do not take more than 1 capsule a day
• do not crush or chew capsules.
• swallow whole. Do not crush or chew capsules.
• do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
• you may repeat a 14-day course every 4 months
• do not take for more than 14 days or more often than every 4 months unless directed by a doctor
• children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information
• read the directions and warnings before use
• keep the card. It contains important information.
• store at 20-25°C (68-77°F)

Inactive ingredients
hydroxypropyl cellulose, hypromellose, magnesium stearate, no. 1, FD&C red no. 40, ferric oxide gelatin, glyceryl monostearate, polyacrylic acid copolymer, pharmaceutical ink, triethyl citrate

Questions or comments? call toll-free
weekdays 9 AM to 5 PM EST at 1-866-226-1600

LOT/EXP
WHITE HOLDOUT

REMOVE VARNISH IN THIS AREA ONLY

DATA MATRIX CODE = 58960

DATA MATRIX CODE

WHITE HOLDOUT QUIET AREA

KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Do Not Use If seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

FPO

3 0573 2450 15 5

PRODUCTION


(NOT COLOR ACCURATE)

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9.0 pt
• DRUG FACTS CONTINUED	8.0 pt
• HEADINGS	8.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.0 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

(b) (4)

090177e18c59d237\Final\Final On: 17-Jul-2017 20:44 (GMT)





090177e18c5d818\Final\Final On: 19-Jul-2017 16:12 (GMT)

PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)

DRUG FACTS MODIFIED TEXT DEFINED	TYPE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5
• # OF CHARACTERS PER INCH	<3
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



090177e18c658815\Final\Final On: 24-Jul-2017 19:22 (GMT)



(b) (4)

PRODUCTION (NOT COLOR ACCURATE)




(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	.5 pt

Report for: PAA091522_422319.pdf

Page: 1



PRODUCTION (NOT COLOR ACCURATE)




(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
+DRUG FACTS TITLE	9 pt
+DRUG FACTS CONTINUED	8 pt
+HEADINGS	9 pt
+SUBHEADING/BODY TEXT	6 pt
+LEADINS	6.5 pt
+# OF CHARACTERS PER INCH	<9
+BULLETS	5 pt
+SPACE BEFORE BULLET	2 ems
+BULLETS, HORIZONTALS	1.5 pt, 5 pt
+SPACE BETWEEN HORIZONTALS AND BOX END	2 spaces

090177e18c5bcd4Final\Final On: 18-Jul-2017 19:59 (GMT)

Report for: PAA091521_420892.pdf

(b) (4)



DRUG FACTS TEST DEFINED

• DRUG FACTS TITLE	
• DRUG FACTS SUBTITLE	
• HEADINGS	
• SUB-HEADINGS/TEXT	
• LOGOS	
• # OF CHARACTERS PER INCH	12PT
• BULLETS	2mm
• SPACE BEFORE BULLET	1.5pt, 5pt
• BARRIERS (VERTICALS AND HORIZONTALS)	1 space

090177e18c59d23c\Final\Final On: 17-Jul-2017 21:05 (GMT)

FRONT



REMOVE VARNISH IN THIS AREA ONLY
WHITE HOLDOUT QUIET AREA
DATA MATRIX CODE 59102

PRINT SIDE

BACK

(b) (4)

090177e18c5541af\Final\Final On: 13-Jul-2017 19:58 (GMT)

PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	10 pt
• DRUG FACTS CONTINUED	6 pt
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	7 pt
• LEADING	7.5 pt
• # OF CHARACTERS PER INCH	60
• BULLETS	10 pt
• SPACE BEFORE BULLET	2
• BARLINES, HAIRLINES	5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	spaces 2

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
01/24/2018