


000079729

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	N/A
• SUBHEADINGS/BODY TEXT	7 pt
• LEADING	7.5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	N/A



000079731

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	N/A
• SUBHEADINGS/BODY TEXT	7 pt
• LEADING	7.5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	N/A



Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain ■ if pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect.

14-Day Course of Treatment ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor.

Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20°-25°C (68°-77°F).

Questions or comments? call to 1-Free weekdays 9 AM to 5 PM EST at 1-888-228-1000

Active ingredient (in each capsule) **Purpose**
Esomeprazole 20 mg Acid reducer
(Each delayed-release capsule corresponds to 22.9 mg esomeprazole magnesium trihydrate)

Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.

Warnings
Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash ■ If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain. These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain.

Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.



Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain
- if pregnant or breast-feeding, ask a health professional before use.

 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment:

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor.

Repeated 14-Day Courses (if needed):

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use.

Heartburn in children may sometimes be caused by a serious condition.

Other information:

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)

Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-800-226-1800

Active ingredient (in each capsule)	Purpose
Esomeprazole 20 mg	Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)	
Uses: <ul style="list-style-type: none"> treats frequent heartburn (occurs 2 or more days a week) not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect 	
Warnings:	
Allergy alert: <ul style="list-style-type: none"> Do not use if you are allergic to esomeprazole. Esomeprazole may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> skin redness blisters rash If an allergic reaction occurs, stop use and seek medical help right away. 	
Do not use if you have: <ul style="list-style-type: none"> trouble or pain swallowing food, vomiting with blood, or bloody or black stools heartburn with lightheadedness, swelling or dizziness chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness frequent chest pain. These may be signs of a serious condition. See your doctor. 	
Ask a doctor before use if you have: <ul style="list-style-type: none"> had heartburn over 3 months. This may be a sign of a more serious condition. 	
<ul style="list-style-type: none"> frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain 	
Ask a doctor or pharmacist before use if you are: <ul style="list-style-type: none"> taking a prescription drug. Acid reducers may interact with certain prescription drugs. 	



THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
 Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.
 Dist. by: GSK CH, Warren, NJ 07059
 ©2022 GSK or licensor
 Made in France
 For most recent product information, visit www.Nexium24HR.com
 000078386

Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain ■ if pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect.
14-Day Course of Treatment ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor.
Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
Other Information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20°-25°C (68°-77°F).
Questions or comments? call 1-866-weekdays 9 AM to 5 PM EST at 1-888-226-1800

Active ingredient (in each capsule) **Purpose**
 Esomeprazole 20 mg Acid reducer
 (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)
Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.
Warnings
Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash ■ If an allergic reaction occurs, stop use and seek medical help right away.
 Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain. These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain
Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

<p>Drug Facts</p> <p>Active ingredient (in each capsule) Purpose Esomeprazole 20 mg Acid reducer (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Uses</p> <ul style="list-style-type: none"> ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect. <p>Warnings</p> <p>Allergy alerts</p> <ul style="list-style-type: none"> ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away. <p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <p>Ask a doctor or pharmacist before use if you are:</p> <ul style="list-style-type: none"> ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs. <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ you get diarrhea ■ you develop a rash or joint pain <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor. <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>Children under 18 years of age: ask a doctor before use.</p> <p>Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other Information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F) <p>Inactive ingredients</p> <p>corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methylacrylic acid copolymer, pharmaceutical ink, polyacrylate 80, sucrose, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1800</p>
---	--



Nexium[®] 24HR
 esomeprazole magnesium
 delayed-release capsules
 20 mg/acid reducer

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
 ©2022 GSK group of companies or its licensor
 Made in France
 For most recent product information, visit www.Nexium24HR.com
 Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

14 CAPSULES
 One 14-day course of treatment



000078381
 3 0573 2450 14 8

See new warning information NDC 0573-2450-14

Treats Frequent Heartburn

1 BOTTLE INSIDE



ACTUAL SIZE

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

<p>Drug Facts</p> <p>Active ingredient (in each capsule) Purpose Esomeprazole 20 mg Acid reducer (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Uses</p> <ul style="list-style-type: none"> ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Warnings</p> <p>Allergy alert:</p> <ul style="list-style-type: none"> ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <p>Ask a doctor or pharmacist before use if you are:</p> <ul style="list-style-type: none"> ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs. <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain 	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor <p>Repeated 14-Day Courses (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. <p>Other Information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F) <p>Inactive ingredients carmellose, corn starch, FD&C blue no. 2, FD&C blue no. 2 aluminum lake, FD&C red no. 3, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polyisobutyl 80, sucrose, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600</p>
---	---



NDC 0573-2452-14

See new warning information

Treats Frequent Heartburn

May take 1 to 4 days for full effect

14 CAPSULES
One 14-day course of treatment

000078392



1 BOTTLE INSIDE



ACTUAL SIZE

Do Not Use if seal under bottle cap is imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts	Drug Facts (continued)	
<p>Active ingredient (in each capsule) Purpose Esomeprazole 20 mg Acid reducer (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days</p>	<p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect</p> <p>14-Day Course of Treatment ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor</p> <p>Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other Information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F)</p> <p>Inactive ingredients corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600</p>	<p>Tips for Managing Heartburn</p> <ul style="list-style-type: none"> • Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables. • Eat slowly and do not eat big meals. • Do not eat late at night or just before bedtime. • Do not lie flat or bend over soon after eating. • Raise the head of your bed. • Wear loose-fitting clothing around your stomach. • If you are overweight, lose weight. • If you smoke, quit smoking.

Nexium[®] 24HR
esomeprazole magnesium delayed-release capsules 20 mg/acid reducer

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
©2022 GSK group of companies or its licensor
Made in France
For most recent product information, visit www.Nexium24HR.com
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

Nexium[®] 24HR
Capsules

See new warning information
Treats Frequent Heartburn

Nexium[®] 24HR
esomeprazole magnesium delayed-release capsules 20 mg/acid reducer

28 CAPSULES

Two 14-day courses of treatment

Easy Open Cap!

2 BOTTLES INSIDE

EASY OPEN CAP!

ACTUAL SIZE

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

May take 1 to 4 days for full effect

000078385

3 0573 2450 28 5

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

<p>Drug Facts</p> <p>Active ingredient (in each capsule) Purpose Esomeprazole 20 mg Acid reducer (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days</p>	<p>Drug Facts (continued) ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions 14-Day Course of Treatment ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect 14-Day Course of Treatment ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F)</p> <p>Inactive ingredients corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600</p>
--	---



esomeprazole magnesium
delayed-release capsules
20 mg/acid reducer

Distributed by:
GSK Consumer Healthcare, Warren, NJ 07059
©2022 GSK group of companies or its licensor
Made in France
For most recent product information, visit
www.Nexium24HR.com
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

42
CAPSULES

Three 14-day courses of treatment

May take 1 to 4 days for full effect



3 0573 2450 42 1

3 BOTTLES INSIDE



ACTUAL SIZE

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

See new warning information

Treats Frequent Heartburn

Nexium[®]
esomeprazole magnesium
delayed-release capsules **24HR**
20 mg/acid reducer

May take 1 to 4 days for full effect

3 Pack Capsules

Flange of Blister
Area covered by
PRODUCT

42
CAPSULES
Three 14-day courses
of treatment

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

Nexium[®]
esomeprazole magnesium
delayed-release capsules **24HR**
20 mg/acid reducer

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

<p>Drug Facts</p> <p>Active ingredient (in each capsule) Esomeprazole 20 mg (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Purpose Acid reducer</p> <p>Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are: ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days</p>	<p>Drug Facts (continued)</p> <p>■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect</p> <p>14-Day Course of Treatment ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor</p> <p>Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other Information ■ read the directions and warnings before use ■ keep the card. It contains important information. ■ store at 20-25°C (68-77°F)</p> <p>Inactive ingredients corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-800-245-1040</p>
---	---

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
©2022 GSK group of companies or its licensor
Made in France
For most recent product information, visit www.Nexium24HR.com
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

LOT/EXP
WHITE HOLDOUT
REMOVE VARNISH

3 0573 2450 43 8

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

<p>Drug Facts</p> <p>Active ingredient (in each capsule) Purpose Esomeprazole 20 mg Acid reducer (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months</p>	<p>Drug Facts (continued) ■ you get diarrhea ■ you develop a rash or joint pain If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect</p> <p>14-Day Course of Treatment ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor</p> <p>Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other Information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F)</p> <p>Inactive ingredients carmine, corn starch, FD&C blue no. 2, FD&C blue no. 2 aluminum lake, FD&C red no. 3, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1 866-226-1600</p>
--	---

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Nexium[®] 24HR
esomeprazole magnesium delayed-release capsules (20 mg/acid reducer)

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059 ©2022 GSK group of companies or its licensor
Made in France
For most recent product information, visit www.Nexium24HR.com
Nexium is a registered trademark of AstraZeneca AB and is used under license.

42 CAPSULES

Three 14-day courses of treatment

000078397



Nexium[®] 24HR
esomeprazole magnesium delayed-release capsules (20 mg/acid reducer)

Clinically Proven to Treat Frequent Heartburn

See new warning information

Treats Frequent Heartburn

NDC 0573-2452-42

3 BOTTLES INSIDE



ACTUAL SIZE

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
09/06/2022 11:09:55 AM