

**naproxen sodium, 220 mg (NSAID)/
 diphenhydramine HCl, 25 mg
 Pain reliever/nighttime sleep-aid**

**SLEEP AID PLUS
 12 HOUR PAIN RELIEVING
 STRENGTH OF ALEVE**

20 CAPSULETS*
*CAPSULE-SHAPED TABLETS

Drug Facts
Active Ingredients
 (in each capsule)
 Diphenhydramine hydrochloride 25 mg.....Nighttime sleep-aid
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses
 • for relief of occasional sleeplessness when associated with minor aches and pains
 • helps you fall asleep and stay asleep

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 • hives • facial swelling • asthma (wheezing)
 • shock • skin redness • rash • blisters
 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • take 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • if you are pregnant or breastfeeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • **do not take more than directed** • drink a full glass of water with each dose • adults and children 12 years and over: take 2 capsules at bedtime • do not take more than 2 capsules in 24 hours • if taken with food, this product may take longer to work

Other information • **read all warnings and directions before use. Keep outer carton.** • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients: carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-955-0889 (Mon-Fri 9AM-5PM EST)

PS

Drug Facts (continued)

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • if you are pregnant or breastfeeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • **do not take more than directed** • drink a full glass of water with each dose • adults and children 12 years and over: take 2 capsules at bedtime • do not take more than 2 capsules in 24 hours • if taken with food, this product may take longer to work

Other information • **read all warnings and directions before use. Keep outer carton.** • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients: carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-955-0889 (Mon-Fri 9AM-5PM EST)

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • if you are pregnant or breastfeeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • **do not take more than directed** • drink a full glass of water with each dose • adults and children 12 years and over: take 2 capsules at bedtime • do not take more than 2 capsules in 24 hours • if taken with food, this product may take longer to work

Other information • **read all warnings and directions before use. Keep outer carton.** • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients: carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-955-0889 (Mon-Fri 9AM-5PM EST)

PS

Drug Facts (continued)

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • if you are pregnant or breastfeeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • **do not take more than directed** • drink a full glass of water with each dose • adults and children 12 years and over: take 2 capsules at bedtime • do not take more than 2 capsules in 24 hours • if taken with food, this product may take longer to work

Other information • **read all warnings and directions before use. Keep outer carton.** • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients: carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-955-0889 (Mon-Fri 9AM-5PM EST)

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • if you are pregnant or breastfeeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • **do not take more than directed** • drink a full glass of water with each dose • adults and children 12 years and over: take 2 capsules at bedtime • do not take more than 2 capsules in 24 hours • if taken with food, this product may take longer to work

Other information • **read all warnings and directions before use. Keep outer carton.** • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients: carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-955-0889 (Mon-Fri 9AM-5PM EST)

PS

Drug Facts (continued)

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • if you are pregnant or breastfeeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • **do not take more than directed** • drink a full glass of water with each dose • adults and children 12 years and over: take 2 capsules at bedtime • do not take more than 2 capsules in 24 hours • if taken with food, this product may take longer to work

Other information • **read all warnings and directions before use. Keep outer carton.** • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients: carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-955-0889 (Mon-Fri 9AM-5PM EST)

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • if you are pregnant or breastfeeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • **do not take more than directed** • drink a full glass of water with each dose • adults and children 12 years and over: take 2 capsules at bedtime • do not take more than 2 capsules in 24 hours • if taken with food, this product may take longer to work

Other information • **read all warnings and directions before use. Keep outer carton.** • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients: carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-955-0889 (Mon-Fri 9AM-5PM EST)

PS

Drug Facts (continued)

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • if you are pregnant or breastfeeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • **do not take more than directed** • drink a full glass of water with each dose • adults and children 12 years and over: take 2 capsules at bedtime • do not take more than 2 capsules in 24 hours • if taken with food, this product may take longer to work

Other information • **read all warnings and directions before use. Keep outer carton.** • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients: carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-955-0889 (Mon-Fri 9AM-5PM EST)

Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • if you are pregnant or breastfeeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • **do not take more than directed** • drink a full glass of water with each dose • adults and children 12 years and over: take 2 capsules at bedtime • do not take more than 2 capsules in 24 hours • if taken with food, this product may take longer to work

Other information • **read all warnings and directions before use. Keep outer carton.** • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients: carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-955-0889 (Mon-Fri 9AM-5PM EST)

PS

Actual Size 100%

MODIFIED DFL





(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



Actual Size 100%





Bayer, the Bayer Cross and Aleve are registered trademarks of Bayer.

Drug Facts (continued)

- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions


- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours
- if taken with food, this product may take longer to work

Other information

- read all warnings and directions before use. Keep outer carton.
- each caplet contains sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients carmellose, FD&C blue #2/aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-395-0889 (Mon - Fri 9AM - 5PM EST)



PS

Drug Facts (continued)

- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, or an one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have a trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks.

Insomnia may be a symptom of a serious underlying medical illness.

- redness or swelling is present in the painful area

Drug Facts (continued)

Active Ingredients **Purposes**
 (in each caplet)

Diphenhydramine hydrochloride, 25 mg.....Nighttime sleep-aid
 Naproxen sodium, 220 mg.....Pain reliever
 (naproxen 200 mg) (NSAID).....Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives (wheezing)
- facial swelling
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.


Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks a day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer

XXXXXXXXXX



Do not use if carbon is open or if foil seal is printed with "SAFETY SQUARE" on bottle opening is missing or broken.

Bayer HealthCare LLC
 Consumer Care
 P.O. Box 1910
 Morrisville, NJ 07962-1910 USA
MADE IN GERMANY

Bayer

LOT LOT & EXP
 EXP CODE AREA


Actual Size 100%



This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



Actual Size 100%



Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours
- if taken with food, this product may take longer to work

Other Information

- read all warnings and directions before use. **Keep outer carton.**
- each caplet contains sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive Ingredients carnauba wax, FD&C Blue #2 aluminum lake, hydroxyethylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyethylene glycol, purified water, talc, titanium dioxide

Questions or comments?
 1-800-393-0889 (Mon-Fri 9AM-5PM EST)

Bayer

Drug Facts (continued)

Directions

- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have diabetes
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When using this product

- crossiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

Bayer HealthCare LLC
 Safety Squeeze is a registered trademark of The Procter & Gamble
 P.O. Box 1910
 Morristown, NJ 07962-1910 USA
MADE IN GERMANY

LOT
 EXP

LOT & EXP CODE AREA

Drug Facts (continued)

Directions

- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have diabetes
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When using this product

- crossiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

Bayer HealthCare LLC
 Safety Squeeze is a registered trademark of The Procter & Gamble
 P.O. Box 1910
 Morristown, NJ 07962-1910 USA
MADE IN GERMANY

LOT
 EXP

LOT & EXP CODE AREA

Drug Facts

Active ingredients (in each caplet)

Diphenhydramine hydrochloride 25 mg... Nighttime sleep-aid
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) ... Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin redness
- itchiness
- facial swelling
- streak
- rash


If an allergic reaction occurs, stop use and seek medical help right away. This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:


- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep
- in children under 12 years of age


XXXXXXXXXX





**naproxen sodium, 220 mg (NSAID)/
 diphenhydramine HCl, 25 mg
 Pain reliever/nighttime sleep-aid**

**SLEEP AID PLUS
 12 HOUR PAIN RELIEVING
 STRENGTH OF ALEVE**



86 CAPSULES
 *CAPSULE-SHAPED TABLETS

NEW PRODUCT

Actual Size 100%



(b) (4)

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



Actual Size 100%