

MODIFIED DFL

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>7.0 pt	7.1 pt
Drug Facts	N/A	7 pt	7 pt
(continued):	N/A	7 pt	7 pt
Headings:	N/A	7 pt	7 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39


Actual Size 100%

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugs>



Actual Size 100%

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugs>



Actual Size 100%

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ● do not take more than directed ● drink a full glass of water with each dose ● adults and children 12 years and over: take 2 caplets at bedtime ● do not take more than 2 caplets in 24 hours ● if taken with food, this product may take longer to work

Other information ● read all warnings and directions before use. Keep outer carton. ● each caplet contains: sodium 20 mg ● store at 20-25°C (68-77°F) ● avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients carnauba wax, FD&C blue #2, aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polydione, purified water, talc, titanium dioxide

Questions or comments?
1-800-395-0689 (Mon - Fri 9AM - 5PM EST)

PS

Bayer, the Bayer Cross and Aleve are registered trademarks of Bayer.

Drug Facts (continued)

Do not use ● if you have ever had an allergic reaction to any other pain reliever/fever reducer ● unless you have time for a full night's sleep ● in children under 12 years of age ● right before or after heart surgery ● with any other product containing diphenhydramine, even one used on skin ● if you have sleeplessness without pain

Ask a doctor before use if ● stomach bleeding warning applies to you ● you have problems or serious side effects from taking pain relievers or fever reducers ● you have a history of stomach problems, such as heartburn ● you have high blood pressure, heart disease, liver disease, kidney disease, or asthma ● you are taking a diuretic ● you have a breathing problem such as emphysema or chronic bronchitis ● you have glaucoma ● you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are ● taking sedatives or tranquilizers, or any other sleep-aid ● under a doctor's care for any serious condition ● taking any other anti-inflammatories ● taking any other drug

When using this product ● drowsiness will occur ● avoid alcoholic drinks ● do not drive a motor vehicle or operate machinery ● take with food or milk if stomach upset occurs ● the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if ● you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ● pain gets worse or lasts more than 10 days ● sleeplessness persists continuously for more than 2 weeks ● insomnia may be a symptom of a serious underlying medical illness. ● redness or swelling is present in the painful area ● any new symptoms appear ● you have difficulty swallowing ● it feels like the pill is stuck in your throat

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEEZE" on bottle opening is missing or broken.

Disk by:
Bayer HealthCare LLC
Whispery, NJ 07081
MADE IN GERMANY

LOT LOT & EXP
 CODE AREA

Drug Facts

Active ingredients (in each caplet)
Diphenhydramine hydrochloride 25 mg, Nighttime sleep-aid
Naproxen sodium 220 mg (NSAID).....Pain reliever
*nonsteroidal anti-inflammatory drug

Uses
● for relief of occasional sleeplessness when associated with minor aches and pains
● helps you fall asleep and stay asleep

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
● hives ● face, arm, or leg swelling (wheezing)
● shock ● skin redness ● rash ● blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinner (anticoagulant) or steroid drug ● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● take more or for a longer time than directed

XXXXXXX
0000


MODIFIED DFL

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>7.0 pt	7.1 pt
Drug Facts	N/A	7 pt	7 pt
(continued):	N/A	7 pt	7 pt
Headings:	N/A	7 pt	7 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	≤ 39

Actual Size 100%

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugs>



Actual Size 100%

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drug>



Actual Size 100%

Reference ID: 3792020

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drug>



Actual Size 100%

Reference ID: 3792020


This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drug>



Actual Size 100%

Reference ID: 3792020

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drug>



Actual Size 100%

Reference ID: 3792020



Bayer, the Bayer Cross, and Aleve are registered trademarks of Bayer.

Drug Facts (continued)

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours
- if taken with food, this product may take longer to work

Other information

- read all warnings and directions before use. Keep outer carton.
- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients camellia wax, FD&C blue #2 aluminum lake, hydroxypropyl methylcellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
1-800-395-0689 (Mon - Fri 9AM - 5PM EST)

Bayer

Drug Facts (continued)

- right before or after heart surgery with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn, kidney disease, or asthma
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antacids
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - belly pain
 - vomiting blood
 - black stools
 - have stomach pain that does not get better
- stomach pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks

Insomnia may be a symptom of a serious underlying medical illness.

- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

PS



Drug Facts

Active ingredients (in each caplet)

Diphenhydramine hydrochloride 25 mg.....Nighttime sleep-aid
Naproxen sodium 220 mg (NSAID).....Pain reliever
Nonsteroidal anti-inflammatory drug

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin redness
- skin blistering
- facial swelling
- stomach pain
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age

XXXXXXXX



Do not use if carbon is open or if foil seal imprinted with "SAFETY SQUEEZE" on bottle opening; is missing or broken.

Dist. By:
Bayer HealthCare LLC
Whippany, NJ 07981
MADE IN GERMANY

LOT LOT & EXP
 CODE AREA
EXP

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts (continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

Actual Size 100%

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drug>



Actual Size 100%

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drug>



Actual Size 100%

Reference ID: 3792020

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

KAREN M MAHONEY
07/15/2015