


FRONT



BACK


Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	—

• if you have sleeplessness without pain Ask doctor before use if • stomach bleeding warning applies to you • you have problems or serious side effects from taking pain relievers or fever reducers • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma • you are taking a diuretic • you have a breathing problem such as emphysema or chronic bronchitis • you have glaucoma • you have trouble urinating due to an enlarged prostate gland Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers, or any other sleep-aid • under a doctor's care for any serious condition • taking any other antihistamines • taking any other drug When using this product • drowsiness will occur • avoid alcoholic drinks • do not drive a motor vehicle or operate machinery • take with food or milk if stomach upset occurs • the risk of heart attack or stroke may increase if you use more than directed or for longer than directed Stop use and ask a doctor if • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. • redness or swelling is present in the painful area • any new symptoms appear • you have difficulty swallowing • it feels like the pill is stuck in your throat If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** • do not take more than directed • drink a full glass of water with each dose • adults and children 12 years and over: take 2 caplets at bedtime • do not take more than 2 caplets in 24 hours • If taken with food, this product may take longer to work **Other information** • read all warnings and directions before use • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F) **Inactive ingredients** carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide **Questions?** 1-800-395-0689 (Mon-Fri 9-5 EST) Bayer HealthCare LLC, Morrisstown, NJ 07960 **MADE IN GERMANY**

While folded on line, tear open at dot or use scissors.
Do not use if pouch is torn or open.



Active ingredients (in each caplet)

Diphenhydramine hydrochloride 25 mg	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Nighttime sleep-aid
	*nonsteroidal anti-inflammatory drug

Use • for relief of occasional sleeplessness when associated with minor aches and pains • helps you fall asleep and stay asleep **Warnings Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • facial swelling • asthma (wheezing) • shock • skin reddening • rash • blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • unless you have time for a full night's sleep • in children under 12 years of age • right before or after heart surgery • with any other product containing diphenhydramine, even one used on skin

Actual Size 100%

(b) (4)

CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	5.5 pt
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	5.5 pt
Subheadings:	N/A	6 pt	5.5 pt
Text:	N/A	6 pt	5.5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	6 pt
Characters Per Inch:	N/A	No more than 39	36

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
02/25/2015