

product containing diphenhydramine, even one used on skin • if you have sleeplessness without pain **Ask doctor before use if** • stomach bleeding warning applies to you • you have problems or serious side effects from taking pain relievers or fever reducers • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke • you are taking a diuretic • you have a breathing problem such as emphysema or chronic bronchitis • you have glaucoma • you have trouble urinating due to an enlarged prostate gland **Ask a doctor or pharmacist before use if you are** • taking sedatives or tranquilizers, or any other sleep-aid • under a doctor's care for any serious condition • taking any other antihistamines • taking any other drug **When using his product** • drowsiness will occur • avoid alcoholic drinks • do not drive a motor vehicle or operate machinery • take with food or milk if stomach upset occurs **Stop use and ask a doctor if** • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • you have symptoms of heart problems or stroke: • chest pain • trouble breathing • weakness in one part or side of body • slurred speech • leg swelling • pain gets worse or lasts more than 10 days • sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. • redness or swelling is present in the painful area • any new symptoms appear • you have difficulty swallowing • it feels like the pill is stuck in your throat **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** • **do not take more than directed** • drink a full glass of water with each dose • adults and children 12 years and over: take 2 caplets at bedtime • do not take more than 2 caplets in 24 hours • if taken with food, his product may take longer to work **Other information** • read all warnings and directions before use • each caplet contains: sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F) **Inactive ingredients** carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide **Questions or comments?** 1-800-395-0689 (Mon - Fri 9AM - 5PM EST)
 Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981 **MADE IN GERMANY** XXXXX

While folded on line, tear open at dot or use scissors.
 Do not use if pouch is torn or open



Active ingredients (in each caplet)

Diphenhydramine hydrochloride 25 mg	Nighttime sleep-aid
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever

nonsteroidal anti-inflammatory drug




Uses • for relief of occasional sleeplessness when associated with minor aches and pains • helps you fall asleep and stay asleep **Warnings Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • facial swelling • asthma (wheezing) • shock • skin reddening • rash • blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** • if you have ever had an allergic reaction to any other pain reliever/fever reducer • unless you have time for a full night's sleep • in children under 12 years of age • right before or after heart surgery • with any other



This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



Reference ID: 4151496



Bayer
Registered Trademarks of Bayer

Drug Facts

Active ingredients (in each caplet)

Diphenhydramine hydrochloride 25 mg Nighttime sleep aid
Naproxen sodium 220 mg Pain reliever
Naproxen 200 mg (NSAID) Nonsteroidal anti-inflammatory drug

Purposes

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters.

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinner (an anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer, unless you have time for a full night's sleep, in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver or kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk
- stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better
- you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks
- insomnia may be a symptom of a serious underlying medical illness
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

Drug Facts (continued)

If pregnant or breast feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours
- taken with food, this product may take longer to work

Other information

- read all warnings and directions before use. Keep out of reach of children.
- each caplet contains: sodium 20 mg
- store at 20° to 25°C (68° to 77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients

camellia wax, FD&C blue #2 aluminum lake, hydroxypropyl methylcellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
1 800 395 0689 (Mon - Fri 9AM - 5PM EST)

Bayer

XXXXXXXXXX

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken

Dat. by:
Bayer HealthCare LLC
Whippany, NJ 07981




MADE IN GERMANY

LOT/EXP

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>






Reference ID: 4151496



This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>






Reference ID: 4151496



This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



Reference ID: 4151496



Bayer, the Bayer Cross and Alevine are registered trademarks of Bayer.

Drug Facts

Active ingredients (in each caplet)
 Diphenhydramine hydrochloride 25 mg Nighttime sleep-aid
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) Pain reliever
 *nonsteroidal anti-inflammatory drug

Purposes

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- blisters
- facial swelling
- shock
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks.

Insomnia may be a symptom of a serious underlying medical illness.

Bayer, the Bayer Cross and Alevine are registered trademarks of Bayer.

XXXXXXX

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

Dist. by:
 Bayer HealthCare LLC
 Whippany, NJ 07981

MADE IN GERMANY

LOT/EXP

Safety Squease is a registered trademark of The Procter & Gamble Company. Used with permission.

Bayer

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



Reference ID: 4151496

Drug Facts

Active ingredients (in each caplet)

Diphenhydramine hydrochloride 25 mg Nighttime sleep-aid
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* Pain reliever
 *nonsteroidal anti-inflammatory drug

Purposes

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Uses

• for relief of occasional sleeplessness when associated with minor aches and pains

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

• hives	• facial swelling
• asthma (wheezing)	• shock
• skin reddening	• rash
• blisters	

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

XXXXXXX
0000

6 80355 99852 3

Bayer, the Bayer Cross and Aleve are registered trademarks of Bayer.

XXXXXXX
0000

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours
- if taken with food, this product may take longer to work

Other information

- read all warnings and directions before use. Keep out carton.
- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-395-0689 (Mon - Fri 9AM - 5PM EST)

Bayer



TWO 80-COUNT
BOTTLES INSIDE

ALEVE[®] pm

naproxen sodium, 220 mg (NSAID)/
diphenhydramine HCl, 25 mg
Pain reliever/nighttime sleep-aid

ALEVE[®] pm

naproxen sodium, 220 mg (NSAID)/
diphenhydramine HCl, 25 mg
Pain reliever/nighttime sleep-aid

160 CAPLETS*
*CAPSULE SHAPED TABLETS

**SLEEP AID PLUS
12 HOUR PAIN RELIEVING
STRENGTH OF ALEVE**



Do not tear! Carefully open or fold seal imprinted with
scored notches on outer opening to remove or broken.

Safety Science is a registered
trademark of the Pharmacia & Upjohn
Company. Use with permission.

Dist. by:
Pharmacia Inc.
Whisper, NJ 07981

MADE IN GERMANY

Bayer, the Bayer Cross and Name
are registered trademarks of Bayer.

Bayer



XXXXX

LOT
EXP

Drug Facts

Active Ingredients (in each caplet)

Diphenhydramine hydrochloride 25 mg.....Night time sleep-aid
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....Pain reliever
*nonsteroidal anti-inflammatory drug

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- facial swelling
- hives
- asthma (wheezing)
- skin reddening
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Purposes

- relieve fever
- relieve pain
- help you sleep

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers or any other sleep-aid
- under a doctor's care for any serious condition
- taking any other antihistamines
- taking any other drug

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better

Drug Facts (continued)

- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the pill is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours
- if taken with food, this product may take longer to work

Other information

- read all warnings and directions before use. Keep out of reach of children.
- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients: carnuba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, titanium dioxide

Questions or comments? 1-800-395-0689 (Mon - Fri 9AM - 5PM EST)

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



Reference ID: 4151496



16 FREE CAPLETS WITH THIS PURCHASE



SLEEP AID PLUS 12 HOUR PAIN RELIEVING STRENGTH OF ALEVE

96 CAPLETS*
*4 CAPLETS SHAPED TABLETS

ALEVE pm
 naproxen sodium, 220 mg (NSAID) /
 diphenhydramine HCl, 25 mg
 Pain reliever/nighttime sleep-aid

Drug Facts

Active ingredients (in each caplet)
 Diphenhydramine hydrochloride 25 mg.....Nighttime sleep-aid
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID).....Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses
 • for relief of occasional sleeplessness when associated with minor aches and pains
 • helps you fall asleep and stay asleep

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 • hives • facial swelling
 • asthma (wheezing) • shock
 • skin reddening • rash
 • blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 • are age 60 or older
 • have had stomach ulcers or bleeding problems
 • take a blood thinning (anticoagulant) or steroid drug
 • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 • have 3 or more alcoholic drinks every day while using this product
 • take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

Do not use
 • if you have ever had an allergic reaction to any other pain reliever/fever reducer
 • unless you have time for a full night's sleep
 • in children under 12 years of age
 • right before or after heart surgery
 • w th any other product containing diphenhydramine, even one used on skin
 • if you have sleeplessness without pain

Ask a doctor before use if
 • stomach bleeding warning applies to you
 • you have problems or serious side effects from taking pain relievers or fever reducers
 • you have a history of stomach problems, such as heartburn
 • you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 • you are taking a diuretic
 • you have a breathing problem such as emphysema or chronic bronchitis
 • you have glaucoma
 • you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are
 • taking sedatives or tranquilizers, or any other sleep-aid
 • under a doctor's care for any serious condition
 • taking any other antihistamines
 • taking any other drug

When using this product
 • drowsiness will occur
 • avoid alcoholic drinks
 • do not drive a motor vehicle or operate machinery
 • take with food or milk if stomach upset occurs

Stop use and ask a doctor if
 • you experience any of the following signs of stomach bleeding:
 • feel faint • vomit blood • have bloody or black stools
 • have stomach pain that does not get better
 • you have symptoms of heart problems or stroke:
 • chest pain • trouble breathing
 • weakness in one part or side of body
 • slurred speech • leg swelling
 • pain gets worse or lasts more than 10 days
 • sleeplessness persists continuously for more than 2 weeks.
 Insomnia may be a symptom of a serious underlying medical illness.
 • redness or swelling is present in the painful area
 • any new symptoms appear
 • you have difficulty swallowing
 • it feels like the pill is stuck in your throat

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 • do not take more than directed
 • drink a full glass of water with each dose
 • adults and children 12 years and over: take 2 caplets at bedtime
 • do not take more than 2 caplets in 24 hours
 • if taken with food, this product may take longer to work

Other information
 • read all warnings and directions before use. Keep outer carton.
 • each caplet contains: sodium 20 mg
 • store at 20-25°C (68-77°F)
 • avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide

Questions or comments?
 1-800-395-0689 (Mon - Fri 9AM - 5PM EST)

Bayer

Do not use if carton is open or if foil seal imprinted with "SAFETY SQUEASE" on bottle opening is missing or broken.

Dist. by:
 Bayer HealthCare LLC
 Whippany, NJ 07981
MADE IN GERMANY
 LOT/EXP

Safety Squeeze is a registered trademark of The Procter & Gamble Company. Used with permission.

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



Reference ID: 4151496

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
09/12/2017