

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ZYDELIG safely and effectively. See full prescribing information for ZYDELIG.

ZYDELIG® (idelalisib) tablets, for oral use  
Initial U.S. Approval: 2014

**WARNING: FATAL AND SERIOUS TOXICITIES: HEPATIC, SEVERE DIARRHEA, COLITIS, PNEUMONITIS, INFECTIONS, and INTESTINAL PERFORATION**

*See full prescribing information for complete boxed warning.*

- Fatal and/or serious hepatotoxicity occurred in 16% to 18% of Zydelig-treated patients. Monitor hepatic function prior to and during treatment. Interrupt and then reduce or discontinue Zydelig. (5.1)
- Fatal and/or serious and severe diarrhea or colitis occurred in 14% to 20% of Zydelig-treated patients. Monitor for the development of severe diarrhea or colitis. Interrupt and then reduce or discontinue Zydelig. (5.2)
- Fatal and/or serious pneumonitis occurred in 4% of Zydelig-treated patients. Monitor for pulmonary symptoms and bilateral interstitial infiltrates. Interrupt or discontinue Zydelig. (5.3)
- Fatal and/or serious infections occurred in 21% to 48% of Zydelig-treated patients. Monitor for signs and symptoms of infection. Interrupt Zydelig if infection is suspected. (5.4)
- Fatal and serious intestinal perforation can occur in Zydelig-treated patients across clinical trials. Discontinue Zydelig if intestinal perforation is suspected. (5.5)

## RECENT MAJOR CHANGES

|                                                         |           |
|---------------------------------------------------------|-----------|
| Boxed Warning                                           | (01/2018) |
| Indications and Usage (1.2)                             | (01/2018) |
| Dosage and Administration (2.2)                         | (01/2018) |
| Warnings and Precautions (5.1, 5.2, 5.3, 5.4, 5.8, 5.9) | (01/2018) |

## INDICATIONS AND USAGE

Zydelig is a kinase inhibitor indicated for the treatment of patients with:

- Relapsed chronic lymphocytic leukemia (CLL), in combination with rituximab, in patients for whom rituximab alone would be considered appropriate therapy due to other co-morbidities. (1.1)
- Relapsed follicular B-cell non-Hodgkin lymphoma (FL) in patients who have received at least two prior systemic therapies. (1.2)
- Relapsed small lymphocytic lymphoma (SLL) in patients who have received at least two prior systemic therapies. (1.3)

### Limitation of use:

Zydelig is not indicated and is not recommended for first-line treatment of any patient. (1.1, 1.2, 1.3)

Zydelig is not indicated and is not recommended in combination with bendamustine and/or rituximab for the treatment of FL. (1.2)

Accelerated approval was granted for FL and SLL based on overall response rate. Improvement in patient survival or disease related symptoms has not been established. Continued approval for these indications may be contingent upon verification of clinical benefit in confirmatory trials.

## DOSAGE AND ADMINISTRATION

Recommended starting dose: 150 mg orally, twice daily. (2.1)

## DOSAGE FORMS AND STRENGTHS

Tablets: 150 mg, 100 mg. (3)

## CONTRAINDICATIONS

History of serious allergic reactions including anaphylaxis and toxic epidermal necrolysis. (4)

## WARNINGS AND PRECAUTIONS

- Severe cutaneous reactions: Monitor patients for the development of severe cutaneous reactions and discontinue Zydelig. (5.6)
- Anaphylaxis: Monitor patients for anaphylaxis and discontinue Zydelig. (5.7)
- Neutropenia: monitor blood counts. (5.8)
- Embryo-fetal toxicity: Zydelig may cause fetal harm. Advise women of potential risk to a fetus and to avoid pregnancy while taking Zydelig. (5.9, 8.1, 8.3)

## ADVERSE REACTIONS

The most common adverse reactions (incidence  $\geq 20\%$ ) in patients treated with Zydelig in the monotherapy trial are diarrhea, fatigue, nausea, cough, pyrexia, abdominal pain, pneumonia, and rash. (6.1)

The most common adverse reactions (incidence  $\geq 30\%$ ) in patients treated with Zydelig in combination trials are diarrhea, pneumonia, pyrexia, fatigue, rash, cough, and nausea. (6.1)

Common laboratory abnormalities include neutropenia, ALT elevations and AST elevations. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Gilead Sciences, Inc. at 1-800-GILEAD-5 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## DRUG INTERACTIONS

- Strong CYP3A Inhibitors: Additional monitoring required if alternative therapy is not available. (7.1)
- Strong CYP3A inducers: Avoid coadministration of strong CYP3A inducers. (7.1)
- CYP3A substrates: Avoid coadministration of sensitive CYP3A substrates. (7.2)

## USE IN SPECIFIC POPULATIONS

Lactation: Advise women not to breastfeed. (8.2)

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

Revised: 01/2018

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**WARNING: FATAL AND SERIOUS TOXICITIES:  
HEPATIC, SEVERE DIARRHEA, COLITIS, PNEUMONITIS,  
INFECTIONS, and INTESTINAL PERFORATION**

**1 INDICATIONS AND USAGE**

- 1.1 Relapsed Chronic Lymphocytic Leukemia
- 1.2 Relapsed Follicular B-cell non-Hodgkin Lymphoma
- 1.3 Relapsed Small Lymphocytic Lymphoma

**2 DOSAGE AND ADMINISTRATION**

- 2.1 Recommended Dosage
- 2.2 Dose Modification

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

- 5.1 Hepatotoxicity
- 5.2 Severe Diarrhea or Colitis
- 5.3 Pneumonitis
- 5.4 Infections
- 5.5 Intestinal Perforation
- 5.6 Severe Cutaneous Reactions
- 5.7 Anaphylaxis
- 5.8 Neutropenia
- 5.9 Embryo-fetal Toxicity

**6 ADVERSE REACTIONS**

- 6.1 Clinical Trial Experience
- 6.2 Postmarketing Experience

**7 DRUG INTERACTIONS**

7.1 Effects of Other Drugs on Zydelig

7.2 Effects of Zydelig on Other Drugs

**8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

**14 CLINICAL STUDIES**

- 14.1 Relapsed Chronic Lymphocytic Leukemia
- 14.2 Relapsed Follicular B-cell non-Hodgkin Lymphoma
- 14.3 Relapsed Small Lymphocytic Lymphoma

**16 HOW SUPPLIED/STORAGE AND HANDLING**

**17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

## FULL PRESCRIBING INFORMATION

### **WARNING: FATAL AND SERIOUS TOXICITIES: HEPATIC, SEVERE DIARRHEA, COLITIS, PNEUMONITIS, INFECTIONS, and INTESTINAL PERFORATION**

Fatal and/or serious hepatotoxicity occurred in 16% to 18% of Zydelig-treated patients. Monitor hepatic function prior to and during treatment. Interrupt and then reduce or discontinue Zydelig as recommended [see *Dosage and Administration (2.2)*, *Warnings and Precautions (5.1)*].

Fatal and/or serious and severe diarrhea or colitis occurred in 14% to 20% of Zydelig-treated patients. Monitor for the development of severe diarrhea or colitis. Interrupt and then reduce or discontinue Zydelig as recommended [see *Dosage and Administration (2.2)*, *Warnings and Precautions (5.2)*].

Fatal and/or serious pneumonitis occurred in 4% of Zydelig-treated patients. Monitor for pulmonary symptoms and bilateral interstitial infiltrates. Interrupt or discontinue Zydelig as recommended [see *Dosage and Administration (2.2)*, *Warnings and Precautions (5.3)*].

Fatal and/or serious infections occurred in 21% to 48% of Zydelig-treated patients. Monitor for signs and symptoms of infection. Interrupt Zydelig if infection is suspected [see *Dosage and Administration (2.2)*, *Warnings and Precautions (5.4)*].

Fatal and serious intestinal perforation can occur in Zydelig-treated patients across clinical trials. Discontinue Zydelig for intestinal perforation [see *Warnings and Precautions (5.5)*].

## 1 INDICATIONS AND USAGE

### 1.1 Relapsed Chronic Lymphocytic Leukemia

Zydelig is indicated, in combination with rituximab, for the treatment of patients with relapsed chronic lymphocytic leukemia (CLL) for whom rituximab alone would be considered appropriate therapy due to other co-morbidities.

#### Limitation of Use

Zydelig is not indicated and is not recommended for first line treatment of patients with CLL.

### 1.2 Relapsed Follicular B-cell non-Hodgkin Lymphoma

Zydelig is indicated for the treatment of patients with relapsed follicular B-cell non-Hodgkin lymphoma (FL) who have received at least two prior systemic therapies.

Accelerated approval was granted for this indication based on Overall Response Rate [see *Clinical Studies (14.2)*]. An improvement in patient survival or disease related

symptoms has not been established. Continued approval for this indication may be contingent upon verification of clinical benefit in confirmatory trials.

#### Limitation of Use

Zydelig is not indicated and is not recommended for first line treatment of patients with FL.

Zydelig is not indicated and is not recommended in combination with bendamustine and/or rituximab for the treatment of FL.

### **1.3 Relapsed Small Lymphocytic Lymphoma**

Zydelig is indicated for the treatment of patients with relapsed small lymphocytic lymphoma (SLL) who have received at least two prior systemic therapies.

Accelerated approval was granted for this indication based on Overall Response Rate [see *Clinical Studies (14.3)*]. An improvement in patient survival or disease related symptoms has not been established. Continued approval for this indication may be contingent upon verification of clinical benefit in confirmatory trials.

#### Limitation of Use

Zydelig is not indicated and is not recommended for first line treatment of patients with SLL.

## **2 DOSAGE AND ADMINISTRATION**

### **2.1 Recommended Dosage**

The recommended maximum starting dose of Zydelig is 150 mg administered orally twice daily.

Zydelig can be taken with or without food. Tablets should be swallowed whole.

Continue treatment until disease progression or unacceptable toxicity. The optimal and safe dosing regimen for patients who receive treatment longer than several months is unknown.

### **2.2 Dose Modification**

See Table 1 for dose modification instructions for specific toxicities related to Zydelig. For other severe or life-threatening toxicities related to Zydelig, withhold drug until toxicity is resolved. If resuming Zydelig after interruption for other severe or life-threatening toxicities, reduce the dose to 100 mg twice daily. Discontinue Zydelig permanently for recurrence of other severe or life-threatening Zydelig-related toxicity upon rechallenge.

**Table 1 Dose Modifications for Toxicities Due to Zydelig**

| <b>Pneumonitis</b>      | <b>Any symptomatic pneumonitis</b>                                                                                                                                                                                                                               |                                                                                                                          |                                                                                                                            |
|-------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|
|                         | Discontinue Zydelig in patients with any severity of symptomatic pneumonitis                                                                                                                                                                                     |                                                                                                                          |                                                                                                                            |
| <b>ALT/AST</b>          | <b>&gt;3-5 × ULN</b>                                                                                                                                                                                                                                             | <b>&gt;5-20 × ULN</b>                                                                                                    | <b>&gt;20 × ULN</b>                                                                                                        |
|                         | Maintain Zydelig dose. Monitor at least weekly until $\leq 1 \times$ ULN.                                                                                                                                                                                        | Withhold Zydelig. Monitor at least weekly until ALT/AST are $\leq 1 \times$ ULN, then may resume Zydelig at 100 mg BID.  | Discontinue Zydelig permanently.                                                                                           |
| <b>Bilirubin</b>        | <b>&gt;1.5-3 × ULN</b>                                                                                                                                                                                                                                           | <b>&gt;3-10 × ULN</b>                                                                                                    | <b>&gt;10 × ULN</b>                                                                                                        |
|                         | Maintain Zydelig dose. Monitor at least weekly until $\leq 1 \times$ ULN.                                                                                                                                                                                        | Withhold Zydelig. Monitor at least weekly until bilirubin is $\leq 1 \times$ ULN, then may resume Zydelig at 100 mg BID. | Discontinue Zydelig permanently.                                                                                           |
| <b>Diarrhea*</b>        | <b>Moderate diarrhea</b>                                                                                                                                                                                                                                         | <b>Severe diarrhea or hospitalization</b>                                                                                | <b>Life-threatening diarrhea</b>                                                                                           |
|                         | Maintain Zydelig dose. Monitor at least weekly until resolved.                                                                                                                                                                                                   | Withhold Zydelig. Monitor at least weekly until resolved, then may resume Zydelig at 100 mg BID.                         | Discontinue Zydelig permanently.                                                                                           |
| <b>Neutropenia</b>      | <b>ANC 1.0 to &lt;1.5 Gi/L</b>                                                                                                                                                                                                                                   | <b>ANC 0.5 to &lt;1.0 Gi/L</b>                                                                                           | <b>ANC &lt;0.5 Gi/L</b>                                                                                                    |
|                         | Maintain Zydelig dose.                                                                                                                                                                                                                                           | Maintain Zydelig dose. Monitor ANC at least weekly.                                                                      | Interrupt Zydelig. Monitor ANC at least weekly until ANC $\geq 0.5$ Gi/L, then may resume Zydelig at 100 mg BID.           |
| <b>Thrombocytopenia</b> | <b>Platelets 50 to &lt;75 Gi/L</b>                                                                                                                                                                                                                               | <b>Platelets 25 to &lt;50 Gi/L</b>                                                                                       | <b>Platelets &lt;25 Gi/L</b>                                                                                               |
|                         | Maintain Zydelig dose.                                                                                                                                                                                                                                           | Maintain Zydelig dose. Monitor platelet counts at least weekly.                                                          | Interrupt Zydelig. Monitor platelet count at least weekly. May resume Zydelig at 100 mg BID when platelets $\geq 25$ Gi/L. |
| <b>Infections</b>       | <b>Grade 3 or higher sepsis or pneumonia</b>                                                                                                                                                                                                                     |                                                                                                                          |                                                                                                                            |
|                         | Interrupt Zydelig until infection has resolved.                                                                                                                                                                                                                  |                                                                                                                          |                                                                                                                            |
|                         | <b>Evidence of CMV infection or viremia</b>                                                                                                                                                                                                                      |                                                                                                                          |                                                                                                                            |
|                         | Interrupt Zydelig in patients with evidence of active CMV infection of any grade or viremia (positive PCR or antigen test) until the viremia has resolved. If Zydelig is resumed, monitor patients by PCR or antigen test for CMV reactivation at least monthly. |                                                                                                                          |                                                                                                                            |

|                                                                                                                                                                                                                                 | Evidence of PJP infection                                                                                                                  |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                                                                                                                                                                                 | Interrupt Zydelig in patients with suspected PJP infection of any grade.<br>Permanently discontinue Zydelig if PJP infection is confirmed. |
| Abbreviations: ALT, alanine aminotransferase; AST, aspartate aminotransferase; BID, twice daily; ULN, upper limit of normal; CMV, cytomegalovirus; PCR: polymerase chain reaction; PJP: <i>Pneumocystis jirovecii</i> pneumonia |                                                                                                                                            |

\*Moderate diarrhea: increase of 4–6 stools per day over baseline; severe diarrhea: increase of  $\geq 7$  stools per day over baseline.

No dose modification is required for lymphocytosis, which has been observed in some patients taking Zydelig. This observed lymphocytosis is a pharmacodynamic effect and should not be considered progressive disease in the absence of other clinical findings.

### 3 DOSAGE FORMS AND STRENGTHS

150 mg tablets: pink, oval-shaped, film-coated tablet debossed with “GSI” on one side and “150” on the other side.

100 mg tablets: orange, oval-shaped, film-coated tablet debossed with “GSI” on one side and “100” on the other side.

### 4 CONTRAINDICATIONS

History of serious allergic reactions including anaphylaxis and toxic epidermal necrolysis [see *Warnings and Precautions* (5.6, 5.7)] .

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Hepatotoxicity

Fatal and/or serious hepatotoxicity occurred in 18% of patients treated with Zydelig monotherapy and 16% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies. Elevations in ALT or AST greater than 5 times the upper limit of normal have occurred [see *Adverse Reactions* (6.1)]. These findings were generally observed within the first 12 weeks of treatment and were reversible with dose interruption. After resumption of treatment at a lower dose, 26% of patients had recurrence of ALT and AST elevations. Discontinue Zydelig for recurrent hepatotoxicity.

Avoid concurrent use of Zydelig with other drugs that may cause liver toxicity.

Monitor ALT and AST in all patients every 2 weeks for the first 3 months of treatment, every 4 weeks for the next 3 months, then every 1 to 3 months thereafter. Monitor weekly for liver toxicity if the ALT or AST rises above 3 times the upper limit of normal until resolved. Withhold Zydelig if the ALT or AST is greater than 5 times the upper limit of normal, and continue to monitor AST, ALT and total bilirubin weekly until the abnormality is resolved [see *Dosage and Administration* (2.2)].

## 5.2 Severe Diarrhea or Colitis

Severe diarrhea or colitis (Grade 3 or higher) occurred in 14% of patients treated with Zydelig monotherapy and 20% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies [see *Adverse Reactions (6.1)*]. Diarrhea can occur at any time. Avoid concurrent use of Zydelig and other drugs that cause diarrhea. Diarrhea due to Zydelig responds poorly to antimotility agents. Median time to resolution ranged between 1 week and 1 month across trials, following interruption of Zydelig therapy and in some instances, use of corticosteroids [see *Dosage and Administration (2.2)*].

## 5.3 Pneumonitis

Fatal and serious pneumonitis occurred in patients treated with Zydelig. Clinical manifestations included interstitial infiltrates and organizing pneumonia. In randomized clinical trials of combination therapies, pneumonitis occurred in 4% of patients treated with Zydelig compared to 1% on the comparator arms. Time to onset of pneumonitis ranged from <1 to 15 months. Monitor patients on Zydelig for pulmonary symptoms. In patients taking Zydelig who present with pulmonary symptoms such as cough, dyspnea, hypoxia, interstitial infiltrates on a radiologic exam, or a decline by more than 5% in oxygen saturation, interrupt Zydelig until the etiology has been determined. If symptomatic pneumonitis or organizing pneumonia is diagnosed, initiate appropriate treatment with corticosteroids and permanently discontinue Zydelig [see *Dosage and Administration (2.2)*].

## 5.4 Infections

Fatal and/or serious infections occurred in 21% of patients treated with Zydelig monotherapy and 48% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies [see *Adverse Reactions (6.1)*]. The most common infections were pneumonia, sepsis, and febrile neutropenia. Treat infections prior to initiation of Zydelig therapy. Monitor patients on Zydelig for signs and symptoms of infection, and interrupt Zydelig for Grade 3 or higher infection [see *Dosage and Administration (2.2)*].

Serious or fatal *Pneumocystis jirovecii* pneumonia (PJP) or cytomegalovirus (CMV) occurred in <1% of patients treated with Zydelig. Provide PJP prophylaxis during treatment with Zydelig. Interrupt Zydelig in patients with suspected PJP infection of any grade, and permanently discontinue Zydelig if PJP infection of any grade is confirmed. Regular clinical and laboratory monitoring for CMV infection is recommended in patients with history of CMV infection or positive CMV serology at the start of treatment with Zydelig. Interrupt Zydelig in the setting of positive CMV PCR or antigen test until the viremia has resolved. If Zydelig is subsequently resumed, patients should be monitored by PCR or antigen test for CMV reactivation at least monthly [see *Dosage and Administration (2.2)*].



## 5.5 Intestinal Perforation

Fatal and serious intestinal perforation occurred in Zydelig-treated patients. At the time of perforation, some patients had moderate to severe diarrhea. Advise patients to promptly report any new or worsening abdominal pain, chills, fever, nausea, or vomiting. Discontinue Zydelig permanently in patients who experience intestinal perforation.

## 5.6 Severe Cutaneous Reactions

Fatal cases of Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) have occurred in patients treated with Zydelig. If SJS or TEN is suspected, interrupt Zydelig until the etiology of the reaction has been determined. If SJS or TEN is confirmed, permanently discontinue Zydelig.

Other severe or life-threatening (Grade  $\geq 3$ ) cutaneous reactions, including dermatitis exfoliative, rash, rash erythematous, rash generalized, rash macular, rash maculopapular, rash papular, rash pruritic, exfoliative rash, and skin disorder, have been reported in Zydelig-treated patients. Monitor patients for the development of severe cutaneous reactions and discontinue Zydelig.

## 5.7 Anaphylaxis

Serious allergic reactions, including anaphylaxis, have been reported in patients on Zydelig. In patients who develop serious allergic reactions, discontinue Zydelig permanently and institute appropriate supportive measures.

## 5.8 Neutropenia

Treatment-emergent Grade 3 or 4 neutropenia occurred in 25% of patients treated with Zydelig monotherapy and 58% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies. Monitor blood counts at least every 2 weeks for the first 6 months of therapy, and at least weekly in patients while neutrophil counts are less than 1.0 Gi/L [see *Dosage and Administration* (2.2)].

## 5.9 Embryo-fetal Toxicity

Based on findings in animals and its mechanism of action, Zydelig may cause fetal harm when administered to a pregnant woman. In animal reproduction studies, administration of idelalisib to pregnant rats during organogenesis caused decreased fetal weight and congenital malformations at systemic exposures 12 times those reported in patients at the recommended dose of 150 mg twice daily. Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment and for at least 1 month after the last dose. [see *Use in Specific Populations* (8.1, 8.3), *Clinical Pharmacology* (12.1), and *Nonclinical Toxicology* (13.1)].



## 6 ADVERSE REACTIONS

The following serious adverse reactions have been associated with Zydelig in clinical trials and are discussed in greater detail in other sections of the prescribing information.

- Hepatotoxicity [see *Warnings and Precautions* (5.1)]
- Severe Diarrhea or Colitis [see *Warnings and Precautions* (5.2)]
- Pneumonitis [see *Warnings and Precautions* (5.3)]
- Infections [see *Warnings and Precautions* (5.4)]
- Intestinal Perforation [see *Warnings and Precautions* (5.5)]
- Severe Cutaneous Reactions [see *Warnings and Precautions* (5.6)]
- Anaphylaxis [see *Warnings and Precautions* (5.7)]
- Neutropenia [see *Warnings and Precautions* (5.8)]

### 6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

#### **Summary of Clinical Trials in Chronic Lymphocytic Leukemia**

The safety data reflect exposure to Zydelig from two randomized, double-blind clinical trials (Studies 312-0116 and 312-0115) in 634 patients with relapsed CLL [see *Clinical Studies* (14.1)] and one randomized, open-label trial in 259 patients with relapsed CLL (Study 312-0119).

#### ***Zydelig with Rituximab (Study 312-0116; NCT01539512)***

Patients with relapsed CLL received up to 8 doses of rituximab (R) with or without Zydelig 150 mg twice daily. The median duration of exposure to Zydelig was 8 months.

Serious adverse reactions were reported in 65 (59%) patients treated with Zydelig + R. The most frequent serious adverse reactions reported for patients treated with Zydelig + R were pneumonia (23%), diarrhea (10%), pyrexia (9%), sepsis (8%), and febrile neutropenia (5%). Adverse reactions that led to discontinuation of Zydelig occurred in 19 (17%) patients. The most common adverse reactions that led to treatment discontinuations were hepatotoxicity and diarrhea/colitis.

Forty-two (38%) patients had dose interruptions and sixteen (15%) patients had dose reductions due to adverse reactions or laboratory abnormalities. The most common reasons for dose interruptions or reductions were pneumonia, diarrhea or colitis, rash, and elevated transaminases.

Table 2 and Table 3 summarize common adverse reactions and laboratory abnormalities reported for Zydelig + R and placebo + R arms.

**Table 2 Adverse Reactions Reported in ≥5% of Patients with CLL and Occurred at ≥2% Higher Incidence in Patients Receiving Zydelig in Study 312-0116**

|                                                      | Zydelig + R<br>N=110 (%) |          | Placebo + R<br>N=108 (%) |          |
|------------------------------------------------------|--------------------------|----------|--------------------------|----------|
| Adverse Reaction                                     | Any Grade                | Grade ≥3 | Any Grade                | Grade ≥3 |
| General disorders and administration site conditions |                          |          |                          |          |
| pyrexia                                              | 44 (40)                  | 3 (3)    | 20 (19)                  | 1 (1)    |
| chills                                               | 27 (25)                  | 2 (2)    | 17 (16)                  | 0        |
| pain                                                 | 8 (7)                    | 0        | 1 (1)                    | 0        |
| Gastrointestinal disorders                           |                          |          |                          |          |
| diarrhea <sup>(a)</sup>                              | 35 (32)                  | 12 (11)  | 20 (19)                  | 0        |
| nausea                                               | 30 (27)                  | 1 (1)    | 25 (23)                  | 0        |
| abdominal pain <sup>(b)</sup>                        | 20 (18)                  | 1 (1)    | 17 (16)                  | 2 (2)    |
| vomiting                                             | 17 (15)                  | 0        | 9 (8)                    | 0        |
| gastroesophageal reflux disease                      | 11 (10)                  | 1 (1)    | 0                        | 0        |
| stomatitis                                           | 7 (6)                    | 2 (2)    | 1 (1)                    | 0        |
| Respiratory, thoracic, and mediastinal disorders     |                          |          |                          |          |
| pneumonia <sup>(c)</sup>                             | 33 (30)                  | 23 (21)  | 20 (19)                  | 14 (13)  |
| Skin and subcutaneous tissue disorders               |                          |          |                          |          |
| rash <sup>(d)</sup>                                  | 27 (25)                  | 4 (4)    | 7 (6)                    | 1 (1)    |
| Metabolism and Nutrition Disorders                   |                          |          |                          |          |
| decreased appetite                                   | 18 (16)                  | 2 (2)    | 12 (11)                  | 2 (2)    |
| dehydration                                          | 7 (6)                    | 3 (3)    | 0                        | 0        |
| Infections and infestations                          |                          |          |                          |          |
| sepsis <sup>(e)</sup>                                | 10 (9)                   | 10 (9)   | 4 (4)                    | 4 (4)    |
| sinusitis                                            | 9 (8)                    | 0        | 6 (6)                    | 0        |
| urinary tract infection                              | 9 (8)                    | 1 (1)    | 4 (4)                    | 2 (2)    |
| bronchitis                                           | 8 (7)                    | 1 (1)    | 5 (5)                    | 1 (1)    |
| oral herpes                                          | 6 (5)                    | 1 (1)    | 3 (3)                    | 0        |
| Psychiatric disorders                                |                          |          |                          |          |
| insomnia                                             | 10 (9)                   | 0        | 7 (6)                    | 0        |
| Musculoskeletal and connective tissue disorders      |                          |          |                          |          |
| arthralgia                                           | 9 (8)                    | 1 (1)    | 4 (4)                    | 0        |
| Nervous system disorders                             |                          |          |                          |          |

|                  | Zydelig + R<br>N=110 (%) |          | Placebo + R<br>N=108 (%) |          |
|------------------|--------------------------|----------|--------------------------|----------|
| Adverse Reaction | Any Grade                | Grade ≥3 | Any Grade                | Grade ≥3 |
| lethargy         | 6 (5)                    | 0        | 2 (2)                    | 0        |

- (a) Diarrhea includes the following preferred terms: diarrhea, colitis.  
(b) Abdominal pain includes the following preferred terms: abdominal pain, abdominal pain upper, abdominal pain lower.  
(c) Pneumonia includes the terms: pneumonia, pneumonitis, lung infection, lung infiltration, pneumocystis jiroveci pneumonia, pneumonia legionella, lung infection pseudomonal, pneumonia fungal, respiratory tract infection, lower respiratory tract infection, and lower respiratory tract infection bacterial.  
(d) Rash includes the following preferred terms: dermatitis exfoliative, drug eruption, rash, rash erythematous, rash generalized, rash macular, rash maculo-papular, rash papular, rash pruritic, rash morbilliform, and exfoliative rash.  
(e) Sepsis includes the terms: sepsis, septic shock, neutropenic sepsis, and sepsis syndrome

**Table 3 Hematologic and Hepatic Laboratory Abnormalities Reported in ≥10% of Patients with CLL and Occurred at ≥5% Higher Incidence in Patients Receiving Zydelig in Study 312-0116**

|                               | Zydelig + R<br>N=110 (%) |           | Placebo + R<br>N=108 (%) |           |
|-------------------------------|--------------------------|-----------|--------------------------|-----------|
| Laboratory Parameter          | Any Grade                | Grade 3–4 | Any Grade                | Grade 3–4 |
| Hematology abnormalities      |                          |           |                          |           |
| neutropenia                   | 71 (65)                  | 46 (42)   | 61 (56)                  | 33 (31)   |
| leukopenia                    | 34 (31)                  | 9 (8)     | 25 (23)                  | 9 (8)     |
| lymphocytopenia               | 23 (21)                  | 11 (10)   | 13 (12)                  | 4 (4)     |
| Serum chemistry abnormalities |                          |           |                          |           |
| ALT increased                 | 43 (39)                  | 10 (9)    | 13 (12)                  | 1 (1)     |
| AST increased                 | 31 (28)                  | 6 (5)     | 16 (15)                  | 0         |

After closure of Study 312-0116, 71 patients continued treatment with Zydelig on an extension study (Study 312-0117; NCT01539291). The median duration of exposure was 18 months. Serious adverse reactions occurred in 48 (68%) patients. The most frequent serious adverse reactions reported were pneumonia (30%), diarrhea (15%), and pyrexia (11%).

The most frequent adverse reactions were pneumonia (51%), pyrexia (46%), and cough (45%). The most frequent Grade 3 or greater adverse reactions were pneumonia (30%), diarrhea (15%), and sepsis (10%).

#### ***Zydelig with Ofatumumab (Study 312-0119; NCT01659021)***

In Study 312-0119, 259 patients with relapsed CLL received up to 12 doses of ofatumumab with or without Zydelig 150 mg twice daily. The median duration of exposure to Zydelig was 13.9 months.

Serious adverse reactions were reported in 133 (77%) patients treated with Zydelig + ofatumumab. The most frequent serious adverse reactions reported were pneumonia (14%), pyrexia (13%), and diarrhea (12%).

Adverse reactions that led to discontinuation of Zydelig occurred in 71 (41%) patients. One hundred and ten (64%) patients had dose interruptions and 42 (24%) patients had dose reductions due to adverse reactions or laboratory abnormalities. The most common reasons for dose discontinuations, reductions, or interruptions were diarrhea and colitis. The most common adverse reactions were diarrhea (55%), pyrexia (38%), nausea (34%), and fatigue (34%).

### ***Zydelig with Bendamustine and Rituximab (Study 312-0115; NCT01569295)***

In Study 312-0115, patients with relapsed CLL received up to 6 cycles of bendamustine and rituximab (BR) with or without Zydelig 150 mg twice daily. The median duration of exposure to Zydelig was 18.2 months.

Serious adverse reactions were reported in 147 (71%) patients treated with Zydelig + BR. The most frequent serious adverse reactions reported for patients treated with Zydelig + BR were febrile neutropenia (21%), pneumonia (17%), pyrexia (12%), and diarrhea (6%).

Adverse reactions that led to discontinuation of Zydelig occurred in 68 (33%) patients. The most common adverse reactions that led to treatment discontinuations were pneumonia, diarrhea, and pyrexia.

One hundred twenty-two (59%) patients treated with Zydelig + BR had dose interruptions and 34 (16%) patients had dose reductions due to adverse reactions. The most common reasons for dose interruptions or reductions were increased ALT and diarrhea. The most common adverse reactions were neutropenia (64%), pyrexia (43%), and diarrhea (41%).

### **Summary of Clinical Trials in Indolent Non-Hodgkin Lymphoma**

The safety data reflect exposure to Zydelig from three open-label clinical trials (Studies 101-09 (NCT01282424), 101-02 (NCT00710528), and 101-10 (NCT01306643)) in 146 patients with indolent non-Hodgkin lymphoma (iNHL) treated with Zydelig 150 mg twice daily [see *Clinical Studies* (14.2, 14.3)]. The median duration of exposure was 6.1 months (range 0.3 to 26.4 months).

Serious adverse reactions were reported in 73 (50%) patients. The most frequent serious adverse reactions that occurred were pneumonia (15%), diarrhea (11%), and pyrexia (9%).

Adverse reactions resulted in interruption or discontinuation for 78 (53%) patients. The most common reasons for interruption or discontinuations were diarrhea (11%), pneumonia (11%), and elevated transaminases (10%).

Table 6 provides the adverse reactions occurring in at least 10% of patients receiving Zydelig monotherapy, and Table 7 provides the hematologic and hepatic laboratory abnormalities.

**Table 6 Adverse Reactions Reported in  $\geq 10\%$  of Patients with Indolent NHL Treated with Zydelig 150 mg BID**

| Adverse Reaction                                     | Zydelig Monotherapy<br>N=146 (%) |                |
|------------------------------------------------------|----------------------------------|----------------|
|                                                      | Any Grade                        | Grade $\geq 3$ |
| Gastrointestinal disorders                           |                                  |                |
| diarrhea <sup>(a)</sup>                              | 68 (47)                          | 20 (14)        |
| nausea                                               | 42 (29)                          | 2 (1)          |
| abdominal pain <sup>(b)</sup>                        | 38 (26)                          | 3 (2)          |
| vomiting                                             | 22 (15)                          | 2 (1)          |
| General disorders and administration site conditions |                                  |                |
| fatigue                                              | 44 (30)                          | 2 (1)          |
| pyrexia                                              | 41 (28)                          | 3 (2)          |
| asthenia                                             | 17 (12)                          | 3 (2)          |
| peripheral edema                                     | 15 (10)                          | 3 (2)          |
| Respiratory, thoracic, and mediastinal disorders     |                                  |                |
| cough                                                | 42 (29)                          | 1 (1)          |
| pneumonia <sup>(c)</sup>                             | 37 (25)                          | 23 (16)        |
| dyspnea                                              | 25 (17)                          | 6 (4)          |
| Skin and subcutaneous disorders                      |                                  |                |
| rash <sup>(d)</sup>                                  | 31 (21)                          | 4 (3)          |
| night sweats                                         | 18 (12)                          | 0              |
| Metabolism and nutrition disorders                   |                                  |                |
| decreased appetite                                   | 24 (16)                          | 1 (1)          |
| Infections and infestations                          |                                  |                |
| upper respiratory tract infection                    | 18 (12)                          | 0              |
| Psychiatric disorders                                |                                  |                |
| insomnia                                             | 17 (12)                          | 0              |
| Nervous system disorders                             |                                  |                |
| headache                                             | 16 (11)                          | 1 (1)          |

(a) Diarrhea includes the following preferred terms: diarrhea, colitis, enterocolitis, and gastrointestinal inflammation.

(b) Abdominal pain includes the following preferred terms: abdominal pain, abdominal pain upper, abdominal pain lower, and abdominal discomfort.

(c) Pneumonia includes the terms: pneumonia, pneumonitis, interstitial lung disease, lung infiltration, pneumonia aspiration, respiratory tract infection, atypical pneumonia, lung infection, pneumocystis jiroveci pneumonia, bronchopneumonia, pneumonia necrotizing, lower respiratory tract infection, pneumonia pneumococcal, pneumonia staphylococcal, pneumonia streptococcal, pneumonia cytomegaloviral, and respiratory syncytial virus infection.

(d) Rash includes the following preferred terms: dermatitis exfoliative, rash, rash erythematous, rash macular, rash maculo-papular, rash pruritic, and exfoliative rash

**Table 7 Hematologic and Hepatic Laboratory Abnormalities in Patients with Indolent non-Hodgkin Lymphoma Treated with Zydelig 150 mg BID**

|                               | Zydelig Monotherapy<br>N=146 (%) |         |         |
|-------------------------------|----------------------------------|---------|---------|
| Laboratory Abnormality        | Any Grade                        | Grade 3 | Grade 4 |
| Serum chemistry abnormalities |                                  |         |         |
| ALT increased                 | 73 (50)                          | 20 (14) | 7 (5)   |
| AST increased                 | 60 (41)                          | 12 (8)  | 6 (4)   |
| Hematology abnormalities      |                                  |         |         |
| neutrophils decreased         | 78 (53)                          | 20 (14) | 16 (11) |
| hemoglobin decreased          | 41 (28)                          | 3 (2)   | 0       |
| platelets decreased           | 38 (26)                          | 4 (3)   | 5 (3)   |

Grades were obtained per CTCAE version 4.03.

### **Summary of Discontinued Clinical Trials in First-Line CLL and Early Line iNHL**

Safety data described below reflect exposure to Zydelig in three randomized, double-blind clinical trials (Studies 312-0123, 313-0124, and 313-0125) in patients with CLL and iNHL.

In Study 312-0123 (NCT01980888), 311 patients with previously untreated CLL received up to 6 cycles of BR with or without Zydelig 150 mg twice daily.

In Study 313-0124 (NCT01732913), 295 patients with previously treated iNHL received 8 doses of R with or without Zydelig 150 mg twice daily. Patients had a median of one prior therapy.

In Study 313-0125 (NCT01732926), 475 patients with previously treated iNHL received up to 6 cycles of BR with or without Zydelig 150 mg twice daily. Patients had a median of two prior therapies.

These three studies were terminated early due to a higher incidence of fatal and/or serious adverse reactions observed in patients treated with Zydelig in combination with R or BR. The most frequent serious adverse reactions were in the system organ classes of infections and infestations, blood and lymphatic system disorders, and gastrointestinal disorders.

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during post-approval use of Zydelig. Because postmarketing reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Skin and Subcutaneous Disorders - Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN)

## 7 DRUG INTERACTIONS

### 7.1 Effects of Other Drugs on Zydelig

**Table 8 Drug Interactions with Zydelig that affect Idelalisib Concentrations**

|                                |
|--------------------------------|
| <b>Strong CYP3A Inhibitors</b> |
| Clinical Impact                |
| Prevention or Management       |
| <b>Strong CYP3A Inducers</b>   |
| Clinical Impact                |
| Prevention or Management       |

### 7.2 Effects of Zydelig on Other Drugs

The coadministration of Zydelig with a CYP3A substrate may increase the concentrations of this CYP3A substrate. Avoid coadministration of Zydelig with sensitive CYP3A substrates [see *Clinical Pharmacology (12.3)*].

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### *Risk Summary*

Based on findings in animal studies (see *Data*) and the mechanism of action [see *Clinical Pharmacology (12.1)*], Zydelig may cause fetal harm when administered to a pregnant woman.

There are no available data in pregnant women to inform the drug-associated risk. In animal reproduction studies, administration of idelalisib to pregnant rats during organogenesis resulted in decreased fetal weight and congenital malformations in rats at maternal exposures (AUC) 12 times those reported in patients at the recommended dose of 150 mg twice daily (see *Data*).

All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. The background risk of major birth defects and miscarriage for the indicated population is unknown. However, the background risk of major birth defects is 2-4% and of miscarriage is 15-20% of clinically recognized pregnancies in the U.S. general population.



## *Data*

### *Animal Data*

In an embryo-fetal development study in rats, pregnant animals receiving oral doses of idelalisib during the period of organogenesis (implantation to closure of the hard palate), embryo-fetal toxicities were observed at the mid- and high-doses that also resulted in maternal toxicity, based on reductions in maternal body weight gain. Adverse findings at idelalisib doses  $\geq 75$  mg/kg/day included decreased fetal weights, external malformations (short tail), and skeletal variations (delayed ossification and/or unossification of the skull, vertebrae, and sternebrae). Additional findings were observed at 150 mg/kg/day dose of idelalisib and included urogenital blood loss, complete resorption, increased post-implantation loss, and malformations (vertebral agenesis with anury, hydrocephaly, and microphthalmia/anophthalmia). The dose of 75 and 150 mg/kg/day of idelalisib in rats resulted in exposures (AUC) of approximately 12 and 30 times, respectively, the human exposure at the recommended dose of 150 mg twice daily.

## **8.2 Lactation**

### **Risk Summary**

No data are available regarding the presence of idelalisib or its metabolites in human milk or its effects on the breastfed child or on milk production. Because of the potential for serious adverse reactions from Zydelig in a breastfed child, advise lactating women not to breastfeed while taking Zydelig and for at least 1 month after the last dose.

## **8.3 Females and Males of Reproductive Potential**

### *Pregnancy*

Based on animal studies, Zydelig may cause fetal harm when administered to a pregnant woman [see *Use in Specific Populations* (8.1)]. Females of reproductive potential should have a pregnancy test prior to starting treatment with Zydelig.

### *Contraception*

#### **Females**

Based on animal studies, Zydelig can cause fetal harm when administered to a pregnant woman [see *Use in Specific Populations* (8.1)]. Advise females of reproductive potential to use effective contraception during treatment with Zydelig and for at least 1 month after the last dose.

#### **Males**

Based on findings in animal reproduction studies, advise male patients with female partners of reproductive potential to use effective contraception during treatment and for 3 months after the last dose of Zydelig [see *Use in Specific Populations* (8.1)].

## 8.4 Pediatric Use

Safety and effectiveness of Zydelig in children less than 18 years of age have not been established.

## 8.5 Geriatric Use

In clinical trials of Zydelig in 615 patients with FL, SLL, and CLL, 327 (53%) patients were age 65 and older. No major differences in effectiveness were observed. When comparing patients 65 years of age or older to younger patients with indolent non-Hodgkin lymphoma, older patients had a higher incidence of discontinuation due to an adverse reaction (28% vs 20%), higher incidence of serious adverse reactions (64% vs 37%), and higher incidence of death (11% vs 5%). When comparing patients 65 years of age or older to younger patients with CLL, older patients had a higher incidence of discontinuation due to an adverse reaction (36% vs 28%), higher incidence of serious adverse reactions (73% vs 67%), and higher incidence of death (13% vs 9%).

## 8.6 Hepatic Impairment

Dose adjustment is not recommended for patients with ALT or AST or bilirubin > upper limit of normal (ULN); however, limited safety and efficacy data are available for patients with baseline AST or ALT > 2.5 x ULN or bilirubin > 1.5 x ULN. Monitor patients with baseline hepatic impairment for signs of Zydelig toxicity [see *Warnings and Precautions* (5)]. Follow dose modifications for adverse reactions [see *Dosage and Administration* (2.2)].

## 11 DESCRIPTION

Idelalisib is an inhibitor of phosphatidylinositol 3-kinase, PI3K $\delta$ .

The chemical name for idelalisib is 5-fluoro-3-phenyl-2-[(1S)-1-(9H-purin-6-ylamino)propyl]quinazolin-4(3H)-one. It has a molecular formula of C<sub>22</sub>H<sub>18</sub>FN<sub>7</sub>O and a molecular weight of 415.42 g/mol. Idelalisib has the following structural formula:



Idelalisib is a white to off-white solid with a pH-dependent aqueous solubility ranging from <0.1 mg/mL at pH 5-7 to over 1 mg/mL at pH 2 under ambient conditions.

Zydelig (idelalisib) tablets are for oral administration. Each tablet contains either 100 mg or 150 mg of idelalisib with the following inactive ingredients: microcrystalline cellulose, hydroxypropyl cellulose, croscarmellose sodium, sodium starch glycolate, magnesium stearate and a tablet coating. The tablet coating consists of polyethylene glycol, talc, polyvinyl alcohol, and titanium dioxide and of FD&C Yellow #6/Sunset Yellow FCF Aluminum Lake (for the 100 mg tablet) and red iron oxide (for the 150 mg tablet).

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Idelalisib is an inhibitor of PI3K $\delta$  kinase, which is expressed in normal and malignant B-cells. Idelalisib induced apoptosis and inhibited proliferation in cell lines derived from malignant B-cells and in primary tumor cells. Idelalisib inhibits several cell signaling pathways, including B-cell receptor (BCR) signaling and the CXCR4 and CXCR5 signaling, which are involved in trafficking and homing of B-cells to the lymph nodes and bone marrow. Treatment of lymphoma cells with idelalisib resulted in inhibition of chemotaxis and adhesion, and reduced cell viability.

### **12.2 Pharmacodynamics**

#### **Cardiac Electrophysiology**

At a dose 2.7 times the maximum recommended dose, Zydelig did not prolong the QT/QTc interval (i.e.,  $\leq 10$  ms).

### **12.3 Pharmacokinetics**

Idelalisib exposure increased in a less than dose-proportional manner over a dose range of 50 mg to 350 mg twice daily (0.3 to 2.3 times the approved recommended dosage).

Following 150 mg twice daily administration of idelalisib, average (% coefficient of variation) maximum concentrations ( $C_{\max}$ ) and area under the curve (AUC) at steady-state were 1861 (43%) ng/mL and 10598 (41%) ng•h/mL for idelalisib.

#### **Absorption**

The median time to peak concentration ( $T_{\max}$ ) was observed at 1.5 hours.

#### *Food Effect*

The administration of a single dose of Zydelig with a high-fat meal (900 calories: 525 calories fat, 250 calories carbohydrates, and 125 calories protein) increased idelalisib AUC 1.4-fold relative to fasting conditions. Zydelig can be administered without regard to food.

#### **Distribution**

Protein binding of idelalisib is  $\geq 84\%$  with no concentration dependence.

The mean blood-to-plasma ratio was 0.7.

The apparent central volume of distribution at steady state is 23 L (%CV ~85%).

Idelalisib is a substrate of P-glycoprotein (P-gp) and BCRP in vitro.

### **Elimination**

The population apparent systemic clearance at steady-state is 14.9 L/hr (%CV ~ 38%).

The population terminal elimination half-life of idelalisib is 8.2 hours.

### **Metabolism**

Idelalisib is metabolized via aldehyde oxidase and CYP3A with additional minor metabolism by UGT1A4.

### **Excretion**

Following a single 150 mg dose of radiolabeled idelalisib, 78% of the radioactivity was excreted in feces and 14% was excreted in the urine. Idelalisib is not a substrate of OATP1B1, OATP1B3, OAT1, OAT3, or OCT2.

### **Specific Populations**

Age (18 to 91 years), sex, race (White, and non-Whites), renal impairment (creatinine clearance  $\geq 15$  mL/min) and weight (38 to 148 kg) had no effect on idelalisib exposure.

### **Pediatric Patients**

The pharmacokinetics of idelalisib in pediatric patients is unknown.

### **Patients with Hepatic Impairment**

The mean AUC increased up to 1.7-fold in patients with hepatic impairment (defined as ALT or AST or bilirubin values  $\geq$  ULN) compared to patients with normal hepatic function. There is limited information on idelalisib exposure in patients with baseline AST or ALT  $> 2.5 \times$  ULN or bilirubin  $> 1.5 \times$  ULN [see *Specific Populations (8.6)*].

### **Drug Interaction Studies**

#### ***Effect of Other Drugs on Idelalisib***

The coadministration of rifampin (strong CYP3A inducer and P-gp inducer) to healthy subjects decreased the mean idelalisib AUC by 75% and the mean  $C_{\max}$  by 58% [see *Drug Interactions (7.1)*].

The coadministration of ketoconazole (strong CYP3A inhibitor and P-gp inhibitor) to healthy subjects increased the mean idelalisib AUC by 1.8-fold. No changes in the mean  $C_{\max}$  were observed [see *Drug Interactions (7.1)*].

In vitro studies suggest that idelalisib inhibits CYP2C8, CYP2C19, and UGT1A1 and induces CYP2B6.

### ***Effect of Idelalisib on Other Drugs***

The mean  $C_{\max}$  of midazolam increased by 2.4-fold and the mean AUC of midazolam increased by 5.4-fold when midazolam (sensitive CYP3A substrate) was coadministered with Zydelig [see *Drug Interactions* (7.2)].

No changes in exposure to rosuvastatin (OATP1B1 and OATP1B3 substrate) or digoxin (P-glycoprotein substrate) were observed when coadministered with Zydelig.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Idelalisib was not carcinogenic in a 26-week study in transgenic mice when administered daily by oral gavage at doses up to 500 mg/kg/day in males and 1000 mg/kg/day in females. Idelalisib was not carcinogenic in a 2-year study in rats when administered daily by oral gavage at exposures 0.40/0.62-fold (male/female), compared to the exposure in patients with hematologic malignancies administered the recommended dose of 150 mg twice daily.

Idelalisib did not induce mutations in the bacterial mutagenesis (Ames) assay and was not clastogenic in the in vitro chromosome aberration assay using human peripheral blood lymphocytes. Idelalisib was genotoxic in males in the in vivo rat micronucleus study at a high dose of 2000 mg/kg.

Idelalisib may impair fertility in humans. In a fertility study, treated male rats (25, 50, or 100 mg/kg/day of idelalisib) were mated with untreated females. Decreased epididymidal and testicular weights were observed at all dose levels and reduced sperm concentration at the mid- and high doses; however, there were no adverse effects on fertility parameters. The low dose in males resulted in an exposure (AUC) that is approximately 50% of the exposure in patients at the recommended dose of 150 mg twice daily.

In a separate fertility study, treated female rats (25, 50, or 100 mg/kg/day of idelalisib) were mated with untreated males. There were no adverse effects on fertility parameters; however, there was a decrease in the number of live embryos at the high dose. The high dose in females resulted in an exposure (AUC) that is approximately 17-fold the exposure in patients at the recommended dose of 150 mg twice daily.

### **13.2 Animal Toxicology and/or Pharmacology**

Toxicities observed in animals and not reported in patients include cardiac toxicity (cardiomyopathy, inflammation, and increased heart weight) and pancreatic findings (inflammation, hemorrhage, and low-incidence acinar degeneration and hyperplasia). These findings were observed in Sprague-Dawley rats in toxicology studies at exposures (AUCs) higher than those reported in patients at the recommended dose of 150 mg twice daily. Cardiac inflammation was mainly seen in a 28-day study in rats, the other findings were observed in the 13-week and/or 6-month studies.

## 14 CLINICAL STUDIES

### 14.1 Relapsed Chronic Lymphocytic Leukemia

#### Study 312-0116

Zydelig was evaluated in a randomized, double-blind, placebo-controlled study GS-US-312-0116 (referred to as 312-0116) (NCT01539512) in 220 patients with relapsed CLL who required treatment and were unable to tolerate standard chemoimmunotherapy due to coexisting medical conditions, reduced renal function as measured by creatinine clearance  $< 60$  mL/min, or NCI CTCAE Grade  $\geq 3$  neutropenia or Grade  $\geq 3$  thrombocytopenia resulting from myelotoxic effects of prior therapy with cytotoxic agents. Patients were randomized 1:1 to receive 8 doses of rituximab (first dose at  $375$  mg/m<sup>2</sup>, subsequent doses at  $500$  mg/m<sup>2</sup> every 2 weeks for 4 infusions and every 4 weeks for an additional 4 infusions) in combination with either an oral placebo twice daily or with Zydelig  $150$  mg taken twice daily until disease progression or unacceptable toxicity.

In Study 312-0116, the median age was 71 years (range 47, 92) with 78% over 65, 66% were male, and 90% were Caucasian. The median time since diagnosis was 8.5 years. The median number of prior therapies was 3. Nearly all (96%) patients had received prior anti-CD20 monoclonal antibodies. The most common ( $>15\%$ ) prior regimens were: bendamustine + rituximab (BR) (98 patients, 45%), fludarabine + cyclophosphamide + rituximab (75 patients, 34%), single-agent rituximab (67 patients, 31%), fludarabine + rituximab (37 patients, 17%), and chlorambucil (36 patients, 16%). The median CIRS (Cumulative Illness Rating Scale) score was 8 (range 0-17), and 85% of patients had a score of  $>6$ . Median Karnofsky score was 80. Median estimated Creatinine Clearance (eCrCl) was  $63.6$  mL/min, with 41% of patients having an eCrCl of  $<60$  mL/min. At screening, 19.5% of patients had a platelet count of  $<50 \times 10^9$ /L, and 13.2% had an absolute neutrophil count (ANC) of  $<1 \times 10^9$ /L.

The efficacy of Zydelig was evaluated by progression free survival (PFS), as assessed by an independent review committee (IRC). The trial was stopped for efficacy following the first pre-specified interim analysis. Results of a second interim analysis continued to show a statistically significant improvement for Zydelig + R over placebo + R for the primary endpoint of PFS (HR: 0.18, 95% CI [0.10, 0.32],  $p < 0.0001$ ).

At the final analysis, with a median follow-up of 8.3 months for the Zydelig + R group, and 5.6 months for the placebo + R group, the median PFS for the Zydelig + R group was 19.4 months (95% CI: 12.3, Not Reached) versus 6.5 months (95% CI: 4.0, 7.3) for the placebo + R group (HR: 0.15, 95% CI [0.09, 0.24],  $p < 0.0001$ ).

Updated efficacy results are shown in Table 9, and the Kaplan-Meier curve for PFS is shown in Figure 1.

**Table 9 Efficacy Results from Study 312-0116**

|            |                          | Zydelig + R<br>N=110 | Placebo + R<br>N=110 |
|------------|--------------------------|----------------------|----------------------|
| <b>PFS</b> | Median (months) (95% CI) | 19.4 (12.3, NR)      | 6.5 (4.0, 7.3)       |


|             |                          |                       |                |
|-------------|--------------------------|-----------------------|----------------|
|             | Hazard ratio (95% CI)    | 0.15 (0.09, 0.24)     |                |
|             | P-value                  | < 0.0001 <sup>†</sup> |                |
| <b>ORR*</b> | (All PRs)                | 92 (83.6%)            | 17 (15.5%)     |
|             | 95% CI                   | 75.4, 90.0            | 9.3, 23.6      |
|             | Odds Ratio (95% CI)      | 27.8 (13.4, 57.5)     |                |
|             | P-value                  | <0.0001               |                |
| <b>DOR</b>  | Median (months) (95% CI) | NR (12, NR)           | 6.2 (2.8, 6.5) |

PFS: progression-free survival; NR: not reached; ORR: overall response rate; PR: partial response; DOR: duration of response

<sup>†</sup> The p value for PFS was based on stratified log-rank test.

\* ORR defined as the proportion of patients who achieved a complete response (CR) or PR. All PRs achieved; none of the patients achieved a CR.

**Figure 1 Kaplan-Meier Plot of IRC-Assessed PFS for Study 312-0116**



## 14.2 Relapsed Follicular B-cell non-Hodgkin Lymphoma

### Study 101-09

The safety and efficacy of Zydelig in patients with FL was evaluated in a single-arm, multicenter study 101-09 (NCT01282424) which included 72 patients with follicular B-cell non-Hodgkin lymphoma who had relapsed within 6 months following rituximab and an alkylating agent and had received at least 2 prior treatments. The median age was 62 years (range 33 to 84), 54% were male, and 90% were Caucasian. At enrollment, 92% of patients had a baseline ECOG performance status of 0 or 1. The median time since diagnosis was 4.7 years and the median number of prior treatments was 4 (range



2 to 12). The most common prior combination regimens were R-CHOP (49%), BR (50%), and R-CVP (28%). At baseline, 33% of patients had extranodal involvement and 26% had bone marrow involvement.

Patients received 150 mg of Zydelig orally twice daily until evidence of disease progression or unacceptable toxicity. Tumor response was assessed according to the revised International Working Group response criteria for malignant lymphoma. The primary endpoint was Independent Review Committee-assessed overall response rate (ORR). Efficacy results are summarized in Table 11.

**Table 11 Overall Response Rate (ORR) and Duration of Response (DOR) in Patients with Relapsed Follicular Lymphoma**

|                             | N=72                               |
|-----------------------------|------------------------------------|
| ORR                         | 39 (54%)                           |
| 95% CI                      | (42, 66%)                          |
| CR                          | 6 (8%)                             |
| PR                          | 33 (46%)                           |
| Median* DOR, months (range) | median not evaluable (0.0+, 14.8+) |

CI = confidence interval; CR = complete response; PR = partial response

\* Kaplan-Meier estimate

The median time to response was 1.9 months (range 1.6–8.3).

### 14.3 Relapsed Small Lymphocytic Lymphoma

#### **Study 101-09**

The safety and efficacy of Zydelig in patients with SLL was evaluated in a single-arm, multicenter study 101-09 (NCT01282424) which included 26 patients with small lymphocytic lymphoma who had relapsed within 6 months following rituximab and an alkylating agent and had received at least 2 prior treatments. The median age was 65 years (range 50 to 87), 73% were male, and 81% were Caucasian. At enrollment, 96% of patients had a baseline ECOG performance status of 0 or 1. The median time since diagnosis was 6.7 years and the median number of prior treatments was 4 (range 2 to 9). The most common prior combination regimens were BR (81%), FCR (62%) and R-CHOP (35%). At baseline, 27% of patients had extranodal involvement.

Patients received 150 mg of Zydelig orally twice daily until evidence of disease progression or unacceptable toxicity. Tumor response was assessed according to the revised International Working Group response criteria for malignant lymphoma. The primary endpoint was Independent Review Committee-assessed overall response rate (ORR). Efficacy results are summarized in Table 12.

**Table 12 Overall Response Rate (ORR) and Duration of Response (DOR) in Patients with Relapsed Small Lymphocytic Lymphoma**

|                             | N=26                  |
|-----------------------------|-----------------------|
| ORR<br>95% CI               | 15 (58%)<br>(37, 77%) |
| CR<br>PR                    | 0<br>15 (58%)         |
| Median* DOR, months (range) | 11.9 (0.0+, 14.7+)    |

CI = confidence interval; CR = complete response; PR = partial response

\* Kaplan-Meier estimate

The median time to response was 1.9 months (range 1.6–8.3).

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Zydelig tablets supplied as follows:

| Tablet Strength | Package Configuration                                                                                                                                    | NDC No.      | Description of Tablet; Debossed on Tablet                          |
|-----------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|--------------|--------------------------------------------------------------------|
| 150 mg          | High density polyethylene (HDPE) bottle with a polyester fiber coil, capped with a child-resistant closure. Each bottle contains 60 film-coated tablets. | 61958-1702-1 | Oval shaped; pink; "150" on one side and "GSI" on the other side   |
| 100 mg          |                                                                                                                                                          | 61958-1701-1 | Oval-shaped; orange; "100" on one side and "GSI" on the other side |

Store between 20–30 °C (68–86 °F) with excursions permitted 15–30 °C (59–86 °F).

- Dispense only in original container.
- Do not use if seal over bottle opening is broken or missing.

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Physicians and health care professionals are advised to discuss the following with patients prior to treatment with Zydelig:

- **Hepatotoxicity**

Advise patients that Zydelig can cause significant elevations in liver enzymes, and that serial testing of serum liver tests (ALT, AST, and bilirubin) are recommended while

taking Zydelig [see *Warnings and Precautions* (5.1)]. Advise patients to report symptoms of liver dysfunction including jaundice, bruising, abdominal pain, or bleeding.

- **Severe Diarrhea or Colitis**

Advise patients that Zydelig may cause severe diarrhea or colitis and to notify their healthcare provider immediately if the number of bowel movements in a day increases by six or more [see *Warnings and Precautions* (5.2)].

- **Pneumonitis**

Advise patients of the possibility of pneumonitis, and to report any new or worsening respiratory symptoms including cough or dyspnea [see *Warnings and Precautions* (5.3)].

- **Infections**

Advise patients that Zydelig can cause serious infections that may be fatal. Advise patients to immediately report symptoms of infection (e.g. pyrexia) [see *Warnings and Precautions* (5.4)].

- **Intestinal Perforation**

Advise patients of the possibility for intestinal perforation and to notify their healthcare provider immediately if they experience severe abdominal pain [see *Warnings and Precautions* (5.5)].

- **Severe Cutaneous Reactions**

Advise patients that Zydelig may cause severe cutaneous reactions and to notify their healthcare provider immediately if they develop a severe skin reaction [see *Warnings and Precautions* (5.6)].

- **Anaphylaxis**

Advise patients that anaphylaxis can occur during treatment with Zydelig and to notify their healthcare provider immediately if they experience symptoms of anaphylaxis [see *Warnings and Precautions* (5.7)].

- **Neutropenia**

Advise patients of the need for periodic monitoring of blood counts. Advise patients to notify their healthcare provider immediately if they develop a fever or any signs of infection [see *Warnings and Precautions* (5.8)].

- **Embryo-Fetal Toxicity**

Advise females to inform their healthcare provider if they are pregnant or become pregnant. Inform female patients of the risk to a fetus and potential loss of the pregnancy [see *Use in Specific Populations* (8.1)].

Advise females of reproductive potential to use effective contraception during treatment and for 1 month after receiving the last dose of Zydelig [see *Warnings and Precautions* (5.9) and *Use in Specific Populations* (8.1, 8.3)].

Advise lactating women not to breastfeed during treatment with Zydelig and for at least 1 month after the last dose [see *Use in Specific Populations* (8.2)].

- **Instructions for Taking Zydelig**

Advise patients to take Zydelig exactly as prescribed and not to change their dose or to stop taking Zydelig unless they are told to do so by their healthcare provider. Zydelig may be taken with or without food. Zydelig tablets should be swallowed whole. Advise patients that if a dose of Zydelig is missed by less than 6 hours, to take the missed dose right away and take the next dose as usual. If a dose of Zydelig is missed by more than 6 hours, advise patients to wait and take the next dose at the usual time.

Manufactured and distributed by:

Gilead Sciences, Inc.

Foster City, CA 94404

GSI and Zydelig are trademarks or registered trademarks of Gilead Sciences, Inc., or its related companies. All other trademarks referenced herein are the property of their respective owners.

© 2018 Gilead Sciences, Inc. All rights reserved.

205858-GS-00X-PI

**MEDICATION GUIDE**  
**ZYDELIG® (zye-DEL-ig)**  
**(idelalisib)**  
**tablets**

**What is the most important information I should know about Zydelig?**

**Zydelig can cause serious side effects that can lead to death, including:**

- **Liver problems.** Abnormal liver blood test results are common during treatment with Zydelig. Zydelig can cause severe liver problems. Your doctor will do blood tests before and during your treatment with Zydelig to check for liver problems. Tell your doctor right away if you get any of the following symptoms of liver problems:
  - o yellowing of your skin or the white part of your eyes (jaundice)
  - o dark or brown (tea colored) urine
  - o pain in the upper right side of your stomach area (abdomen)
  - o bleeding or bruising more easily than normal
- **Severe diarrhea.** Diarrhea is common during treatment with Zydelig and can sometimes be severe. Tell your doctor right away if the number of bowel movements you have in a day increases by six or more. Ask your doctor about medicines you can take to treat your diarrhea.
- **Lung problems (pneumonitis).** Your doctor may do tests to check your lungs if you have breathing problems during treatment with Zydelig. Tell your doctor right away if you get new or worsening cough, shortness of breath, difficulty breathing, or wheezing. Your doctor may treat you with a corticosteroid medicine if you develop lung problems.
- **Infections.** Zydelig can cause serious infections that may lead to death. Tell your doctor right away if you have a fever or any signs of an infection during treatment with Zydelig.
- **Tear in intestinal wall (perforation).** Tell your doctor or get medical help right away if you get new or worsening stomach area (abdomen) pain, chills, fever, nausea, or vomiting.
- **Severe skin reactions.** Tell your doctor right away if you get any of the following symptoms during treatment with Zydelig:
  - o painful sores or ulcers on your skin, lips, or in your mouth
  - o severe rash with blisters or peeling skin
  - o rash with itching

If you have any of the above serious side effects during treatment with Zydelig, your doctor may completely stop your treatment, stop your treatment for a period of time, or change your dose of Zydelig.

See “**What are the possible side effects of Zydelig?**” for more information about side effects.

**What is Zydelig?**

Zydelig is a prescription medicine used to treat people with:

- **Chronic Lymphocytic Leukemia (CLL)** in combination with rituximab when CLL comes back after prior cancer treatment and when rituximab treatment alone may be used due to other health problems.
- **Follicular B-cell non-Hodgkin Lymphoma (FL)** when the disease has come back after treatment with at least two prior medicines.
- **Small Lymphocytic Lymphoma (SLL)** when the disease comes back after treatment with at least two prior medicines.

Zydelig should not be used as the first medicine to treat people with CLL, FL, or SLL.

Zydelig should not be used in combination with bendamustine and/or rituximab to treat people with FL.

It is not known if Zydelig is safe and effective in children less than 18 years of age.

**Do not take Zydelig** if you have a history of serious allergic reactions including a severe skin reaction called toxic epidermal necrolysis (TEN).

**Before taking Zydelig, tell your doctor about all of your medical conditions, including if you:**

- have liver problems
- have lung problems
- have an infection
- are pregnant or plan to become pregnant. Zydelig may harm your unborn baby. Females who are able to become pregnant should have a pregnancy test before starting treatment with Zydelig.
  - o **Females** who are able to become pregnant should use effective birth control (contraception) during treatment with Zydelig and for at least 1 month after the last dose of Zydelig. Talk to your doctor about birth control methods that may be right for you. Tell your doctor right away if you become pregnant or think you are pregnant during treatment with Zydelig.
  - o **Males** with female partners who are able to become pregnant should use effective birth control (contraception) during treatment with Zydelig and for 3 months after the last dose.
- are breastfeeding or plan to breastfeed. It is not known if Zydelig passes into your breast milk. Do not breastfeed during your treatment with Zydelig and for at least 1 month after the last dose.

**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Zydelig and certain other medicines may affect each other. Know the medicines you take. Keep a list of your medicines and show it to your doctor and pharmacist when you get a new medicine.

**How should I take Zydelig?**

- Take Zydelig exactly as your doctor tells you to take it.
- Your doctor may change your dose of Zydelig or tell you to stop taking Zydelig. Do not change your dose or stop taking Zydelig without first talking to your doctor.
- Take Zydelig 2 times a day.
- You may take Zydelig with or without food.
- Take Zydelig tablets whole.
- Do not miss a dose of Zydelig. If you miss a dose of Zydelig by less than 6 hours, take the missed dose right away. Then take your next dose as usual. If you miss a dose of Zydelig by more than 6 hours, wait and take the next dose of Zydelig at your usual time.

**What are the possible side effects of Zydelig?**

Zydelig can cause serious side effects, including:

- See **“What is the most important information I should know about Zydelig?”**
- **Anaphylaxis.** Tell your doctor or get medical help right away if you have a serious allergic reaction during treatment with Zydelig.
- **Low white blood cell count (neutropenia).** Neutropenia is common during treatment with Zydelig and can sometimes be severe. Your doctor will check your blood counts regularly during treatment with Zydelig. Tell your doctor right away if you have a fever or any signs of an infection during treatment with Zydelig.

**The most common side effects of Zydelig when used alone include:**

- |             |                               |
|-------------|-------------------------------|
| • tiredness | • stomach area (abdomen) pain |
| • nausea    | • pneumonia                   |
| • cough     | • rash                        |
| • fever     |                               |

**The most common side effects of Zydelig when used in combination with rituximab or with bendamustine and rituximab include:**

- |             |          |
|-------------|----------|
| • pneumonia | • rash   |
| • fever     | • cough  |
| • tiredness | • nausea |

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of Zydelig. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store Zydelig?**

- Store Zydelig between 68°F to 86°F (20°C to 30°C).
- Keep Zydelig in its original container.
- Do not use Zydelig if the seal over the bottle opening is broken or missing.

**Keep Zydelig and all medicines out of reach of children.**

**General information about the safe and effective use of Zydelig.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Zydelig for a condition for which it was not prescribed. Do not give Zydelig to other people, even if they have the same symptoms you have. It may harm them. You can ask your doctor or pharmacist for information about Zydelig that is written for health professionals.

**What are the ingredients in Zydelig?**

**Active ingredient:** idelalisib

**Inactive ingredients:** microcrystalline cellulose, hydroxypropyl cellulose, croscarmellose sodium, sodium starch glycolate, and magnesium stearate. The tablet coating contains polyethylene glycol, talc, polyvinyl alcohol, titanium dioxide and FD&C Yellow #6 or Sunset Yellow FCF Aluminum Lake (for the 100 mg tablet) and red iron oxide (for the 150 mg tablet).

Manufactured and distributed by: Gilead Sciences, Inc. Foster City, CA 94404

©2018 Gilead Sciences, Inc. All rights reserved

For more information, call 1-800-445-3235 or go to [www.Zydelig.com](http://www.Zydelig.com).

MG

205858-GS-00X-

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: 01 2018

**Initial REMS approval: 07/23/2014**

**Most recent modification: 01/2018**

**NDA 205858**

**ZYDELIG® (idelalisib) tablets**

**Drug Class: Phosphoinositide-3 Kinase-Delta Inhibitor**

Gilead Sciences, Inc.  
333 Lakeside Drive  
Foster City, CA 94404  
Phone: (800) 445-3235

## **RISK EVALUATION AND MITIGATION STRATEGY (REMS)**

### **I. GOAL:**

The goal of the Zydelig REMS is to mitigate the risks of fatal and /or serious hepatotoxicity; fatal and/or serious and severe diarrhea or colitis; fatal and/or serious pneumonitis; fatal and/or serious infections, and fatal and serious intestinal perforation associated with Zydelig treatment by informing healthcare providers of the risks of

- fatal and/or serious hepatotoxicity
- fatal and/or serious and severe diarrhea or colitis
- fatal and/or serious pneumonitis
- fatal and/or serious infections
- fatal and serious intestinal perforation

### **II. REMS ELEMENTS**

#### **A. Communication Plan**

Gilead Sciences, Inc. (Gilead) must implement the following communication plan for healthcare providers who are likely to prescribe Zydelig. The communication plan must consist of the following:

1. **REMS Letter Healthcare Providers** - A *REMS Letter to Healthcare Providers* must be distributed within 30 calendar days of the approval of the REMS modification (01/26/2018). The letter must be distributed electronically to oncologists and hematologists who are likely to prescribe Zydelig. If a targeted healthcare provider's



email address is not available, or if an email is undeliverable, the provider will receive the letter through the mail. The *REMS Letter to Healthcare Providers* must inform healthcare providers of the risks of fatal and/or serious hepatotoxicity; fatal and/or serious and severe diarrhea or colitis; fatal and/or serious pneumonitis; fatal and/or serious infections; and fatal and serious intestinal perforation associated with Zydelig treatment. The letter must be accompanied by the Prescribing Information (with Medication Guide attached) and the *Zydelig REMS Fact Sheet*. The *REMS Letter to Healthcare Providers* must be available from the *Zydelig REMS Website* at the time of distribution and remain on the website for 12 months after approval of the REMS modification (01/26/2018).

2. **REMS Letter for Professional Societies** - A *REMS Letter for Professional Societies* must be distributed electronically within 30 calendar days of the approval of the REMS modification (01/26/2018). If a targeted Professional Society's email address is not available, or if an email is undeliverable, the letter must be sent through the mail. The *REMS Letter for Professional Societies* must inform the leadership of the professional societies described below of the risks of fatal and/or serious hepatotoxicity; fatal and/or serious and severe diarrhea or colitis; fatal and/or serious pneumonitis; fatal and/or serious infections; and fatal and serious intestinal perforation associated with Zydelig treatment. Gilead must request the leadership of the professional societies distribute this risk information to their memberships.

The *REMS Letter for Professional Societies* must be distributed to the following organizations:

- American Society of Clinical Oncology (ASCO)
  - American Society of Hematology (ASH)
  - Oncology Nursing Society (ONS)
  - National Comprehensive Cancer Network (NCCN)
  - Hematology Oncology Pharmacy Association (HOPA)
  - American Pharmacists Association (APhA)
  - American Society of Health-System Pharmacists (ASHP)
3. **REMS Fact Sheet** – A *Zydelig REMS Fact Sheet* must be distributed to Healthcare Providers. The *Zydelig REMS Fact Sheet* must be included in the mailings of the *REMS Letter to Healthcare Providers* and the *REMS Letter for Professional Societies* and must be available on the *Zydelig REMS Website*. Hard copies of the *Zydelig REMS Fact Sheet* must also be distributed by Gilead's sales representatives and medical field-based personnel to healthcare providers during follow-up details/visits with healthcare providers for the first 12 months after the approval of the REMS modification (01/26/2018). Gilead sales representatives and medical field-based personnel must orally review the risk messages contained in the *Zydelig REMS Fact Sheet* during the visit with the healthcare provider.
  4. **Journal Information Piece** - Gilead must publish in the following professional journals an information piece that includes the risks of fatal and/or serious hepatotoxicity, fatal

and/or serious and severe diarrhea or colitis, fatal and serious pneumonitis and serious intestinal perforation associated with Zydelig treatment.

- Journal of Clinical Oncology
- Blood
- New England Journal of Medicine
- Hematology Oncology Today
- Oncology & Hematology Review
- Leukemia and Lymphoma

The information piece will be published quarterly in each publication for one year following the REMS modification (01/26/2018).

5. **Dissemination of REMS information at scientific meetings** – The *Zydelig REMS Fact Sheet* and the Prescribing Information must be prominently displayed at scientific meetings where Gilead has a presence (e.g., booth) through the end of January 2019.
6. **REMS Program Website** –The *Zydelig REMS Program Website* ([www.zydeligrems.com](http://www.zydeligrems.com)) will continue for 12 months after the approval of this REMS modification (01/26/2018). The *Zydelig REMS Program Website* must include the option to print the currently approved Prescribing Information, Medication Guide, *REMS Letter for Healthcare Providers*, *REMS Factsheet*, and *Patient Safety Information Card*. The Zydelig product website must include a prominent REMS-specific link to the *Zydelig REMS Program Website*. All website information must be updated within 60 calendar days of approval of the REMS modification (01/26/2018).
7. **Zydelig Patient Safety Information Card** - A Patient Safety Information Card highlights the risks of Zydelig and includes information on the management of these risks. Gilead's sales representatives or medical field based personnel must distribute the *Patient Safety Information Cards* to prescribers within the first 12 months of approval of the REMS modification (01/26/2018). The *Patient Safety Information Card* must also be available on the Zydelig REMS Program Website for 12 months after the approval of the REMS modification (01/26/2018).

The following are part of the REMS and are appended.

- *REMS Letter to Healthcare Providers*
- *REMS Letter for Professional Societies*
- *REMS Fact Sheet*
- *Journal Information Piece*
- *Zydelig REMS Website*
- *Zydelig Patient Safety Information Card*

---

**B. Timetable for Submission of Assessments**

Gilead must submit REMS assessments to the FDA 18 months, 4 years, and 7 years from the date of the initial approval of the REMS (07/23/2014). To facilitate inclusion of as much information as possible while allowing reasonable time to prepare the submission, the reporting interval covered by each assessment should conclude no earlier than 60 calendar days before the submission date for that assessment so that it will be received by the FDA on or before the due date.

**Initial REMS approval: 07/23/2014**

**Most recent modification: 01/2018**

**NDA 205858**

**ZYDELIG® (idelalisib) tablets**

**Drug Class: Phosphoinositide-3 Kinase-Delta Inhibitor**

Gilead Sciences, Inc.  
333 Lakeside Drive  
Foster City, CA 94404  
Phone: (800) 445-3235

## **RISK EVALUATION AND MITIGATION STRATEGY (REMS)**

### **I. GOAL:**

The goal of the Zydelig REMS is to mitigate the risks of fatal and /or serious hepatotoxicity; fatal and/or serious and severe diarrhea or colitis; fatal and/or serious pneumonitis; fatal and/or serious infections, and fatal and serious intestinal perforation associated with Zydelig treatment by informing healthcare providers of the risks of

- fatal and/or serious hepatotoxicity
- fatal and/or serious and severe diarrhea or colitis
- fatal and/or serious pneumonitis
- fatal and/or serious infections
- fatal and serious intestinal perforation

### **II. REMS ELEMENTS**

#### **A. Communication Plan**

Gilead Sciences, Inc. (Gilead) must implement the following communication plan for healthcare providers who are likely to prescribe Zydelig. The communication plan must consist of the following:

1. **REMS Letter Healthcare Providers** - A *REMS Letter to Healthcare Providers* must be distributed within 30 calendar days of the approval of the REMS modification (01/26/2018). The letter must be distributed electronically to oncologists and hematologists who are likely to prescribe Zydelig. If a targeted healthcare provider's

email address is not available, or if an email is undeliverable, the provider will receive the letter through the mail. The *REMS Letter to Healthcare Providers* must inform healthcare providers of the risks of fatal and/or serious hepatotoxicity; fatal and/or serious and severe diarrhea or colitis; fatal and/or serious pneumonitis; fatal and/or serious infections; and fatal and serious intestinal perforation associated with Zydelig treatment. The letter must be accompanied by the Prescribing Information (with Medication Guide attached) and the *Zydelig REMS Fact Sheet*. The *REMS Letter to Healthcare Providers* must be available from the *Zydelig REMS Website* at the time of distribution and remain on the website for 12 months after approval of the REMS modification (01/26/2018).

2. **REMS Letter for Professional Societies** - A *REMS Letter for Professional Societies* must be distributed electronically within 30 calendar days of the approval of the REMS modification (01/26/2018). If a targeted Professional Society's email address is not available, or if an email is undeliverable, the letter must be sent through the mail. The *REMS Letter for Professional Societies* must inform the leadership of the professional societies described below of the risks of fatal and/or serious hepatotoxicity; fatal and/or serious and severe diarrhea or colitis; fatal and/or serious pneumonitis; fatal and/or serious infections; and fatal and serious intestinal perforation associated with Zydelig treatment. Gilead must request the leadership of the professional societies distribute this risk information to their memberships.

The *REMS Letter for Professional Societies* must be distributed to the following organizations:

- American Society of Clinical Oncology (ASCO)
  - American Society of Hematology (ASH)
  - Oncology Nursing Society (ONS)
  - National Comprehensive Cancer Network (NCCN)
  - Hematology Oncology Pharmacy Association (HOPA)
  - American Pharmacists Association (APhA)
  - American Society of Health-System Pharmacists (ASHP)
3. **REMS Fact Sheet** – A *Zydelig REMS Fact Sheet* must be distributed to Healthcare Providers. The *Zydelig REMS Fact Sheet* must be included in the mailings of the *REMS Letter to Healthcare Providers* and the *REMS Letter for Professional Societies* and must be available on the *Zydelig REMS Website*. Hard copies of the *Zydelig REMS Fact Sheet* must also be distributed by Gilead's sales representatives and medical field-based personnel to healthcare providers during follow-up details/visits with healthcare providers for the first 12 months after the approval of the REMS modification (01/26/2018). Gilead sales representatives and medical field-based personnel must orally review the risk messages contained in the *Zydelig REMS Fact Sheet* during the visit with the healthcare provider.
  4. **Journal Information Piece** - Gilead must publish in the following professional journals an information piece that includes the risks of fatal and/or serious hepatotoxicity, fatal

and/or serious and severe diarrhea or colitis, fatal and serious pneumonitis and serious intestinal perforation associated with Zydelig treatment.

- Journal of Clinical Oncology
- Blood
- New England Journal of Medicine
- Hematology Oncology Today
- Oncology & Hematology Review
- Leukemia and Lymphoma

The information piece will be published quarterly in each publication for one year following the REMS modification (01/26/2018).

5. **Dissemination of REMS information at scientific meetings** – The *Zydelig REMS Fact Sheet* and the Prescribing Information must be prominently displayed at scientific meetings where Gilead has a presence (e.g., booth) through the end of January 2019.
6. **REMS Program Website** –The *Zydelig REMS Program Website* ([www.zydeligrems.com](http://www.zydeligrems.com)) will continue for 12 months after the approval of this REMS modification (01/26/2018). The *Zydelig REMS Program Website* must include the option to print the currently approved Prescribing Information, Medication Guide, *REMS Letter for Healthcare Providers*, *REMS Factsheet*, and *Patient Safety Information Card*. The Zydelig product website must include a prominent REMS-specific link to the *Zydelig REMS Program Website*. All website information must be updated within 60 calendar days of approval of the REMS modification (01/26/2018).
7. **Zydelig Patient Safety Information Card** - A Patient Safety Information Card highlights the risks of Zydelig and includes information on the management of these risks. Gilead's sales representatives or medical field based personnel must distribute the *Patient Safety Information Cards* to prescribers within the first 12 months of approval of the REMS modification (01/26/2018). The *Patient Safety Information Card* must also be available on the Zydelig REMS Program Website for 12 months after the approval of the REMS modification (01/26/2018).

The following are part of the REMS and are appended.

- *REMS Letter to Healthcare Providers*
- *REMS Letter for Professional Societies*
- *REMS Fact Sheet*
- *Journal Information Piece*
- *Zydelig REMS Website*
- *Zydelig Patient Safety Information Card*

**B. Timetable for Submission of Assessments**

Gilead must submit REMS assessments to the FDA 18 months, 4 years, and 7 years from the date of the initial approval of the REMS (07/23/2014). To facilitate inclusion of as much information as possible while allowing reasonable time to prepare the submission, the reporting interval covered by each assessment should conclude no earlier than 60 calendar days before the submission date for that assessment so that it will be received by the FDA on or before the due date.



[Zydelig® logo]

[banded copy]  
**Zydelig REMS**

[headline]

## **FDA REQUIRED UPDATED REMS SAFETY INFORMATION**

[boxed copy]

### **Boxed Warning Risk of:**

- **Fatal and/or serious hepatotoxicity - *updated***
- **Fatal and/or serious and severe diarrhea or colitis - *updated***
- **Fatal and/or serious pneumonitis**
- **Fatal and/or serious infections - *updated***
- **Fatal and serious intestinal perforation**

[body copy]  
Month 20XX

Dear Healthcare Provider:

The FDA has required this safety notice as part of the Zydelig REMS (**Risk Evaluation and Mitigation Strategy**) to inform you about the recent **update to the incidence rates of fatal and serious toxicities in the Zydelig Boxed Warning** as follows:

[boxed copy]

### **WARNING: FATAL AND SERIOUS TOXICITIES: HEPATIC, SEVERE DIARRHEA, COLITIS, PNEUMONITIS, INFECTIONS and INTESTINAL PERFORATION**

- Fatal and/or serious hepatotoxicity occurred in 16% to 18% of Zydelig-treated patients. Monitor hepatic function prior to and during treatment. Interrupt and then reduce or discontinue Zydelig.
- Fatal and/or serious and severe diarrhea or colitis occurred in 14% to 20% of Zydelig-treated patients. Monitor for the development of severe diarrhea or colitis. Interrupt and then reduce or discontinue Zydelig.
- Fatal and/or serious pneumonitis occurred in 4% of Zydelig-treated patients. Monitor for pulmonary symptoms and interstitial infiltrates. Interrupt or discontinue Zydelig.
- Fatal and/or serious infections occurred in 21% to 48% of Zydelig-treated

patients. Monitor for signs and symptoms of infection. Interrupt Zydelig if infection is suspected.

- Fatal and serious Intestinal perforation can occur in Zydelig-treated patients across clinical trials. Discontinue Zydelig if intestinal perforation is suspected.

The updated incidence rates of fatal and serious toxicities in the Zydelig Boxed Warning reflect data from patients treated with Zydelig in combination with rituximab or with unapproved combination therapies.

Please see the enclosed **Zydelig REMS Fact Sheet**, a non-promotional fact sheet reviewed by the FDA, for more detailed safety information. Be sure to give the **Zydelig Patient Safety Information Card** to all patients. This card, additional copies of the fact sheet, and other important information are available at: [www.ZydeligREMS.com](http://www.ZydeligREMS.com).

Zydelig is a kinase inhibitor indicated for the treatment of patients with:

- Relapsed chronic lymphocytic leukemia (CLL) in combination with rituximab, in patients for whom rituximab alone would be considered appropriate therapy due to other co-morbidities.
- Relapsed follicular B-cell non-Hodgkin lymphoma (FL) in patients who have received at least two prior systemic therapies.
- Relapsed small lymphocytic lymphoma (SLL) in patients who have received at least two prior systemic therapies.

Limitation of use:

Zydelig is not indicated and is not recommended for first-line treatment of any patient. Zydelig is not indicated and is not recommended in combination with bendamustine and/or rituximab for the treatment of FL.

This letter does not contain the complete safety profile for Zydelig. Please review the enclosed Prescribing Information.

[subhead]

**Reporting Adverse Events**

[body copy]

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088. Healthcare Providers should report all suspected adverse events associated with Zydelig to the FDA or to Gilead at 1-800-445-3235.

Sincerely,

[signature]

William Guyer, Pharm.D.  
Senior Vice President, Medical Affairs

[signoff]

**Gilead Sciences, Inc.** 333 Lakeside Drive Foster City, CA  
94404 USA

[www.ZydeligREMS.com](http://www.ZydeligREMS.com)

*Phone* 650 574 3000 *facsimile* 650 578 9264  
REMS-ZYD-0034

**From:** Gilead Sciences, Inc.

**To:** <Healthcare Provider Name>

**Subject:** Revised Boxed Warning for Zydelig: Update to Incidence Rates of Fatal and Serious Toxicities

[ZYDELIG® logo]

[banded copy]

**Zydelig REMS**

[headline]

**FDA REQUIRED UPDATED REMS SAFETY INFORMATION**

[boxed copy]

**Boxed Warning includes the Risk of:**

- Fatal and/or serious hepatotoxicity - *updated*
- Fatal and/or serious and severe diarrhea or colitis - *updated*
- Fatal and/or serious pneumonitis
- Fatal and/or serious infections - *updated*
- Fatal and serious intestinal perforation

Month 20XX

Dear Healthcare Provider:

The FDA has required this safety notice as part of the Zydelig REMS (**R**isk **E**valuation and **M**itigation **S**trategy) to inform you about the recent update to the **incidence rates of fatal and serious toxicities in the Zydelig Boxed Warning** as follows:

[boxed copy]

**WARNING: FATAL AND SERIOUS TOXICITIES: HEPATIC, SEVERE DIARRHEA, COLITIS, PNEUMONITIS, INFECTIONS, and INTESTINAL PERFORATION**

- Fatal and/or serious hepatotoxicity occurred in 16 to 18% of Zydelig-treated patients. Monitor hepatic function prior to and during treatment. Interrupt and then reduce or discontinue Zydelig.
- Fatal and/or serious and severe diarrhea or colitis occurred in 14% to 20% of Zydelig-treated patients. Monitor for the development of severe diarrhea or

colitis. Interrupt or discontinue Zydelig.

- Fatal and/or serious pneumonitis occurred in 4% of Zydelig-treated patients. Monitor for pulmonary symptoms and interstitial infiltrates. Interrupt or discontinue Zydelig.
- Fatal and/or serious infections occurred in 21% to 48% of Zydelig-treated patients. Monitor for signs and symptoms of infection. Interrupt Zydelig if infection is suspected.
- Fatal and serious intestinal perforation can occur in Zydelig-treated patients across clinical trials. Discontinue Zydelig if intestinal perforation is suspected.

The updated incidence rates of fatal and serious toxicities in the Zydelig Boxed Warning reflect data from patients treated with Zydelig in combination with rituximab or with unapproved combination therapies.

Please see the **Zydelig REMS Fact Sheet** [insert direct link], a non-promotional fact sheet reviewed by the FDA, for more detailed safety information. Please give the **Zydelig Patient Safety Information Card** [insert direct link] to all patients. The card, the fact sheet, and other important information are available at: [www.ZydeligREMS.com](http://www.ZydeligREMS.com).

Zydelig is a kinase inhibitor indicated for the treatment of patients with

- Relapsed chronic lymphocytic leukemia (CLL) in combination with rituximab, in patients for whom rituximab alone would be considered appropriate therapy due to other co-morbidities.
- Relapsed follicular B-cell non-Hodgkin lymphoma (FL) in patients who have received at least two prior systemic therapies.
- Relapsed small lymphocytic lymphoma (SLL) in patients who have received at least two prior systemic therapies.

Limitation of use:

Zydelig is not indicated and is not recommended for first-line treatment of any patient. Zydelig is not indicated and is not recommended in combination with bendamustine and/or rituximab for the treatment of FL.

This email does not contain the complete safety profile for Zydelig. To review the Prescribing Information, including complete **BOXED WARNING** and Medication Guide, see links below:

[links]

**Prescribing Information      Medication Guide**

[subhead]

**Reporting Adverse Events**

[body copy]

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [link] [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088. Healthcare Providers should report all suspected adverse events associated with Zydelig to the FDA or to Gilead at 1-800-445-3235.

Sincerely,

[signature]

William Guyer, Pharm.D.  
Senior Vice President, Medical Affairs

[signoff]

**Gilead Sciences, Inc.** 333 Lakeside Drive Foster City, CA 94404 USA  
*Phone* 650 574 3000 *facsimile* 650 578 9264  
REMS-ZYD-0035

[Zydelig® logo]

[banded copy]

**Zydelig REMS**

[headline]

## **FDA REQUIRED UPDATED REMS SAFETY INFORMATION**

[boxed copy]

### **Boxed Warning Risk of:**

- Fatal and/or serious hepatotoxicity - *updated*
- Fatal and/or serious and severe diarrhea or colitis - *updated*
- Fatal and/or serious pneumonitis
- Fatal and/or serious infections - *updated*
- Fatal and serious intestinal perforation

[body copy]

Month 20XX

Dear <name>:

The FDA has required Gilead Sciences to distribute this safety notice to the [Professional Organization] as part of the Zydelig REMS (**R**isk **E**valuation and **M**itigation **S**trategy). The **incidence rates of fatal and serious toxicities have been updated in the Boxed Warning** for Zydelig. We request that you distribute this information to your members about the following serious risks of Zydelig:

[boxed copy]

### **WARNING: FATAL AND SERIOUS TOXICITIES: HEPATIC, SEVERE DIARRHEA or COLITIS, PNEUMONITIS, INFECTIONS and INTESTINAL PERFORATION**

- Fatal and/or serious hepatotoxicity occurred in 16 to 18% of Zydelig-treated patients. Monitor hepatic function prior to and during treatment. Interrupt and then reduce or discontinue Zydelig.
- Fatal and/or serious and severe diarrhea or colitis occurred in 14% to 20% of Zydelig-treated patients. Monitor for the development of severe diarrhea or colitis. Interrupt and then reduce or discontinue Zydelig.
- Fatal and/or serious pneumonitis occurred in 4% of Zydelig-treated patients. Monitor for pulmonary symptoms and bilateral interstitial infiltrates. Interrupt and or discontinue Zydelig.



- Fatal and/or serious infections occurred in 21% to 48% of Zydelig-treated patients. Monitor for signs and symptoms of infection. Interrupt Zydelig if infection is suspected.
- Fatal and serious intestinal perforation can occur in Zydelig-treated patients across clinical trials. Discontinue Zydelig if intestinal perforation is suspected.

The updated incidence rates of fatal and serious toxicities in the Zydelig Boxed Warning reflect data from patients treated with Zydelig in combination with rituximab or with unapproved combination therapies.

Please see the enclosed **Zydelig REMS Fact Sheet**, a non-promotional fact sheet reviewed by the FDA, for more detailed safety information. Please encourage your members to provide the **Zydelig Patient Safety Information Card** to all patients. This card, additional copies of the fact sheet, and other important information are available at: [www.ZydeligREMS.com](http://www.ZydeligREMS.com).

Zydelig is a kinase inhibitor indicated for the treatment of patients with

- Relapsed chronic lymphocytic leukemia (CLL) in combination with rituximab, in patients for whom rituximab alone would be considered appropriate therapy due to other co-morbidities.
- Relapsed follicular B-cell non-Hodgkin lymphoma (FL) in patients who have received at least two prior systemic therapies.
- Relapsed small lymphocytic lymphoma (SLL) in patients who have received at least two prior systemic therapies.

Limitation of use:

Zydelig is not indicated and is not recommended for first-line treatment of any patient. Zydelig is not indicated and is not recommended in combination with bendamustine and/or rituximab for the treatment of FL.

This letter does not contain the complete safety profile for Zydelig. Please review the enclosed Prescribing Information.

[subhead]

**Reporting Adverse Events**

[body copy]

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088. Healthcare Providers should report all suspected adverse events associated with Zydelig to the FDA or to Gilead at 1-800-445-3235.

Sincerely,

[signature]

William Guyer, Pharm.D.  
Senior Vice President, Medical Affairs

[signoff]

**Gilead Sciences, Inc.** 333 Lakeside Drive Foster City, CA  
94404 USA

[www.ZydeligREMS.com](http://www.ZydeligREMS.com)

*Phone* 650 574 3000 *facsimile* 650 578 9264

REMS-ZYD-0036

From: Gilead Sciences, Inc.

To: < Name, Professional Society>

Subject: Revised Boxed Warning for Zydelig: Update to Incidence Rates of Fatal and Serious Toxicities

[Zydelig® logo]

[banded copy]

**Zydelig REMS**

[headline]

**FDA REQUIRED UPDATED REMS SAFETY INFORMATION**

[boxed copy]

**Boxed Warning includes the Risk of:**

- Fatal and/or serious hepatotoxicity - *updated*
- Fatal and/or serious and severe diarrhea or colitis - *updated*
- Fatal and/or serious pneumonitis
- Fatal and/or serious infections - *updated*
- Fatal and serious intestinal perforation

[body copy]

Month 20XX

**IMPORTANT SAFETY NOTICE**

Dear [name]:

The FDA has required Gilead Sciences to distribute this safety notice to the [Professional Organization] as part of the Zydelig REMS (**R**isk **E**valuation and **M**itigation **S**trategy). The **incidence rates of fatal and serious toxicities have been updated in the Boxed Warning** for Zydelig. We request that you distribute this information to your members about the following serious risks of Zydelig:

[boxed copy]

**WARNING: FATAL AND SERIOUS TOXICITIES: HEPATIC, SEVERE DIARRHEA, COLITIS, PNEUMONITIS, INFECTIONS and INTESTINAL PERFORATION**

- Fatal and/or serious hepatotoxicity occurred in 16% to 18% of Zydelig-treated patients. Monitor hepatic function prior to and during treatment. Interrupt and

then reduce or discontinue Zydelig.

- Fatal and/or serious and severe diarrhea or colitis occurred in 14% to 20% of Zydelig-treated patients. Monitor for the development of severe diarrhea or colitis. Interrupt and then reduce or discontinue Zydelig.
- Fatal and/or serious pneumonitis occurred in 4% of Zydelig-treated patients. Monitor for pulmonary symptoms and interstitial infiltrates. Interrupt or discontinue Zydelig.
- Fatal and/or serious infections occurred in 21% to 48% of Zydelig-treated patients. Monitor for signs and symptoms of infection. Interrupt Zydelig if infection is suspected.
- Fatal and serious intestinal perforation can occur in Zydelig-treated patients across clinical trials. Discontinue Zydelig if intestinal perforation is suspected.

The updated incidence rates of fatal and serious toxicities in the Zydelig Boxed Warning reflect data from patients treated with Zydelig in combination with rituximab or with unapproved combination therapies.

Please see the **Zydelig REMS Fact Sheet** [insert direct link], a non-promotional fact sheet reviewed by the FDA, for more detailed safety information. Please encourage your members to provide the **Zydelig Patient Safety Information Card** [insert direct link] to all patients. The card, the fact sheet, and other important information are available at: [link] [www.ZydeligREMS.com](http://www.ZydeligREMS.com).

Zydelig is a kinase inhibitor indicated for the treatment of patients with

- Relapsed chronic lymphocytic leukemia (CLL) in combination with rituximab, in patients for whom rituximab alone would be considered appropriate therapy due to other co-morbidities.
- Relapsed follicular B-cell non-Hodgkin lymphoma (FL) in patients who have received at least two prior systemic therapies.
- Relapsed small lymphocytic lymphoma (SLL) in patients who have received at least two prior systemic therapies.

Limitation of use:

Zydelig is not indicated and is not recommended for first-line treatment of any patient. Zydelig is not indicated and is not recommended in combination with bendamustine and/or rituximab for the treatment of FL.

This letter does not contain the complete safety profile for Zydelig. To review the Prescribing Information, including complete **BOXED WARNING** and Medication Guide, see links below:

[links]

**Prescribing Information      Medication Guide**

[subhead]

**Reporting Adverse Events**

[body copy]

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [link] [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088. Healthcare Providers should report all suspected adverse events associated with Zydelig to the FDA or to Gilead at 1-800-445-3235.

Sincerely,

[signature]

William Guyer, Pharm.D.  
Senior Vice President, Medical Affairs

[signoff]

**Gilead Sciences, Inc.** 333 Lakeside Drive Foster City, CA  
94404 USA  
*Phone* 650 574 3000 *facsimile* 650 578 9264  
REMS-ZYD-0037

[www.ZydeligREMS.com](http://www.ZydeligREMS.com)

[Zydelig® logo]

[banded copy]

## **Zydelig REMS Fact Sheet**

[headline]

### **FDA REQUIRED Zydelig (idelalisib) UPDATED REMS SAFETY INFORMATION**

[boxed copy]

#### **Boxed Warning for the Risk of:**

- **Fatal and/or serious hepatotoxicity - *updated***
- **Fatal and/or serious and severe diarrhea or colitis - *updated***
- **Fatal and/or serious pneumonitis**
- **Fatal and /or serious infections - *updated***
- **Fatal and serious intestinal perforation**

[body copy]

#### **Fatal and/or Serious Hepatotoxicity**

- Fatal and/or serious hepatotoxicity occurred in 18% of patients treated with Zydelig monotherapy and 16% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies.
- Elevations in ALT or AST greater than 5 times the upper limit of normal have occurred. These findings were generally observed within the first 12 weeks of treatment and were reversible with dose interruption. After resumption of treatment at a lower dose, 26% of patients had recurrence of ALT and AST elevations. Discontinue Zydelig for recurrent hepatotoxicity.
- Avoid concurrent use of Zydelig with other drugs that may cause liver toxicity.
- Monitor ALT and AST in all patients every 2 weeks for the first 3 months of treatment, every 4 weeks for the next 3 months, then every 1 to 3 months thereafter. Monitor weekly for liver toxicity if the ALT or AST rises above 3 times the upper limit of normal until resolved. Withhold Zydelig if the ALT or AST is greater than 5 times the upper limit of normal, and continue to monitor AST, ALT, and total bilirubin weekly until the abnormality is resolved.

#### **Fatal and/or Serious and Severe Diarrhea or Colitis**

- Severe diarrhea or colitis (Grade 3 or higher) occurred in 14% of patients treated with Zydelig monotherapy and 20% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies. Diarrhea can occur at any time.
- Avoid concurrent use of Zydelig and other drugs that cause diarrhea. Diarrhea due to Zydelig responds poorly to antimotility agents. Median time to resolution

ranged between one week and one month across trials following interruption of Zydelig therapy and in some instances, use of corticosteroids.

[body copy]

**Fatal and/or Serious Pneumonitis**

- Fatal and serious pneumonitis occurred in patients treated with Zydelig. Clinical manifestations included interstitial infiltrates and organizing pneumonia. In randomized clinical trials of combination therapies, pneumonitis occurred in 4% of patients treated with Zydelig compared to 1% on the comparator arms. Time to onset of pneumonitis ranged from <1 to 15 months. Monitor patients on Zydelig for pulmonary symptoms.
- In patients taking Zydelig who present with pulmonary symptoms such as cough, dyspnea, hypoxia, interstitial infiltrates on a radiologic exam, or a decline by more than 5% in oxygen saturation, interrupt Zydelig until the etiology has been determined. If symptomatic pneumonitis or organizing pneumonia is diagnosed, initiate appropriate treatment with corticosteroids and permanently discontinue Zydelig.

[body copy]

**Fatal and/or Serious Infections**

- Fatal and/or serious infections occurred in 21% of patients treated with Zydelig monotherapy and 48% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies. The most common infections were pneumonia, sepsis, and febrile neutropenia. Treat infections prior to initiation of Zydelig therapy.
- Monitor patients on Zydelig for signs and symptoms of infection and interrupt Zydelig for Grade 3 or higher infection.
- Serious or fatal *Pneumocystis jirovecii* pneumonia (PJP) or cytomegalovirus (CMV) occurred in <1% of patients treated with Zydelig. Provide PJP prophylaxis during treatment with Zydelig. Interrupt Zydelig in patients with suspected PJP infection of any grade, and permanently discontinue Zydelig if PJP infection of any grade is confirmed. Regular clinical and laboratory monitoring for CMV infection is recommended in patients with history of CMV infection or positive CMV serology at the start of treatment with Zydelig. Interrupt Zydelig in the setting of positive CMV PCR or antigen test until the viremia has resolved. If Zydelig is subsequently resumed, patients should be monitored by PCR or antigen test for CMV reactivation at least monthly.

[body copy]

**Fatal and Serious Intestinal Perforation**

- Fatal and serious intestinal perforation occurred in Zydelig-treated patients. At the time of perforation, some patients had moderate to severe diarrhea.

- Advise patients to promptly report any new or worsening abdominal pain, chills, fever, nausea, or vomiting.
- Discontinue Zydelig permanently in patients who experience intestinal perforation.

### INDICATION:

Zydelig is a kinase inhibitor indicated for the treatment of patients with:

- Relapsed chronic lymphocytic leukemia (CLL) in combination with rituximab, in patients for whom rituximab alone would be considered appropriate therapy due to other co-morbidities.
- Relapsed follicular B-cell non-Hodgkin lymphoma (FL) in patients who have received at least two prior systemic therapies.
- Relapsed small lymphocytic lymphoma (SLL) in patients who have received at least two prior systemic therapies

#### Limitation of use:

Zydelig is not indicated and is not recommended for first-line treatment of any patient. Zydelig is not indicated and is not recommended in combination with bendamustine and/or rituximab for the treatment of FL.

[subhead]

### WHAT IS THE ZYDELIG REMS?

[body copy]

A REMS (**R**isk **E**valuation and **M**itigation **S**trategy) is a program required by the FDA to manage known or potential serious risks associated with a drug product. FDA has determined that a REMS is necessary to ensure that the benefits of Zydelig outweigh its risks. The purpose of the Zydelig REMS is to inform Healthcare Providers of fatal and serious risks of hepatotoxicity, severe diarrhea or colitis, pneumonitis, infections, and intestinal perforation. This fact sheet is required by the FDA as part of the Zydelig REMS program.

A **Patient Safety Information Card** is also available for you to provide to your patients. The card reminds patients of the symptoms of the serious risks of Zydelig and should be shown to any healthcare provider involved in your patient's care. The card and other information and resources are available at [www.ZydeligREMS.com](http://www.ZydeligREMS.com)

This fact sheet does not contain the complete safety profile for Zydelig. Please see the Prescribing Information, including the **BOXED WARNING** and Medication Guide.

[subhead]

### REPORTING ADVERSE EVENTS



[body copy]

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit [link] [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088. Healthcare Providers should report all suspected adverse events associated with Zydelig to the FDA or to Gilead at 1-800-445-3235.

[signoff]

© 201X Gilead Sciences, Inc. All rights reserved. REMS-ZYD-0038 XX/20XX

Gilead, the Gilead logo, and ZYDELIG are trademarks of Gilead Sciences, Inc., or one of its related companies.

[Zydelig® logo]

[banded copy]

**FDA REQUIRED Updated Safety Information for Zydelig® (idelalisib)**

**Safety information updates were made to the Boxed Warning, including incidence rates of fatal and serious toxicities.**

[boxed copy]

**BOXED WARNING For Zydelig (Idelalisib)**

**Risk of:**

- **Fatal and/or serious hepatotoxicity - *updated***
- **Fatal and/or serious and severe diarrhea or colitis - *updated***
- **Fatal and/or serious pneumonitis**
- **Fatal and/or serious infections - *updated***
- **Fatal and serious intestinal perforation**

**Fatal and/or Serious Hepatotoxicity**

- Fatal and/or serious hepatotoxicity occurred in 18% of patients treated with Zydelig monotherapy and 16% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies.
- Elevations in ALT or AST greater than 5 times the upper limit of normal have occurred. These findings were generally observed within the first 12 weeks of treatment and were reversible with dose interruption. After resumption of treatment at a lower dose, 26% of patients had recurrence of ALT and AST elevations. Discontinue Zydelig for recurrent hepatotoxicity.
- Avoid concurrent use of Zydelig with other drugs that may cause liver toxicity.
- Monitor ALT and AST in all patients every 2 weeks for the first 3 months of treatment, every 4 weeks for the next 3 months, then every 1 to 3 months thereafter. Monitor weekly for liver toxicity if the ALT or AST rises above 3 times the upper limit of normal until resolved. Withhold Zydelig if the ALT or AST is greater than 5 times the upper limit of normal, and continue to monitor AST, ALT and total bilirubin weekly until the abnormality is resolved.

**Fatal and/or Serious and Severe Diarrhea or Colitis**

- Severe diarrhea or colitis (Grade 3 or higher) occurred in 14% of patients treated with Zydelig monotherapy and 20% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies. Diarrhea can occur at any time.
- Avoid concurrent use of Zydelig and other drugs that cause diarrhea. Diarrhea due to Zydelig responds poorly to antimotility agents. Median time to resolution

ranged between one week and one month across trials, following interruption of Zydelig therapy and in some instances, use of corticosteroids.

[body copy]

#### **Fatal and/or Serious Pneumonitis**

- Fatal and serious pneumonitis occurred in patients treated with Zydelig. Clinical manifestations included interstitial infiltrates and organizing pneumonia.
- In randomized clinical trials of combination therapies, pneumonitis occurred in 4% of patients treated with Zydelig compared to 1% on the comparator arms. Time to onset of pneumonitis ranged from <1 to 15 months. Monitor patients on Zydelig for pulmonary symptoms.
- In patients taking Zydelig who present with pulmonary symptoms such as cough, dyspnea, hypoxia, interstitial infiltrates on a radiologic exam, or a decline by more than 5% in oxygen saturation, interrupt Zydelig until the etiology has been determined. If symptomatic pneumonitis or organizing pneumonia is diagnosed, initiate appropriate treatment with corticosteroids and permanently discontinue Zydelig.

#### **Fatal and/or Serious Infections**

- Fatal and/or serious infections occurred in 21% of patients treated with Zydelig monotherapy and 48% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies. The most common infections were pneumonia, sepsis, and febrile neutropenia. Treat infections prior to initiation of Zydelig therapy.
- Monitor patients on Zydelig for signs and symptoms of infection and interrupt Zydelig for Grade 3 or higher infection.
- Serious or fatal *Pneumocystis jirovecii* pneumonia (PJP) or cytomegalovirus (CMV) occurred in <1% of patients treated with Zydelig. Provide PJP prophylaxis during treatment with Zydelig. Interrupt Zydelig in patients with suspected PJP infection of any grade, and permanently discontinue Zydelig if PJP infection of any grade is confirmed. Regular clinical and laboratory monitoring for CMV infection is recommended in patients with history of CMV infection or positive CMV serology at the start of treatment with Zydelig. Interrupt Zydelig in the setting of positive CMV PCR or antigen test until the viremia has resolved. If Zydelig is subsequently resumed, patients should be monitored by PCR or antigen test for CMV reactivation at least monthly.

#### **Fatal and Serious Intestinal Perforation**

- Fatal and serious intestinal perforation occurred in Zydelig-treated patients.
- At the time of perforation, some patients had moderate to severe diarrhea.
- Advise patients to promptly report any new or worsening abdominal pain, chills, fever, nausea, or vomiting.
- Discontinue Zydelig permanently in patients who experience intestinal perforation.

## INDICATION:

Zydelig is a kinase inhibitor indicated for the treatment of patients with:

- Relapsed chronic lymphocytic leukemia (CLL) in combination with rituximab, in patients for whom rituximab alone would be considered appropriate therapy due to other co-morbidities.
- Relapsed follicular B-cell non-Hodgkin lymphoma (FL) in patients who have received at least two prior systemic therapies.
- Relapsed small lymphocytic lymphoma (SLL) in patients who have received at least two prior systemic therapies.

Limitation of use:

Zydelig is not indicated and is not recommended for first-line treatment of any patient. Zydelig is not indicated and is not recommended in combination with bendamustine and/or rituximab for the treatment of FL.

|                                                                                                                                                                                                    |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| You are encouraged to report negative side effects of Zydelig to Gilead at 1-800-445-3235 and/or the FDA at <a href="http://www.fda.gov/medwatch">www.fda.gov/medwatch</a> or call 1-800-FDA-1088. |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

[disclaimer]

*This journal piece is part of the FDA-required Zydelig REMS. Visit [www.ZydeligREMS.com](http://www.ZydeligREMS.com) for more information.*

*For complete safety information, see the Prescribing Information available at [www.ZydeligREMS.com](http://www.ZydeligREMS.com).*

[signoff]

© 201X Gilead Sciences, Inc. All rights reserved. REMS-ZYD-0040 XX/20XX

Gilead, the Gilead logo, and ZYDELIG are trademarks of Gilead Sciences, Inc., or one of its related companies.

## **Zydelig Patient Safety Information Card**

### **Important Safety Information for Patients Taking Zydelig**

There are serious risks to understand when taking Zydelig.

Be certain to get regular blood tests as scheduled by your doctor.

Also, if you experience any of the following symptoms, you should immediately call your doctor and seek emergency medical care.

These symptoms include:

- stomach (abdominal) pain or swelling
- persistent or worsening nausea or vomiting
- severe diarrhea
- dark urine color
- bloody or tar-colored stool
- shortness of breath, difficulty breathing, or wheezing
- new or worsening cough
- persistent fever (temperature over 101°F)

Remember! Call your doctor and get emergency medical care right away if you have any of these symptoms and show this card to any doctor involved in your care.

## **Zydelig Patient Safety Card**

### **Important Safety Information for Healthcare Providers**

This patient is taking Zydelig (idelalisib) therapy for the treatment of leukemia or lymphoma. These are some of the serious risks associated with Zydelig:

- Fatal and/or serious hepatotoxicity
- Fatal and/or serious and severe diarrhea or colitis
- Fatal and/or serious pneumonitis
- Fatal and/or serious infections
- Fatal and serious intestinal perforation

Healthcare Providers:

- Evaluate urgently
- Provide all supportive care
- Contact the prescribing physician (see contact information at left) as soon as possible to coordinate care if a patient presents with signs and symptoms of the risks noted here

For more information about Zydelig, please refer to the full Prescribing Information. In case of safety concerns, call Gilead at 1-800-445-3235.

**Patients receiving Zydelig should carry this card at all times. Show this card to any doctor involved in your health care.**

Patient Name \_\_\_\_\_

Zydelig Prescriber Name \_\_\_\_\_

Prescriber Phone Number \_\_\_\_\_

© 201X Gilead Sciences, Inc.

All rights reserved. REMS-ZYD-0039 XX/20XX

Gilead and ZYDELIG are trademarks of Gilead Sciences, Inc., or one of its related companies.

[Zydelig® logo]

[links]

| PRESCRIBING<br>INFORMATION | MEDICATION GUIDE |
|----------------------------|------------------|
|----------------------------|------------------|

[header copy]

Zydelig (idelalisib) REMS (**Risk Evaluation and Mitigation Strategy**)

[body copy]

### **What is the Zydelig REMS?**

A REMS (**Risk Evaluation and Mitigation Strategy**) is a program required by the Food and Drug Administration (FDA) to manage known or potential serious risks associated with a drug product.

The purpose of the Zydelig REMS is to inform healthcare providers about the following serious **risks** of Zydelig:

#### **Fatal and/or Serious Hepatotoxicity**

- Fatal and/or serious hepatotoxicity occurred in 18% of patients treated with Zydelig monotherapy and 16% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies.

#### **Fatal and/or Serious and Severe Diarrhea or Colitis**

- Fatal and/or serious and severe diarrhea or colitis (Grade 3 or higher) occurred in 14% of patients treated with Zydelig monotherapy and 20% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies. Diarrhea can occur at any time.

#### **Fatal and/or Serious Pneumonitis**

Fatal and serious pneumonitis occurred in patients treated with Zydelig. Clinical manifestations included interstitial infiltrates and organizing pneumonia. In randomized clinical trials of combination therapies, pneumonitis occurred in 4% of patients treated with Zydelig compared to 1% on the comparator arms. Time to onset of pneumonitis ranged from <1 to 15 months. Monitor patients on Zydelig for pulmonary symptoms.

#### **Fatal and/or Serious Infections**

- Fatal and/or serious infections occurred in 21% of patients treated with Zydelig monotherapy and 48% of patients treated with Zydelig in combination with rituximab or with unapproved combination therapies. The most common infections were pneumonia, sepsis, and febrile neutropenia.

[body copy]

**Fatal and Serious Intestinal Perforation**

- Fatal and serious intestinal perforation occurred in Zydelig-treated patients.
- At the time of perforation, some patients had moderate to severe diarrhea.

**Zydelig Fact Sheet:** A non-promotional fact sheet, reviewed by the FDA, with more detailed safety information on these risks is available. (See link in the box labeled "Materials for Healthcare Providers")

**Zydelig Patient Safety Information Card:** This card should be given to all patients by Zydelig prescribers and should be carried by patients on Zydelig at all times. Patients should show this card to any healthcare professional that sees them in a health-related encounter. (See link in the box labeled "Materials for Patients")

[body copy]

[boxed links]

**Materials for Healthcare Providers**

Zydelig REMS Letter to Healthcare Providers

[link] **Download PDF**

Zydelig REMS Fact Sheet

[link] **Download PDF**

**Materials for Patients**

Medication Guide

[link] **Download PDF**

Zydelig Patient Safety Information Card

[link] **Download PDF**

**INDICATION:**

Zydelig is a kinase inhibitor indicated for the treatment of patients with

- Relapsed chronic lymphocytic leukemia (CLL) in combination with rituximab, in patients for whom rituximab alone would be considered appropriate therapy due to other co-morbidities.
- Relapsed follicular B-cell non-Hodgkin lymphoma (FL) in patients who have received at least two prior systemic therapies.
- Relapsed small lymphocytic lymphoma (SLL) in patients who have received at least two prior systemic therapies.



Limitation of use:

Zydelig is not indicated and is not recommended for first-line treatment of any patient.  
Zydelig is not indicated and is not recommended in combination with bendamustine and/or rituximab for the treatment of FL.

You are encouraged to report negative side effects of Zydelig to Gilead at 1-800-445-3235 and/or the FDA at [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

**This site is intended for US Healthcare Professionals.**

[signoff]

©201X Gilead Sciences, Inc. All rights reserved. REMS-ZYD-0033 XX/201X

Gilead, the Gilead logo, ZYDELIG, and the ZYDELIG logo are trademarks of Gilead Sciences, Inc., or one of its related companies.

[bottom links]

**Privacy Policy**

**Terms of Use**

**Contact Us**