

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VIBERZI safely and effectively. See full prescribing information for VIBERZI.

VIBERZI (eluxadoline) tablets, for oral use, CIV
Initial U.S. Approval: 2015

-----RECENT MAJOR CHANGES-----

Contraindications, patients without a gallbladder (4) 04/2017
Warnings and Precautions (5.1, 5.2) 04/2017

-----INDICATIONS AND USAGE-----

VIBERZI is a mu-opioid receptor agonist, indicated in adults for the treatment of irritable bowel syndrome with diarrhea (IBS-D). (1)

-----DOSAGE AND ADMINISTRATION-----

- The recommended dosage in adults is 100 mg twice daily taken with food. (2)
The recommended dosage is 75 mg twice daily taken with food in patients who:
 - are unable to tolerate the 100 mg dose. (2, 6.1)
 - are receiving concomitant OATP1B1 inhibitors. (2, 7)
 - have mild or moderate hepatic impairment. (2, 8.6)
- Discontinue VIBERZI in patients who develop severe constipation. (2)
- If a dose is missed, take the next dose at the regular time; do not take 2 doses at once. (2)

-----DOSAGE FORMS AND STRENGTHS-----

75 mg and 100 mg tablets (3)

-----CONTRAINDICATIONS-----

- patients without a gallbladder (4)
- known or suspected biliary duct obstruction, or sphincter of Oddi disease or dysfunction (4)

- alcoholism, alcohol abuse, alcohol addiction, or drink more than 3 alcoholic beverages/day (4)
- a history of pancreatitis; structural diseases of the pancreas, including known or suspected pancreatic duct obstruction (4)
- severe hepatic impairment (Child-Pugh Class C) (4, 8.6)
- severe constipation or sequelae from constipation, or known or suspected mechanical gastrointestinal obstruction (4)

-----WARNINGS AND PRECAUTIONS-----

Pancreatitis and Sphincter of Oddi Spasm: Monitor patients for new or worsening abdominal pain, with or without nausea and vomiting, or acute biliary pain with liver or pancreatic enzyme elevations; immediately discontinue VIBERZI and seek medical attention if symptoms develop. (5.1, 5.2)

-----ADVERSE REACTIONS-----

Most common adverse reactions (>5%) are constipation, nausea and abdominal pain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Allergan at 1- 800-433-8871 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----DRUG INTERACTIONS-----

See full prescribing information. (7)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

Revised: 04/2017

FULL PRESCRIBING INFORMATION CONTENTS*

- 1 INDICATIONS AND USAGE
- 2 DOSAGE AND ADMINISTRATION
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
 - 5.1 Pancreatitis
 - 5.2 Sphincter of Oddi Spasm
- 6 ADVERSE REACTIONS
 - 6.1 Clinical Trials Experience
- 7 DRUG INTERACTIONS
- 8 USE IN SPECIFIC POPULATIONS
 - 8.1 Pregnancy
 - 8.2 Lactation
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
 - 8.6 Hepatic Impairment
- 9 DRUG ABUSE AND DEPENDENCE
 - 9.1 Controlled Substance

- 9.2 Abuse
- 9.3 Dependence
- 10 OVERDOSAGE
- 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES
- 16 HOW SUPPLIED/STORAGE AND HANDLING
- 17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

VIBERZI is indicated in adults for the treatment of irritable bowel syndrome with diarrhea (IBS-D).

2 DOSAGE AND ADMINISTRATION

The recommended dosage of VIBERZI is 100 mg taken orally twice daily with food.

The recommended dosage of VIBERZI is 75 mg taken orally twice daily with food in patients who:

- are unable to tolerate the 100 mg dose of VIBERZI [see *Adverse Reactions (6.1)*].
- are receiving concomitant OATP1B1 inhibitors [see *Drug Interactions (7)*].
- have mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment [see *Use in Specific Population (8.6)*, *Clinical Pharmacology (12.3)*].

Discontinue VIBERZI in patients who develop severe constipation.

Instruct patients if they miss a dose, take the next dose at the regular time and not to take 2 doses at the same time to make up for a missed dose.

3 DOSAGE FORMS AND STRENGTHS

- 75 mg tablets: capsule-shaped tablets are coated in pale-yellow to light tan color debossed with “FX75” on one side. Each tablet contains 75 mg eluxadoline.
- 100 mg tablets: capsule-shaped tablets are coated in pink-orange to peach color debossed with “FX100” on one side. Each tablet contains 100 mg eluxadoline.

4 CONTRAINDICATIONS

VIBERZI is contraindicated in patients:

- Without a gallbladder. These patients are at increased risk of developing serious adverse reactions of pancreatitis and/or sphincter of Oddi spasm [see *Warnings and Precautions (5.1, 5.2)*]
- With known or suspected biliary duct obstruction; or sphincter of Oddi disease or dysfunction. These patients are at increased risk for sphincter of Oddi spasm [see *Warnings and Precautions (5.1)*].

- With alcoholism, alcohol abuse or alcohol addiction, or in patients who drink more than 3 alcoholic beverages per day. These patients are at increased risk for acute pancreatitis [see *Warnings and Precautions (5.1)*].
- With a history of pancreatitis; or structural diseases of the pancreas, including known or suspected pancreatic duct obstruction. These patients are at increased risk for acute pancreatitis [see *Warnings and Precautions (5.1)*].
- With severe hepatic impairment (Child-Pugh Class C). These patients are at risk for significantly increased plasma concentrations of eluxadoline [see *Use in Specific Populations (8.6)*].
- With a history of chronic or severe constipation or sequelae from constipation, or known or suspected mechanical gastrointestinal obstruction. These patients may be at risk for severe complications of bowel obstruction. *Warnings and Precautions*

5 WARNINGS AND PRECAUTIONS

5.1 Pancreatitis

Pancreatitis, with or without sphincter of Oddi spasm [see *Warnings and Precautions (5.1)*], has been reported in patients taking either the 75 mg or 100 mg dosage of VIBERZI, including serious cases resulting in hospitalization, primarily in patients without a gallbladder. Fatal cases have also been reported in patients without a gallbladder. VIBERZI is contraindicated in patients without a gallbladder [see *Contraindications (4)*]. Most of the reported cases of serious pancreatitis occurred within a week of starting treatment with VIBERZI and some developed symptoms after one to two doses.

In patients with a gallbladder, evaluate a patient's alcohol intake prior to starting VIBERZI. Instruct patients to avoid chronic or acute excessive alcohol use while taking VIBERZI. Monitor for new or worsening abdominal pain that may radiate to the back or shoulder, with or without nausea and vomiting. Instruct patients to immediately stop VIBERZI and seek medical attention if they experience symptoms suggestive of pancreatitis such as acute abdominal or epigastric pain radiating to the back or shoulder associated with elevations of pancreatic enzymes with or without nausea and vomiting [see *Contraindications (4)*].

5.2 Sphincter of Oddi Spasm

There is a risk of sphincter of Oddi spasm, resulting in pancreatitis or hepatic enzyme elevation associated with acute abdominal pain (e.g., biliary-type pain) in patients taking VIBERZI [see *Adverse Reactions (6.1)*]. Postmarketing serious adverse reactions of sphincter of Oddi spasm with or without pancreatitis resulting in hospitalization have been reported, primarily in patients without a gallbladder [see *Warnings and Precautions (5.1)*]. Most of the reported cases of

serious sphincter of Oddi spasm occurred within a week of starting treatment with VIBERZI and some developed symptoms after one to two doses. VIBERZI is contraindicated in patients without a gallbladder [*see Contraindications (4)*].

Instruct patients to immediately stop VIBERZI and seek medical attention if they experience symptoms suggestive of sphincter of Oddi spasm such as acute worsening of abdominal pain, (e.g. acute epigastric or biliary [i.e., right upper quadrant] pain), that may radiate to the back or shoulder with or without nausea and vomiting, associated with elevations of pancreatic enzymes or liver transaminases. Do not restart VIBERZI in patients who developed biliary duct obstruction or sphincter of Oddi spasm while taking VIBERZI [*see Contraindications (4)*].

6 ADVERSE REACTIONS

The following adverse reactions described below and elsewhere in the labeling include:

- Pancreatitis [*see Warnings and Precautions (5.1)*]
- Sphincter of Oddi Spasm [*see Warnings and Precautions (5.2)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Over 1700 patients with IBS-D have been treated with 75 or 100 mg of VIBERZI twice daily in controlled trials. Exposures from placebo-controlled clinical trials in adult patients with IBS-D included 1391 exposed for 3 months, 1001 exposed for 6 months and 488 exposed for one year.

Demographic characteristics were comparable between the treatment groups [*see Clinical Studies (14)*]. Data described below represent pooled data compared to placebo across the randomized trials.

Pancreatitis

Cases of pancreatitis, not associated with sphincter of Oddi spasm, were reported in 2/807 (0.2%) of patients receiving 75 mg and 3/1032 (0.3%) of patients receiving 100 mg VIBERZI twice daily in clinical trials. Of these 5 cases, 3 were associated with excessive alcohol intake, one was associated with biliary sludge, and in one case the patient discontinued VIBERZI 2 weeks prior to the onset of symptoms. All pancreatic events resolved with lipase normalization upon discontinuation of VIBERZI, with 80% (4/5) resolving within 1 week of treatment discontinuation. The case of sphincter of Oddi spasm-induced pancreatitis resolved within 24 hours of discontinuation.

Sphincter of Oddi Spasm

In clinical trials, sphincter of Oddi spasm occurred in 0.2% (2/807) of patients receiving 75 mg and 0.8% (8/1032) of patients receiving 100 mg VIBERZI twice daily.

- Among patients receiving 75 mg, 1/807 (0.1%) patient experienced a sphincter of Oddi spasm presenting with abdominal pain but with lipase elevation less than 3 times the upper limit of normal (ULN) and 1/ 807 (0.1%) patient experienced a sphincter of Oddi spasm manifested as elevated hepatic enzymes associated with abdominal pain
- Among patients receiving 100 mg, 1/1032 (0.1%) patient experienced a sphincter of Oddi spasm manifested as pancreatitis and 7/1032 (0.7%) patients experienced sphincter of Oddi spasm manifested as elevated hepatic enzymes associated with abdominal pain

Of those patients who experienced a sphincter of Oddi spasm, 80% (8/10) reported their first onset of symptoms within the first week of treatment. The case of sphincter of Oddi spasm-induced pancreatitis occurred within minutes of taking the first dose of VIBERZI. No cases of sphincter of Oddi spasm occurred greater than 1 month after treatment onset. All events resolved upon discontinuation of VIBERZI, with symptoms typically improved by the following day.

Common Adverse Reactions

Table 1 provides the incidence of common adverse reactions reported in > 2% of IBS-D patients in either VIBERZI treatment group and at an incidence greater than in the placebo group.

Table 1: Common* Adverse Reactions in the Placebo-Controlled Studies in IBS-D Patients

Adverse Reactions	VIBERZI 100 mg twice daily (N= 1032) %	VIBERZI 75 mg twice daily (N=807) %	Placebo (N=975) %
Constipation	8	7	2
Nausea	7	8	5
Abdominal Pain**	7	6	4
Upper Respiratory Tract Infection	5	3	4
Vomiting	4	4	1
Nasopharyngitis	3	4	3
Abdominal Distention	3	3	2
Bronchitis	3	3	2
Dizziness	3	3	2
Flatulence	3	3	2
Rash***	3	3	2
Increased ALT	3	2	1
Fatigue	2	3	2
Viral gastroenteritis	1	3	2

* Reported in > 2% of VIBERZI-treated patients at either dose and at an incidence greater than in placebo-treated patients

** "Abdominal Pain" term includes: abdominal pain, abdominal pain lower, and abdominal pain upper

*** "Rash" term includes: dermatitis, dermatitis allergic, rash, rash erythematous, rash generalized, rash maculopapular, rash papular, rash pruritic, urticaria, and idiopathic urticaria

Constipation was the most commonly reported adverse reaction in VIBERZI-treated patients in these trials. Approximately 50% of constipation events occurred within the first 2 weeks of treatment while the majority occurred within the first 3 months of therapy. Rates of severe constipation were less than 1% in patients receiving 75 mg and 100 mg VIBERZI. Similar rates of constipation occurred between the active and placebo arms beyond 3 months of treatment.

Adverse Reactions Leading to Discontinuation

Eight percent of patients treated with 75 mg, 8% of patients treated with 100 mg VIBERZI and 4% of patients treated with placebo discontinued prematurely due to adverse reactions. In the VIBERZI treatment groups, the most common reasons for discontinuation due to adverse reactions were constipation (1% for 75 mg and 2% for 100 mg) and abdominal pain (1% for both 75 mg and 100 mg). In comparison, less than 1% of patients in the placebo group withdrew due to constipation or abdominal pain.

Less Common Adverse Reactions

Adverse reactions that were reported in ≤ 2% of VIBERZI-treated patients are listed below by body system.

Gastrointestinal: gastroesophageal reflux disease

General Disorders and administration site conditions: feeling drunk

Investigations: increased AST

Nervous system: sedation, somnolence

Psychiatric disorders: euphoric mood

Respiratory: asthma, bronchospasm, respiratory failure, wheezing

7 DRUG INTERACTIONS

The metabolism of eluxadoline by CYP pathways has not been clearly established. In addition, the potential of eluxadoline to inhibit CYP3A4 in the gut has not been established.

Tables 2 and 3 include drugs which demonstrated a clinically important drug interaction with VIBERZI or which potentially may result in clinically relevant interactions.

Table 2: Established and Other Potentially Clinically Relevant Interactions Affecting VIBERZI

OATP1B1 Inhibitors	
<i>Clinical Impact:</i>	Increased exposure to eluxadoline when coadministered with cyclosporine [see <i>Clinical Pharmacology (12.3)</i>]
<i>Intervention:</i>	Administer VIBERZI at a dose of 75 mg twice daily [see <i>Dosage and Administration (2)</i>] and monitor patients for impaired mental or physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery and for other eluxadoline-related adverse reactions [see <i>Adverse Reactions (6.1)</i>].
<i>Examples:</i>	cyclosporine, gemfibrozil, antiretrovirals (atazanavir, lopinavir, ritonavir, saquinavir, tipranavir), rifampin, eltrombopag
Strong CYP Inhibitors*	
<i>Clinical Impact:</i>	Potential for increased exposure to eluxadoline [see <i>Clinical Pharmacology (12.3)</i>]
<i>Intervention:</i>	Monitor patients for impaired mental or physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery and for other eluxadoline-related adverse reactions [see <i>Adverse Reactions (6.1)</i>].
<i>Examples:</i>	ciprofloxacin, (CYP1A2), gemfibrozil (CYP2C8), fluconazole, (CYP2C19), clarithromycin (CYP3A4), paroxetine and bupropion, (CYP2D6)
Drugs that Cause Constipation	
<i>Clinical Impact:</i>	Increased risk for constipation related adverse reactions and potential for constipation related serious adverse reactions
<i>Intervention:</i>	Avoid use with other drugs that may cause constipation (see below); loperamide may be used occasionally for acute management of severe diarrhea but avoid chronic use. Discontinue loperamide immediately if constipation occurs.
<i>Examples:</i>	alosetron, anticholinergics, opioids

*As a precautionary measure due to incomplete information on the metabolism of eluxadoline

Table 3: Established and Other Potentially Clinically Relevant Interactions Affecting Drugs Co-Administered with VIBERZI

OATP1B1 and BCRP Substrate	
<i>Clinical Impact:</i>	VIBERZI may increase the exposure of co-administered OATP1B1 and BCRP substrates. Increased exposure to rosuvastatin when co-administered with VIBERZI with a potential for increased risk of myopathy/rhabdomyolysis [see <i>Clinical Pharmacology (12.3)</i>]
<i>Intervention:</i>	Use the lowest effective dose of rosuvastatin (see prescribing information of rosuvastatin for additional information on recommended dosing).
CYP3A Substrates with Narrow Therapeutic Index	
<i>Clinical Impact:</i>	Potential for increased exposure of co-administered drug [see <i>Clinical Pharmacology (12.3)</i>]
<i>Intervention:</i>	Monitor drug concentrations or other pharmacodynamic markers of drug effect when concomitant use with eluxadoline is initiated or discontinued.
<i>Examples:</i>	alfentanil, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, tacrolimus

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no studies with VIBERZI in pregnant women that inform any drug-associated risks. The background risk of major birth defects and miscarriage for the indicated population is unknown. However, the background risk in the U.S. general population of major birth defects is 2 to 4% and of miscarriage is 15 to 20% of clinically recognized pregnancies. In animal reproduction studies, oral and subcutaneous administration of eluxadoline to rats and rabbits during organogenesis at doses approximately 51 and 115 times the human exposure after a single oral dose of 100 mg, respectively, demonstrated no teratogenic effects. In a pre- and postnatal development study in rats, no adverse effects were observed in offspring with oral administration of eluxadoline at doses approximately 10 times the human exposure [see *Data*].

Data

Animal Data

Eluxadoline administered as combined oral (1000 mg/kg/day) and subcutaneous (5 mg/kg/day) doses during the period of organogenesis to rats and rabbits (exposures about 51 and 115 times, respectively, the human AUC of 24 ng.h/mL after a single oral dose of 100 mg) did not cause any adverse effects on embryofetal development. A pre- and postnatal development study in rats showed no evidence of any adverse effect on pre- and postnatal development at oral doses of eluxadoline up to 1000 mg/kg/day (with exposures about 10 times the human AUC of 24 ng.h/mL after a single oral dose of 100 mg). In the same study, eluxadoline was detected in the

milk of lactating rats administered oral doses of 100, 300 and 1000 mg/kg/day (with exposures about 1.8, 3 and 10 times, respectively, the human AUC of 24 ng.h/mL after a single oral dose of 100 mg). Milk samples were collected from six lactating females per group on lactation day 12. Mean concentrations of eluxadoline in the milk of lactating rats on lactation day 12 were 2.78, 5.49 and 44.02 ng/mL at 100, 300 and 1000 mg/kg/day, respectively.

8.2 Lactation

Risk Summary

No data are available regarding the presence of eluxadoline in human milk, the effects of eluxadoline on the breastfed infant, or the effects of eluxadoline on milk production. However, eluxadoline is present in rat milk [*see Use in Specific Populations (8.1)*].

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for VIBERZI and any potential adverse effects on the breastfed infant from VIBERZI or from the underlying maternal condition.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

Juvenile Toxicology Data

Eluxadoline was orally administered to juvenile rats at 500, 750, and 1500 mg/kg/day (about 16, 54 and 30 times, respectively, the human AUC of 24 ng.h/mL after a single oral dose of 100 mg) for 4 weeks. There were no adverse physiologic effects related to eluxadoline. Based on these results, the NOAEL for male and female juvenile rats was 1500 mg/kg/day (about 30 times the human AUC of 24 ng.h/mL after a single oral dose of 100 mg).

8.5 Geriatric Use

Of 1795 IBS-D patients in clinical trials of VIBERZI who received 75 mg or 100 mg twice daily, 139 (7.7%) were at least 65 years of age, while 15 (0.8%) were at least 75 years old. No overall differences in effectiveness were observed between these patients and younger patients. There were no overall differences in the types of adverse reactions observed between elderly and younger patients; however, a higher proportion of elderly patients than younger patients experienced adverse reactions (66% vs 59%), serious adverse reactions (9% vs 4%), and gastrointestinal adverse reactions (39% vs 28%).

8.6 Hepatic Impairment

Plasma concentrations of eluxadoline increase in patients with hepatic impairment [*see Clinical Pharmacology (12.3)*].

VIBERZI is contraindicated in patients with severe hepatic impairment (Child-Pugh Class C) as plasma concentrations of eluxadoline increase significantly (16-fold) and there is no information to support the safety of VIBERZI in these patients.

In patients with mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment, plasma concentrations of eluxadoline increase to a lesser extent (6- and 4-fold, respectively). Administer VIBERZI at a reduced dose of 75 mg twice daily to these patients [*see Dosage and Administration (2)*]. Monitor patients with any degree of hepatic impairment for impaired mental or physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery and for other eluxadoline-related adverse reactions [*see Adverse Reactions (6.1)*].

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Eluxadoline is listed in Schedule IV of the Controlled Substances Act.

9.2 Abuse

In a drug discrimination study in monkeys, intravenous administration of eluxadoline hydrochloride produced full generalization to the morphine cue. In a self-administration study in monkeys, eluxadoline hydrochloride was self-administered to a degree that was less than that of heroin but greater than that of saline.

Adverse reactions of euphoria and feeling drunk were reported in clinical trials of IBS-D evaluating 75 mg and 100 mg doses of VIBERZI. The rate of euphoria was 0% for 75 mg and 0.2% (2/1032) for 100 mg and the rate of feeling drunk was 0.1% (1/807) for 75 mg and 0.1% (1/1032) for 100 mg.

In contrast, in two human abuse potential studies conducted in recreational opioid-experienced individuals, suprathreshold oral doses of VIBERZI (300 mg and/or 1000 mg) and intranasal doses of VIBERZI (100 mg and/or 200 mg) produced the adverse reaction of euphoria (at a rate ranging from 14% to 28%) that was greater than that of placebo (0% to 5%) but less than that of oxycodone (44% to 76%). In the two human abuse potential studies, suprathreshold oral and intranasal doses of VIBERZI produced small but significant increases on positive subjective measures such as Drug Liking and High compared to placebo. Suprathreshold oral and intranasal doses of VIBERZI also produced small but significant increases on negative subjective measures such as Drug Disliking and Dysphoria compared to placebo. In the same studies, oxycodone (30 mg and 60 mg oral, and 15 and 30 mg intranasal) produced significantly greater responses on positive and negative subjective measures than those produced by eluxadoline and placebo.

microcrystalline cellulose, colloidal silica, crospovidone, mannitol, magnesium stearate, and Opadry II (partially hydrolyzed polyvinyl alcohol, titanium dioxide, polyethylene glycol, talc, iron oxide yellow, and iron oxide red).

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Eluxadoline is a mu-opioid receptor agonist; eluxadoline is also a delta opioid receptor antagonist and a kappa opioid receptor agonist. The binding affinities (K_i) of eluxadoline for the human mu and delta opioid receptors are 1.8 nM and 430 nM, respectively. The binding affinity (K_i) of eluxadoline for the human kappa opioid receptor has not been determined; however, the K_i for guinea pig cerebellum kappa opioid receptor is 55 nM. In animals, eluxadoline interacts with opioid receptors in the gut.

12.2 Pharmacodynamics

Cardiac Electrophysiology

At a dose 10 times the maximum recommended dose (100 mg), VIBERZI does not prolong the QT interval to any clinically relevant extent.

12.3 Pharmacokinetics

Following oral administration of 100 mg VIBERZI in healthy subjects, the C_{max} of eluxadoline was approximately 2 to 4 ng/mL and AUC was 12 to 22 ng.h/mL. Eluxadoline has approximately linear pharmacokinetics with no accumulation upon repeated twice daily dosing. The variability of eluxadoline pharmacokinetic parameters ranges from 51% to 98%.

Absorption

Absolute bioavailability of eluxadoline has not been determined. The median T_{max} value was 1.5 hours (range: 1 to 8 hours) under fed conditions and 2 hours (range: 0.5 to 6 hours) under fasting conditions.

The administration of VIBERZI with a high fat meal that contained approximately 800 to 1000 total calories, with 50% of calories being derived from fat content decreased the C_{max} of eluxadoline by 50% and AUC by 60%.

Distribution

Plasma protein binding of eluxadoline was 81%.

Elimination

The mean plasma elimination half-life of eluxadoline ranged from 3.7 hours to 6 hours.

Metabolism

Metabolism of eluxadoline is not clearly established [see *Drug Interactions (7)*]. There is evidence that glucuronidation can occur to form an acyl glucuronide metabolite.

Excretion

Following a single oral dose of 300 mg [¹⁴C] eluxadoline in healthy male subjects, 82.2% of the total radioactivity was recovered in feces within 336 hours and less than 1% was recovered in urine within 192 hours.

Specific Populations

Hepatic Impairment

Following a single oral 100-mg dose in subjects with varying degrees of liver impairment and healthy subjects, mean eluxadoline plasma exposure was 6-fold, 4-fold, and 16-fold higher in mild, moderate, and severe hepatically impaired subjects (Child Pugh Class A, B, C), respectively, compared to the subjects with normal liver function [see *Dosage and Administration (2)*, *Contraindications (4)*, *Use in Specific Populations (8.6)*].

Drug Interactions

In Vitro Assessment of Drug Interactions

In vitro studies indicate that eluxadoline is neither an inducer of CYP1A2, CYP2C9, CYP2C19, and CYP3A4, nor an inhibitor of CYP1A2, CYP2A6, CYP2B6, CYP2C9, CYP2C19, CYP3A4 and CYP2D6 at clinically relevant systemic concentrations. Although CYP2E1 was slightly inhibited by eluxadoline (IC₅₀ of approximately 20 micromolar [11 mcg/mL]), clinically meaningful interactions are unlikely. The *in vitro* studies were not adequate to establish the potential for eluxadoline to inhibit CYP3A4 in the gut [see *Drug Interactions (7)*].

In vitro studies suggest that eluxadoline is a substrate for OAT3, OATP1B1, BSEP and MRP2, but not for OCT1, OCT2, OAT1, OATP1B3, P-gp and BCRP. Based on the *in vitro* studies, clinically meaningful interaction via inhibition of OCT1, OCT2, OAT1, OAT3, OATP1B3, BSEP and MRP2 by eluxadoline is unlikely. However, the *in vitro* studies were not adequate to establish the potential for eluxadoline to inhibit P-gp in the gut.

In Vivo Assessment of Drug Interactions

The following drug interactions were studied in healthy subjects:

Oral Contraceptives

Coadministration of multiple doses of 100 mg VIBERZI with multiple dose administration of an oral contraceptive (norethindrone 0.5 mg/ethinyl estradiol 0.035 mg) does not change the exposure of either drug.

Cyclosporine

Coadministration of a single dose of 100 mg VIBERZI with a single dose of 600 mg cyclosporine resulted in 4.4-fold and 6.2-fold increase in AUC and C_{max} of eluxadoline, respectively, compared to administration of VIBERZI alone [see *Drug Interactions (7)*].

Probenecid

Coadministration of a single dose of 100 mg VIBERZI with a single dose of 500 mg probenecid resulted in a 35% and 31% increase in eluxadoline AUC and C_{max} , respectively, compared to administration of VIBERZI alone. This change in eluxadoline exposures is not expected to be clinically meaningful.

Rosuvastatin

Coadministration of multiple doses of 100 mg VIBERZI twice daily with a single dose 20 mg rosuvastatin resulted in an increase in the AUC (40%) and C_{max} (18%) of rosuvastatin compared to administration of rosuvastatin alone. Similar results were observed with the active, major metabolite, n-desmethyl rosuvastatin [see *Drug Interactions (7)*].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Two-year oral carcinogenicity studies have been conducted with eluxadoline in CD-1 mice at doses up to 1500 mg/kg/day (about 14 times the human AUC of 24 ng.h/mL after a single oral dose of 100 mg) and in Sprague Dawley rats at oral doses up to 1500 mg/kg/day (about 36 times the human AUC of 24 ng.h/mL after a single oral dose of 100 mg). Oral administration of eluxadoline for 104 weeks did not produce tumors in mice and rats.

Mutagenesis

Eluxadoline was negative in the Ames test, chromosome aberration test in human lymphocytes, in the mouse lymphoma cell (L5178Y/TK^{+/+}) forward mutation test and in the *in vivo* rat bone marrow micronucleus test.

Impairment of Fertility

Eluxadoline at oral doses up to 1000 mg/kg/day (about 10 times the human AUC of 24 ng.h/mL after a single oral dose of 100 mg) was found to have no adverse effect on fertility and reproductive performance of male and female rats.

14 CLINICAL STUDIES

The efficacy and safety of VIBERZI in IBS-D patients was established in two randomized, multi-center, multi-national, double-blind, placebo-controlled trials (Studies 1 and 2). A total of 1281 patients in Study 1 and 1145 patients in Study 2 received treatment with VIBERZI 75 mg, VIBERZI 100 mg or placebo twice daily [overall, patients had a mean age of 45 years (range 18 to 80 years with 10% at least 65 years of age or older), 66% female, 86% white, 11% black, and 27% Hispanic].

All patients met Rome III criteria for IBS-D (loose [mushy] or watery stools $\geq 25\%$ and hard or lumpy stools $< 25\%$ of bowel movements) and were required to meet both of the following criteria:

- an average of worst abdominal pain scores in the past 24 hours of > 3.0 on a 0 to 10 scale over the week prior to randomization.
- an average daily stool consistency score (Bristol Stool Scale or BSS) of ≥ 5.5 and at least 5 days with a BSS score ≥ 5 on a 1 to 7 scale over the week prior to randomization.

Pertinent exclusion criteria included: prior pancreatitis, alcohol abuse, cholecystitis prior 6 months, sphincter of Oddi dysfunction, inflammatory bowel disease, intestinal obstruction, gastrointestinal infection or diverticulitis within prior 3 months, lipase greater than 2 xULN, ALT or AST greater than 3 xULN.

Study 1 and Study 2 included identical 26-week double-blind, placebo-controlled treatment periods. Study 1 continued double-blinded for an additional 26 weeks for long-term safety (total of 52 weeks of treatment), followed by a 2-week follow-up. Study 2 included a 4-week single-blinded, placebo-withdrawal period upon completion of the 26-week treatment period. During the double-blind treatment phase and the single-blinded placebo withdrawal phase, patients were allowed to take loperamide rescue medication for the acute treatment of uncontrolled diarrhea, but were not allowed to take any other antidiarrheal, antispasmodic agent or rifaximin for their diarrhea. Additionally, patients were allowed to take aspirin-containing medications or nonsteroidal anti-inflammatory drugs, but no narcotic or opioid containing agents.

Efficacy of VIBERZI was assessed in both trials using an overall composite responder primary endpoint. The primary endpoint was defined by the simultaneous improvement in the daily worst abdominal pain score by $\geq 30\%$ as compared to the baseline weekly average AND a reduction in the BSS to < 5 on at least 50% of the days within a 12-week time interval. Improvement in daily worst abdominal pain in the absence of a concurrent bowel movement was also considered a response day. Results for endpoints were based on electronic daily diary entries by patients.

The proportion of composite responders over 12 weeks is shown in **Table 4**. In both trials, the proportion of patients who were composite responders to VIBERZI was statistically significantly

higher than placebo for both doses. The proportion of patients who were composite responders to VIBERZI was similar for male and female patients in both trials.

Table 4: Efficacy Results in Randomized Clinical Trials

	Study 1			Study 2		
	VIBERZI 100mg twice daily n=426	VIBERZI 75mg twice daily n=427	PBO n=427	VIBERZI 100mg twice daily n=382	VIBERZI 75mg twice daily n=381	PBO n=382
Composite¹ Response over 12 weeks						
Responder rates	25%	24%	17%	30%	29%	16%
Treatment difference	8% ²	7% ⁴		13% ³	13% ³	
95% CI (%)	(2.6, 13.5)	(1.4, 12.2)		(7.5, 19.2)	(6.8, 18.5)	
Composite Response over 26 weeks						
Responder rates	29%	23%	19%	33%	30%	20%
Treatment difference	10%	4%		13%	10%	
95% CI (%)	(4.7, 16.1)	(-1.0, 9.9)		(6.4, 18.8)	(4.2, 16.4)	
Abdominal Pain Response Improved ≥30% over 12 weeks						
Responder rates	43%	42%	40%	51%	48%	45%
Treatment difference	4%	3%		6%	3%	
95% CI (%)	(-3.0, 10.2)	(-3.8, 9.4)		(-1.3, 12.8)	(-4.3, 9.8)	
BSS <5 Response over 12 weeks						
Responder rates	34%	30%	22%	36%	37%	21%
Treatment difference	12%	8%		15%	16%	
95% CI (%)	(6.3, 18.2)	(2.1, 13.8)		(8.4, 21.0)	(9.7, 22.4)	

¹Composite= Simultaneous improvement of Worst Abdominal Pain (WAP) by ≥30% and Bristol Stool Score (BSS) < 5 on the same day for ≥ 50% of days over the interval

² P<0.01

³ P<0.001

⁴ P<0.05

Additionally, the proportion of patients who were composite responders to VIBERZI at each 4-week interval was numerically higher than placebo for both doses as early as month 1 through month 6 demonstrating that efficacy is maintained throughout the course of treatment.

During the 4 week single-blind withdrawal period in Study 2, no evidence of worsening of diarrhea or abdominal pain compared to baseline was demonstrated at either dose.

16 HOW SUPPLIED/STORAGE AND HANDLING

VIBERZI is available as:

- 75 mg tablets: capsule-shaped tablets, coated in pale-yellow to light tan color, debossed with “FX75” on one side.
Bottle of 60: NDC 61874-075-60
- 100 mg tablets: capsule-shaped tablets, coated in pink-orange to peach color, debossed with “FX100” on one side.
Bottle of 60: NDC 61874-100-60

Store VIBERZI tablets at 20°C to 25°C (68°F to 77°F) with excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

Instruct patients to:

- stop VIBERZI immediately and seek medical attention if unusual or severe abdominal pain that may radiate to the back or shoulder with or without nausea and vomiting, develops [see *Warnings and Precautions (5.1, 5.2)*].
- avoid chronic or acute excessive alcohol use while taking VIBERZI [see *Warnings and Precautions (5.1)*].
- take one tablet twice daily with food.
- if they miss a dose, take the next dose at the regular time. Do not take 2 doses at the same time to make up for a missed dose.
- call their healthcare provider if they are unable to tolerate VIBERZI
- discontinue VIBERZI and call their health care provider if they experience severe constipation
- not take alosetron with VIBERZI or not take loperamide on a *chronic* basis with VIBERZI due to the potential for constipation. Loperamide may occasionally be used with VIBERZI for *acute management* of severe diarrhea, but must be discontinued if

constipation develops. Also, instruct patients to avoid taking VIBERZI with other medications that may cause constipation (for example opioids, anticholinergics, etc.).

Distributed by:
Allergan USA, Inc.
Irvine, CA 92612

© 2017 Allergan. All rights reserved.

MEDICATION GUIDE
VIBERZI (vye BER zee), CIV
(eluxadoline) tablets

What is the most important information I should know about VIBERZI?

VIBERZI can cause serious side effects, including:

- **Inflammation of the pancreas (pancreatitis).** Pancreatitis has happened most often in people who do not have a gallbladder and can lead to hospitalization. Pancreatitis has led to death in some people who do not have a gallbladder. Pancreatitis usually happens within the first week of treatment with VIBERZI but can happen after 1 to 2 doses of VIBERZI. Your risk of getting pancreatitis is increased if you drink more than 3 alcoholic drinks a day. Limit your use of alcoholic drinks while you are taking VIBERZI.
- **Sphincter of Oddi spasm.** The sphincter of Oddi is a muscular valve that controls the flow of digestive juices (bile and pancreatic juice) to the first part of your small intestine. A sphincter of Oddi spasm can cause an increase in your liver and pancreas enzymes and inflammation of the pancreas (pancreatitis) that can cause sudden stomach-area (abdomen) pain. Sphincter of Oddi spasm has happened most often in people who do not have a gallbladder and can lead to hospitalization. This spasm usually happens within the first week of treatment with VIBERZI but can happen after 1 or 2 doses of VIBERZI.

Stop taking VIBERZI right away and get emergency medical care if you have new or worsening stomach-area (abdomen) pain or pain in the upper right side of your stomach-area (abdomen) that may move to your back or shoulder, with or without nausea and vomiting.

What is VIBERZI?

VIBERZI is a prescription medicine used to treat adults who have irritable bowel syndrome with diarrhea (IBS-D).

- VIBERZI is a controlled substance (CIV) because it contains eluxadoline and may be abused or lead to drug dependence. Keep your VIBERZI in a safe place, to protect it from theft. Never give your VIBERZI to anyone else, because it may harm them. Selling or giving away this medicine is against the law.

It is not known if VIBERZI is safe and effective in children.

People 65 years old and older have had an increased number of side effects, including serious side effects and stomach problems, while taking VIBERZI than people younger than 65 years old have had.

Do not take VIBERZI if you:

- do not have a gallbladder
- have or may have had a blockage in your gallbladder or a sphincter of Oddi problem
- have or had problems with alcohol abuse, alcohol addiction, or drink more than 3 alcoholic drinks a day
- have had inflammation of your pancreas (pancreatitis) or other pancreas problems, including if you have had or may have had a blockage in your pancreas
- have severe liver problems
- have had long-lasting (chronic) or severe constipation, or problems caused by constipation
- have or may have had a bowel blockage (intestinal obstruction)

Talk to your doctor if you are not sure if you have any of these conditions.

Before taking VIBERZI, tell your doctor about all of your medical conditions, including if you:

- See “**What is the most important information I should know about VIBERZI?**”
- have liver problems
- are pregnant or plan to become pregnant. It is not known if VIBERZI will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if VIBERZI passes into your breast milk or could harm your baby.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of your medicines to show your doctor and pharmacist when you get a new medicine. VIBERZI and other medicines may affect each other causing side effects.

If you are taking VIBERZI you should not take:

- medicines that cause constipation including:
 - Lotronex[®] (alosetron)
 - anticholinergic medicines
 - opioid pain medicines

Ask your doctor or pharmacist for a list of these medicines, if you are not sure.

- Avoid taking loperamide, a medicine used to treat diarrhea, for a long time (chronic use). You may take loperamide occasionally to treat severe diarrhea. **Stop taking loperamide right away if you become constipated.**

How should I take VIBERZI?

- Take VIBERZI exactly as your doctor tells you to take it.
- Take 1 tablet of VIBERZI 2 times each day with food.
- If you miss a dose, take your next dose at your regular time. Do not take 2 doses at the same time to make up for a missed dose.
- Do not change your dose or stop taking VIBERZI unless your doctor tells you to.
- If you take too much VIBERZI, call your doctor or go to the nearest hospital emergency room right away.

What should I avoid while taking VIBERZI?

- Limit your use of alcoholic drinks while you are taking VIBERZI.
- If you have liver problems, **do not** drive, operate machinery, or do other dangerous activities until you know how VIBERZI affects you.

What are the possible side effects of VIBERZI?

See “**What is the most important information I should know about VIBERZI?**”

The most common side effects of VIBERZI include: constipation, nausea, and abdominal pain. Stop taking VIBERZI and call your doctor if you have constipation that is severe.

These are not all the possible side effects of VIBERZI. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store VIBERZI?

Store VIBERZI at room temperature between 68°F to 77°F (20°C to 25°C).

Keep VIBERZI and all medicines out of the reach of children.

General Information about the safe and effective use of VIBERZI

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use VIBERZI for a condition for which it was not prescribed. Do not give VIBERZI to other people, even if they have the same symptoms that you have. It may harm them. You can ask your doctor or pharmacist for information about VIBERZI that is written for health professionals.

What are the ingredients in VIBERZI?

Active ingredient: eluxadoline

Inactive ingredients: silicified microcrystalline cellulose, colloidal silica, crospovidone, mannitol, magnesium stearate, and Opadry II (partially hydrolyzed polyvinyl alcohol, titanium dioxide, polyethylene glycol, talc, iron oxide yellow, and iron oxide red).

Distributed by: Actavis Pharma, Inc., Parsippany, NJ 07054 USA

© 2017 Allergan. All rights reserved.

For more information, go to www.VIBERZI.com or call 1-800-272-5525.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: April 2017