

PRODUCTION (NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	5.5 pt
• SUBHEADINGS/BODY TEXT	5.5 pt
• LEADING	5.5 pt
• BULLETS	4.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	1 pt, .5 pt

090177e18963b1afFinalFinal On: 11-Aug-2016 19:49




DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend

090177e18963b1aa\Final\Final On: 11-Aug-2016 19:50



090177e18963b1b2\Final\Final On: 11-Aug-2016 19:49

PRODUCTION (NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	N/A
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces


**TOP (FACE)
 PANEL 1**



UNVARNISHED AND PRINTED IN BLACK FOR LOT & EXPIRATION AREA
 ART BLEEDS OFF 1/16" (0.0625") OUT FROM DIE (DASH LINE DOES NOT PRINT)
 0.125" X 0.125" DATA MATRIX CODE (ECC 200), PRINTED IN BLACK ON UNPRINTED WHITE BACKGROUND
DATA MATRIX CODE = 078631

**BASE
 PANEL 3**


1/4" HINGE AREA UNVARNISHED (LINES DO NOT PRINT)



ART BLEEDS OFF 1/16" (0.0625") OUT FROM DIE (DASH LINE DOES NOT PRINT)
 THIS LINE INDICATES DIE AND DOES NOT PRINT
 NO COPY 1/16" (0.0625") IN FROM DIE (DASH LINE DOES NOT PRINT)

(b) (4)

**TOP (BACK)
 PANEL 2**



THIS LINE INDICATES DIE AND DOES NOT PRINT
 (b) (4)

1-1/16" (0.0625") CORNER RADIUS TYP

PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	



DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	5.5 pt
• SUBHEADINGS/BODY TEXT	5.5 pt
• LEADING	5.5 pt
• BULLETS	4.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	1 pt, .5 pt

090177e18963a7f0\Final\Final On: 11-Aug-2016 19:47



090177e18963a7ee\Final\Final On: 11-Aug-2016 19:48

PRODUCTION (NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



090177e18963a7f2\Final\Final On: 11-Aug-2016 19:47

128 CODE AREA - SUBSET C 10 MIL DENSITY WITH 4 OR 6 DIGITS (CODE TO EXTEND TO EDGE OF FLAP - UNVARNISHED) PRINTED SIDE OF CARTON
128 CODE = 078786

128 CODE = 078786

LOT & EXP AREA. DARK COLOR FOR LASER PRINTING.

3/16" DATA MATRIX CODE WITH 1/8" QUIET ZONE ON ALL SIDES
DATA MATRIX CODE = 078786

128 CODE = 078786

PRODUCTION (NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



UNCOATED
NO COPY
NO PRINT
128 CODE = 078391

128 CODE
AREA

128 CODE = 078391

128 CODE
AREA

128 CODE = 078391

128 CODE
AREA

MICRO PRINT
LOCATION

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts
Active ingredient (in each tablet)
Esomeprazole 20 mg
(Each delayed-release tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Purpose
Acid reducer

Uses
■ treats frequent heartburn (occurs 2 or more days a week)
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have:
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain
These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have:
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain
Ask a doctor or pharmacist before use if you are taking:
■ warfarin, clopidogrel or clostazolol (blood-thinning medicines)
■ prescription antifungal or anti-yeast medicines
■ digoxin (heart medicine)
■ diazepam (anxiety medicine)
■ tacrolimus or mycophenolate mofetil (immune system medicines)
■ prescription antiretrovirals (medicines for HIV infection)
■ methotrexate (arthritis medicine)
Stop use and ask a doctor if:
■ your heartburn continues or worsens
■ you need to take this product for more than 14 days
■ you need to take more than 1 course of treatment every 4 months
■ you get diarrhea

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ adults 18 years of age and older
■ this product is to be used once a day (every 24 hours), every day for 14 days
■ may take 1 to 4 days for full effect
14-Day Course of Treatment
■ swallow 1 tablet with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 tablet a day
■ swallow whole. Do not crush or chew tablets.
■ do not use for more than 14 days unless directed by your doctor
Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information
■ read the directions and warnings before use
■ keep the carton. It contains important information.
■ store at 20-25°C (68-77°F)

Inactive ingredients
corn starch, croscopolone, D&C red no. 27 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, mica, microcrystalline cellulose, paraffin, polyethylene glycol, polyorbate 80, sodium stearyl fumarate, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600

Tips for Managing Heartburn
• Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
• Eat slowly and do not eat big meals.
• Do not eat late at night or just before bedtime.
• Do not lie flat or bend over soon after eating.
• Raise the head of your bed.
• Wear loose-fitting clothing around your stomach.
• If you are overweight, lose weight.
• If you smoke, quit smoking.

Nexium 24HR
esomeprazole magnesium
delayed-release tablets
20 mg/acid reducer

Nexium 24HR
Clinically Proven to Treat Frequent Heartburn

Treats Frequent Heartburn

Nexium 24HR
esomeprazole magnesium
delayed-release tablets
20 mg/acid reducer

42 TABLETS
Three 14-day courses of treatment

May take 1 to 4 days for full effect

3 BOTTLES
INSIDE



Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

128 CODE AREA

128 CODE AREA - SUBSET C 10 MIL DENSITY WITH 4 OR 6 DIGITS (CODE TO EXTEND TO EDGE OF FLAP - UNVARNISHED) PRINTED SIDE OF CARTON
128 CODE = 078391

128 CODE AREA

MICRO PRINT LOCATION
128 CODE = 078391

LOT & EXP AREA. DARK COLOR FOR LASER PRINTING.

3/16" DATA MATRIX CODE WITH 1/8" QUIET ZONE ON ALL SIDES
DATA MATRIX CODE = 078391

PAA078391.FDA01
FPO
3 0573 2451 42 8

128 CODE AREA

128 CODE = 078391

PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

090177e18963a7f4\Final\Final On: 11-Aug-2016 19:46

NDC 0573-2451-43

Treats Frequent Heartburn



Nexium[®]
esomeprazole magnesium
delayed-release tablets **24HR**
20 mg/acid reducer

May take 1 to 4 days for full effect

3 Pack

Tablets



42 TABLETS

Three 14-day courses
of treatment

REMOVE VARNISH
IN THIS AREA ONLY

WHITE HOLDOUT
QUIET AREA

DATA MATRIX CODE
055414

REMOVE VARNISH
WHITE HOLDOUT
LOT/EXP

Marketed by: Pfizer, Madison, NJ 07940 USA
© 2016 Pfizer Inc.
Made in France
For most recent product information, visit
www.Nexium24HR.com
Nexium is a registered trademark of AstraZeneca
Afl and is used under license.



3 0573 2451 43 5

KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts (continued)

Purpose
Esomeprazole 20 mg (each delayed-release tablet corresponds to 22.3 mg Acid reducer)

Active ingredient (in each tablet)
esomeprazole magnesium trihydrate

Uses
● treats frequent heartburn (occurs 2 or more days a week)
● not intended for immediate relief of heartburn, this drug may take 1 to 4 days for full effect

Warnings
● Do not use if you are allergic to esomeprazole
● trouble or pain swallowing food, vomiting with blood, or bloody or black stools
● heartburn with **lightheadedness, sweating or dizziness**
● chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
● frequent **chest pain**
● frequent **diarrhea**
● frequent **weight loss**
● frequent wheezing, particularly with heartburn

Ask a doctor before use if you have
● had heartburn over 3 months. This may be a sign of a more serious condition.
● frequent wheezing, particularly with heartburn

Ask a doctor or pharmacist before use if you are taking
● warfarin, clopidogrel or clostazol (blood-thinning medicines)
● digoxin (heart medicine)
● prescription antifungal or anti-yeast medicines
● prescription antiretroviral (medicines for HIV infection)
● tacrolimus or mycophenolate mofetil (immune system medicines)
● diazepam (anxiety medicine)
● prescription antibiotic (medicines for HIV infection)
● methicarbonyl (antibiotic medicine)

Stop use and ask a doctor if
● your heartburn continues or worsens
● you need to take this product for more than 14 days
● you need to take more than 1 course of treatment every 4 months
● you get diarrhea

Directions
● adults 18 years of age and older
● this product is to be used once a day (every 24 hours), every day for 14 days
● may take 1 to 4 days for full effect

14-Day Course of Treatment
● swallow 1 tablet with a glass of water before eating in the morning
● take every day for 14 days
● do not take more than 1 tablet a day
● swallow whole. Do not crush or chew tablets.
● do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
● you may repeat a 14-day course every 4 months
● do not use for more than 14 days unless directed by your doctor

Other Information
● read the directions and warnings before use
● keep the card. It contains important information.
● store at 20-25°C (68-77°F)

Inactive ingredients
com starch, croscollon, D&C red no. 27 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, mica, microcrystalline cellulose, paraffin, polyethylene glycol, poly sorbate 80, sodium stearoyl fumarate, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments?
call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.



Nexium[®]
esomeprazole magnesium
delayed-release tablets **24HR**
20 mg/acid reducer

PRODUCTION (NOT COLOR ACCURATE)

Job Information	Color Legend

DRUG FACTS TEXT DEFINED	TYPE SIZE
● DRUG FACTS TITLE	10 pt
● DRUG FACTS CONTINUED	9 pt
● HEADINGS	9 pt
● SUBHEADINGS/BODY TEXT	7 pt
● LEADING	7.5 pt
● # OF CHARACTERS PER INCH	<39
● BULLETS	6 pt
● SPACE BEFORE BULLET	2 ems
● BARLINES, HAIRLINES	1.5 pt, .5 pt
● SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

CODE 128,
 10 MIL DENSITY
CODE 128 = 079072

3/16" x 3/16" Data Matrix
 Code ECC 200, printed in
 black on unprinted white
 background (quiet zone
 of 1 mm). Minimum 2 mm
 from edge of insert.

DATA MATRIX CODE = 079072

UNPRINTED AREA
 (TYP.)

UNPRINTED AREA
 (TYP.)

FRONT

CODE 128,
 10 MIL DENSITY
CODE 128 = 079072

PRODUCTION

(NOT COLOR ACCURATE)

Job Information	Color Legend
(b) (4)	

090177e189741bab\Final\Final On: 18-Aug-2016 15:38