



Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain ■ you are pregnant or breast-feeding, ask a health professional before use. ■ you develop a rash or joint pain

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions: ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect. **14-Day Course of Treatment** ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ swallow whole. Do not crush or chew tablets. ■ do not use for more than 14 days unless directed by your doctor. **Repeated 14-Day Courses (if needed)** ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F)

Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-800-228-1800

Active ingredient (in each tablet) **Purpose(s)**
 Esomeprazole 20 mg Acid reducer

(Each delayed-release tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Use ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If an allergic reaction occurs, stop use and seek medical help right away.
 Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain. These may be signs of a serious condition. See your doctor.
 Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Esomeprazole 20 mg. Acid reducer (Each delayed-release tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p>	<p>Drug Facts (continued)</p> <p>if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect</p> <p>14-Day Course of Treatment ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ swallow whole. Do not crush or chew tablets. ■ do not use for more than 14 days unless directed by your doctor</p> <p>Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other Information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F)</p> <p>Inactive ingredients corn starch, crospovidone, D&C red no. 27 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, mica, microcrystalline cellulose, paraffin, polyethylene glycol, polysorbate 80, sodium stearyl fumarate, sucrose, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600</p>
---	---



Nexium[®] 24HR
esomeprazole magnesium delayed-release tablets
20 mg/acid reducer

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
©2022 GSK group of companies or its licensor
Made in France
For most recent product information, visit www.Nexium24HR.com
Nexium is a registered trademark of AstraZeneca AB and is used under license.



000078388
3 0573 2451 14 5

NDC 0573-2451-14

See new warning information

Treats Frequent Heartburn



Nexium[®] 24HR
esomeprazole magnesium delayed-release tablets
20 mg/acid reducer

Tablets

May take 1 to 4 days for full effect

14 TABLETS
One 14-day course of treatment.

1 BOTTLE INSIDE



Treats Frequent Heartburn

Nexium[®] 24HR
esomeprazole magnesium delayed-release tablets
20 mg/acid reducer

14 Tablets Tablets
One 14-day course of treatment

ACTUAL SIZE

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Reference ID: 5041197

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Esomeprazole 20 mg Acid reducer (Each delayed-release tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are: ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p>	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect</p> <p>14-Day Course of Treatment ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ swallow whole. Do not crush or chew tablets. ■ do not use for more than 14 days unless directed by your doctor</p> <p>Repeated 14-Day Courses (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>Other Information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F)</p> <p>Inactive ingredients corn starch, croscopolone, D&C red no. 27 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, glyceryl monostearate, hydroxypropyl cellulose, hydroxymethylcellulose, magnesium stearate, methacrylic acid copolymer, mica, microcrystalline cellulose, paraffin, polyethylene glycol, polyorbate 80, sodium stearyl fumarate, sucrose, talc, titanium dioxide, triethyl citrate</p> <p>Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600</p>
---	---

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Nexium[®] 24HR
esomeprazole magnesium delayed-release tablets 20 mg/acid reducer

42 TABLETS

Three 14-day courses of treatment

May take 1 to 4 days for full effect

3 BOTTLES INSIDE



Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
09/06/2022 11:10:33 AM