

Drug Facts (continued)

- discard each cloth after a single use
- after package has been opened, discard any unused cloths

Other information

- store product flat
- store between 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

benzalkonium chloride, dimethicone emulsion, glycerin, isopropyl alcohol, propylene glycol, purified water USP



DO NOT FLUSH

Non-Sterile.

Not made with Natural Rubber Latex.

Cloth: 100% Polyester

REF MSC099CHG



www.medline.com

©2022 Medline Industries, LP
Medline is a registered trademark of Medline Industries, LP.

Manufactured by Medline Industries, LP, Three Lakes Drive, Northfield, IL 60093 USA.

Made in USA with domestic and foreign materials.


Patent Pending

Questions, comments, or ordering call: 1-800-MEDLINE

Customer Service Hours:
(Monday to Friday, 7 AM to 7 PM CST)
V1 RJ22MSC



Tear Here


NDC: 53329-244-03

READYPREP CHG[®]

**2% CHLORHEXIDINE
GLUCONATE* CLOTH**

**PATIENT PREOPERATIVE
SKIN PREPARATION**

NON-STERILE

*EQUIVALENT TO 500 MG CHLORHEXIDINE
GLUCONATE PER CLOTH


- ⊕ Provides rapid bactericidal action against a broad spectrum of microorganisms
- ⊕ Significantly reduces the number of microorganisms on intact skin
- ⊕ Demonstrates continued antimicrobial activity for up to 6 hours after application

FRAGRANCE FREE

RINSE FREE

SINGLE USE ONLY.
FOR EXTERNAL USE ONLY.
DO NOT MICROWAVE.

**DO NOT
FLUSH**



**6 9x10.5 in (22.9x26.7 cm)
DISPOSABLE CLOTHS**

REF MSC099CHG

Drug Facts

Active ingredient	Purpose
Chlorhexidine gluconate 2% solution...	Antiseptic

Uses

- helps reduce bacteria that can potentially cause skin infection
- for preparation of skin prior to surgery

Warnings

For external use only

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar punctures or in contact with the meninges
- on open skin wounds
- as a general skin cleanser

When using this product

- keep out of eyes, ears and mouth. May cause serious or permanent injury if chlorhexidine is permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.





Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.


Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- do not microwave
- product and packaging are not sterile. Follow your hospital policy for skin preparation with non-sterile products.
- **to open package:** identify the tear notch labeled on the front of the package. Grasp with hands on both sides of the tear notch and tear to expose cloth.
- transfer contents onto prep table, avoiding contact between cloth and outside of package to reduce risk of cloth contamination
- use a cloth to prepare the skin area indicated for a moist or dry site, making certain to keep the remaining cloths where they will not be contaminated. Use additional cloths to prepare larger areas.
- **dry surgical sites** (such as abdomen or arm): use one cloth to cleanse each 161 cm² area (approximately 5 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to dry for one (1) minute. Do not rinse.
- **moist surgical sites** (such as inguinal fold): use one cloth to cleanse each 65 cm² area (approximately 2 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to dry for one (1) minute. Do not rinse.



 REF MSC099CHG


READYPREP CHG®

2% CHLORHEXIDINE GLUCONATE* CLOTH PATIENT PREOPERATIVE SKIN PREPARATION


NON-STERILE
*EQUIVALENT TO 500 MG CHLORHEXIDINE GLUCONATE PER CLOTH
9×10.5 in (22.9×26.7 cm) Disposable Cloths
Qty: 6 Cloths per Pack/32 Packs per Case

Non-Sterile, Single use only. Store between 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).
Not made with Natural Rubber Latex. Store product flat.

www.medline.com Manufactured in the USA with domestic and foreign materials by Medline Industries, LP, Three Lakes Drive, Northfield, IL 60093 USA. Patent Pending. © 2009 MEDLINE. V181229C.



(01) 10193489189872

 REF MSC099CHG


READYPREP CHG®

2% CHLORHEXIDINE GLUCONATE* CLOTH PATIENT PREOPERATIVE SKIN PREPARATION

NON-STERILE
*EQUIVALENT TO 500 MG CHLORHEXIDINE GLUCONATE PER CLOTH
9×10.5 in (22.9×26.7 cm) Disposable Cloths
Qty: 6 Cloths per Pack/32 Packs per Case

Non-Sterile, Single use only. Store between 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).
Not made with Natural Rubber Latex. Store product flat.

www.medline.com Manufactured in the USA with domestic and foreign materials by Medline Industries, LP, Three Lakes Drive, Northfield, IL 60093 USA. Patent Pending. © 2009 MEDLINE. V181229C.



(01) 10193489189872

 REF MSC099CHG

READYPREP CHG®

2% CHLORHEXIDINE GLUCONATE* CLOTH PATIENT PREOPERATIVE SKIN PREPARATION


NON-STERILE
*EQUIVALENT TO 500 MG CHLORHEXIDINE GLUCONATE PER CLOTH
9×10.5 in (22.9×26.7 cm) Disposable Cloths
Qty: 6 Cloths per Pack/32 Packs per Case

Non-Sterile, Single use only. Store between 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).
Not made with Natural Rubber Latex. Store product flat.

www.medline.com Manufactured in the USA with domestic and foreign materials by Medline Industries, LP, Three Lakes Drive, Northfield, IL 60093 USA. Patent Pending. © 2009 MEDLINE. V181229C.



(01) 10193489189872

 REF MSC099CHG


READYPREP CHG®

2% CHLORHEXIDINE GLUCONATE* CLOTH PATIENT PREOPERATIVE SKIN PREPARATION





NON-STERILE
*EQUIVALENT TO 500 MG CHLORHEXIDINE GLUCONATE PER CLOTH
9×10.5 in (22.9×26.7 cm) Disposable Cloths
Qty: 6 Cloths per Pack/32 Packs per Case

Non-Sterile, Single use only. Store between 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).
Not made with Natural Rubber Latex. Store product flat.

www.medline.com Manufactured in the USA with domestic and foreign materials by Medline Industries, LP, Three Lakes Drive, Northfield, IL 60093 USA. Patent Pending. © 2009 MEDLINE. V181229C.



(01) 10193489189872



This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

PAMELA J HORN
05/24/2023 02:12:17 PM