

8U174 00 V6 100422

**Drug Facts** (continued)

**Directions** ■ adults 18 years of age and older  
■ this product is to be used once a day (every 24 hours), every day for 14 days  
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**  
■ take 1 tablet before eating in the morning

**Do not crush or chew tablets**  
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.  
■ take every day for 14 days ■ do not take more than 1 tablet a day  
■ do not use for more than 14 days unless directed by your doctor  
■ do not take this medicine with alcohol

**Repeated 14-Day Course (if needed)**  
■ you may repeat a 14-day course every 4 months  
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other information** ■ read the directions and warnings before use  
■ keep the carton, it contains important information.  
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
■ protect product from moisture  
■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, copovidone, glycol, silicon dioxide, sodium stearoyl fumarate, maize malto-dextrin, mannitol, magnesium, microcrystalline cellulose, sorbitol, sucralose, talc, titanium dioxide, triethyl citrate

**Questions or comments?** 1-800-919-9260; weekdays 7:30 AM to 5:00 PM EST

**Use** ■ treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
Allergy alert:  
■ do not use if you are allergic to lansoprazole  
■ lansoprazole may cause severe skin reactions. Symptoms may include: skin redness ■ rash  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
■ heartburn with lightheadedness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness  
■ frequent chest pain  
These may be signs of a serious condition. See your doctor.  
**Ask a doctor before use if you have** ■ liver disease  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss ■ nausea or vomiting ■ stomach pain  
**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain  
**If pregnant or breast-feeding,** ask a health care professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Location for lot number and expiration date

**NOT FOR RESALE**

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

**Safety Feature – Do not use if printed tablet blister unit is open or torn.**

Compare to the active ingredient in Prevacid® 24 HR\*



Treats Frequent Heartburn **24 HR**

# Lansoprazole

Delayed Release Orally Disintegrating Tablets 15 mg / Acid Reducer

**Dissolve Tabs**

**Melts in your mouth**  
Dissolves without water



Strawberry Flavor

**14 TABLETS**

One 14-day course of treatment  
May take 1 to 4 days for full effect



Actual Size

**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight. ■ If you smoke, quit smoking.

**FORMAT: STANDARD**

**TYPE SIZES:**

<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

**Drug Facts**

**Active ingredient (in each tablet)** Lansoprazole 15 mg  
Acid Reducer

**Purpose** Acid Reducer

**Use** ■ treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
Allergy alert: ■ do not use if you are allergic to lansoprazole ■ do not use if you cause severe skin reactions. Symptoms may include: ■ rash ■ blisters ■ skin redness ■ If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ chest pain or shoulder pain with shortness of breath; sweating; pain ■ heartburn with lightheadedness, sweating or dizziness ■ frequent chest pain ■ spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain

**Ask a doctor before use if you have** ■ liver disease ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, flavor, hypromellose, hydroxyethylcellulose, mazel, methylcellulose, methylcellulose, microcrystalline cellulose, polyacrylate, polyacrylate 80, polyethylene glycol, silicon dioxide, sodium stearoyl fumarate, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

**Other information** ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity ■ protect product from moisture ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**14-Day Course of Treatment**  
■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ do not crush or chew tablets ■ do not use if you are allergic to lansoprazole ■ do not use if you cause severe skin reactions. Symptoms may include: ■ rash ■ blisters ■ skin redness ■ If an allergic reaction occurs, stop use and seek medical help right away.

**Repeated 14-Day Course (if needed)**  
■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Directions** ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**  
■ take 1 tablet before eating in the morning ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ do not crush or chew tablets ■ do not use if you are allergic to lansoprazole ■ do not use if you cause severe skin reactions. Symptoms may include: ■ rash ■ blisters ■ skin redness ■ If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ chest pain or shoulder pain with shortness of breath; sweating; pain ■ heartburn with lightheadedness, sweating or dizziness ■ frequent chest pain ■ spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain

**Ask a doctor before use if you have** ■ liver disease ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, flavor, hypromellose, hydroxyethylcellulose, mazel, methylcellulose, methylcellulose, microcrystalline cellulose, polyacrylate, polyacrylate 80, polyethylene glycol, silicon dioxide, sodium stearoyl fumarate, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

Location for lot number and expiration date

8U174 FA C8 100422



KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

**Safety Feature –**  
Do not use if printed tablet blister unit is open or torn.

NDC 64239-181-01

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR\*

**24 HR**  
**Lansoprazole**

Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer



Strawberry Flavor

May take 1 to 4 days for full effect

**Dissolve Tabs**

**14 TABLETS**

ONE 14-DAY COURSE OF TREATMENT

**Melts in your mouth**  
Dissolves without water



Actual size

<b>FORMAT: STANDARD</b>	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique (continued)
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight. ■ If you smoke, quit smoking.

**Drug Facts**  
**Active ingredient (in each tablet)** Lansoprazole 15 mg  
**Purpose** Acid Reducer

**Use** ■ Treats frequent heartburn (occurs 2 or more days a week)  
■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
**Allergy alert:** ■ Do not use if you are allergic to lansoprazole  
■ Do not use if you cause severe skin reactions. Symptoms may include: ■ rash ■ blisters ■ skin redness  
■ If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
■ heartburn with lightheadedness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
■ frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have** ■ liver disease  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, flavor, hypromellose, hydroxyethylcellulose, maize malic acid, mannitol, magnesium, microcrystalline cellulose, polyacrylate, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, stearic acid, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

**Other information** ■ Read the directions and warnings before use  
■ Keep the carton. It contains important information.  
■ Store at 20-25°C (68-77°F) ■ Keep product out of high heat and humidity  
■ protect product from moisture  
■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**14-Day Course of Treatment**  
■ take 1 tablet before eating in the morning  
■ place the tablet on tongue; tablet disintegrates, with or without water.  
■ do not crush or chew tablets  
The tablets can also be swallowed whole with water.  
■ take every day for 14 days ■ do not take more than 1 tablet a day  
■ do not use for more than 14 days unless directed by your doctor  
■ If an allergic reaction occurs, stop use and seek medical help right away.

**Repeated 14-Day Course (if needed)**  
■ you may repeat a 14-day course every 4 months  
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

**Directions** ■ adults 18 years of age and older  
■ this product is to be used once a day (every 24 hours), every day for 14 days  
■ relief of symptoms within 24 hours

**Drug Facts (continued)**

Location for lot number and expiration date

8U174 FA C6-J 100422



KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

**Safety Feature –**  
Do not use if printed tablet blister unit is open or torn.

NDC 64239-181-01

Treats Frequent Heartburn



Compare to the active ingredient in Prevacid® 24 HR\*

Child-resistant

24 HR  
**Lansoprazole**

Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer



Strawberry Flavor

May take 1 to 4 days for full effect

Dissolve Tabs

14 TABLETS

ONE 14-DAY COURSE OF TREATMENT

Melts in your mouth  
Dissolves without water



Actual size

<b>FORMAT: STANDARD</b>	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Manufactured by:  
Dexcel Pharma Technologies Ltd  
10 Hakdima Street, Yokneam 2069200, Israel

\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®.

**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- If you smoke, quit smoking.

**Lansoprazole**  
Delayed Release  
Orally Disintegrating Tablets 15 mg

**Drug Facts**  
**Active ingredient (in each tablet)**  
Lansoprazole 15 mg.....Acid Reducer

**Use**  
■ treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
■ skin reddening ■ blisters ■ rash  
■ lansoprazole may cause severe skin reactions. Symptoms may include:  
■ do not use if you are allergic to lansoprazole  
■ do not use if you are allergic to lansoprazole

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
■ heartburn with lightheadedness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

**Ask a doctor before use if you have:** ■ liver disease  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Questions or comments? 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST**

**Drug Facts (continued)**

**Directions** ■ adults 18 years of age and older  
■ this product is to be used once a day (every 24 hours), every day for 14 days  
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**  
■ take 1 tablet before eating in the morning  
■ do not crush or chew tablets  
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.  
■ take every day for 14 days. ■ do not take more than 1 tablet a day  
■ do not use for more than 14 days unless directed by your doctor  
■ do not take this medicine with alcohol

**Repeated 14-Day Course (if needed)**  
■ you may repeat a 14-day course every 4 months  
■ do not take for more than 14 days or more often than every 4 months  
■ unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other information**  
■ read the directions and warnings before use  
■ keep the carton. It contains important information.  
■ store at 20-25°C (68-77°F). ■ Keep product out of high heat and humidity  
■ protect product from moisture  
■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolamide, flavor, hypromellose, hypromellose phthalate, maize polyorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

8U130 FA CB 100422

**Lansoprazole**  
Delayed Release  
Orally Disintegrating Tablets 15 mg

**Safety Feature – Do not use if printed tablet blister unit is open or torn.**

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

NDC 64239-181-02

**Treats Frequent Heartburn**

MELTech™  
Melts In Your Mouth

Compare to the active ingredient in Prevacid® 24 HR\*

**24 HR**

**Lansoprazole**  
Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer

May take 1 to 4 days for full effect

**Dissolve Tabs**

**28 TABLETS** TWO 14-DAY COURSES OF TREATMENT

**Melts in your mouth**  
Dissolves without water

Strawberry Flavor

Actual size

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

**Manufactured by:**  
**Dexcel Pharma Technologies Ltd**  
 10 Hakidma Street, Yokneam 2069200, Israel  
 Made in Israel

\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®.

**Tips for Managing Heartburn**  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Eat slowly and do not eat big meals.  
 ■ Do not eat late at night or just before bedtime.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight.  
 ■ If you smoke, quit smoking.

■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

**Drug Facts**  
 Active ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer

**Use**  
 ■ treats frequent heartburn (occurs 2 or more days a week)  
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
 ■ Allergy alert: lansoprazole may cause severe skin reactions. Symptoms may include: skin redness ■ blisters ■ rash  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
 ■ heartburn with lightheadedness, sweating or dizziness  
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

**Ask a doctor before use if you have:** ■ liver disease  
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.  
 ■ frequent wheezing, particularly with heartburn  
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
 Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:** ■ your heartburn continues or worsens  
 ■ you need to take this product for more than 14 days  
 ■ you need to take more than 1 course of treatment every 4 months  
 ■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

**Drug Facts (continued)**  
**Directions** ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**  
 ■ take 1 tablet before eating in the morning  
 ■ do not crush or chew tablets  
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.  
 ■ take every day for 14 days ■ do not take more than 1 tablet a day  
 ■ do not use for more than 14 days unless directed by your doctor  
 ■ do not take this medicine with alcohol

**Repeated 14-Day Course (if needed)**  
 ■ you may repeat a 14-day course every 4 months  
 ■ do not take for more than 14 days or more often than every 4 months  
**unless directed by a doctor**  
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other information**  
 ■ read the directions and warnings before use  
 ■ keep the carton. It contains important information.  
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
 ■ protect product from moisture  
 ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, flavor, hypromellose, hypromellose phthalate, maize polyorbate 80, propylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

8U130 FA CB-V 100422

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

NDC 64239-181-02

**Treats Frequent Heartburn**


 Melts In Your Mouth

Compare to the active ingredient in Prevacid® 24 HR\*


**24 HR**

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg  
 Acid Reducer

May take 1 to 4 days for full effect

**Dissolve Tabs**

**28 TABLETS** TWO 14-DAY COURSES OF TREATMENT

**Melts in your mouth**  
 Dissolves without water

Orally Disintegrating Tablets  
 Strawberry Flavor  
 Actual size

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Avoid fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods.  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight.  
 ■ If you smoke, quit smoking.

■ This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®.  
 ■ Manufactured by:  
 Dexcel Pharma Technologies Ltd  
 10 Hakdima Street, Yokneam 2069200, Israel  
 Made in Israel

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

**Drug Facts** (continued)

**Active ingredient (in each tablet)**  
Lansoprazole 15 mg.....Acid Reducer

**Use**  
 ■ treats frequent heartburn (occurs 2 or more days a week)  
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
 ■ lansoprazole may cause severe skin reactions. Symptoms may include:  
 ■ skin redness ■ blisters ■ rash  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
 ■ heartburn with lightheadedness, sweating or dizziness  
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

**Ask a doctor before use if you have:** ■ liver disease  
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.  
 ■ frequent wheezing, particularly with heartburn  
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
 Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:** ■ your heartburn continues or worsens  
 ■ you need to take this product for more than 14 days  
 ■ you need to take more than 1 course of treatment every 4 months  
 ■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

**Drug Facts**

**Directions** ■ adults 18 years of age and older  
 ■ this product is to be used once a day (every 24 hours), every day for 14 days  
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours


**14-Day Course of Treatment**  
 ■ take 1 tablet before eating in the morning  
 ■ do not crush or chew tablets  
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.  
 ■ take every day for 14 days. ■ do not take more than 1 tablet a day  
 ■ do not use for more than 14 days unless directed by your doctor  
 ■ do not take this medicine with alcohol

**Repeated 14-Day Course (if needed)**  
 ■ you may repeat a 14-day course every 4 months  
 ■ do not take for more than 14 days or more often than every 4 months  
**unless directed by a doctor**

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other information**  
 ■ read the directions and warnings before use  
 ■ keep the carton. It contains important information.  
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
 ■ protect product from moisture  
 ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, fawn, hypromellose, hypromellose phthalate, maize polyorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate



8U130 FA CB-A 100422

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg



**Safety Feature – Do not use if printed tablet blister unit is open or torn.**

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

NDC 64239-181-02

**Treats Frequent Heartburn**



Compare to the active ingredient in Prevacid® 24 HR\*

**Value 2 Pack**



**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg  
 Acid Reducer

May take 1 to 4 days for full effect

**Dissolve Tabs**

**28 TABLETS** TWO 14-DAY COURSES OF TREATMENT

**Melts in your mouth**  
 Dissolves without water

 Strawberry Flavor  
 Actual size

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Manufactured by:  
**Dexcel Pharma Technologies Ltd**  
10 Hakidma Street, Yokneam 2069200, Israel

\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®.

**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- If you smoke, quit smoking.

**Lansoprazole**  
Delayed Release  
Orally Disintegrating Tablets 15 mg

**Drug Facts**

**Active ingredient (in each tablet)**  
Lansoprazole 15 mg.....Acid Reducer

**Use**  
■ treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
■ lansoprazole may cause severe skin reactions. Symptoms may include:  
■ skin redness ■ blisters ■ rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
■ heartburn with lightheadedness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

**Ask a doctor before use if you have:** ■ liver disease  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

**Other information**  
■ read the directions and warnings before use  
■ keep the carton. It contains important information.  
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
■ protect product from moisture  
■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolamide, flavor, hypromellose, hypromellose phthalate, maize polyorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

**Directions** ■ adults 18 years of age and older  
■ this product is to be used once a day (every 24 hours), every day for 14 days  
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**  
■ take 1 tablet before eating in the morning  
■ do not crush or chew tablets  
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.  
■ take every day for 14 days. ■ do not take more than 1 tablet a day  
■ do not use for more than 14 days unless directed by your doctor  
■ do not take this medicine with alcohol

**Repeated 14-Day Course (if needed)**  
■ you may repeat a 14-day course every 4 months  
■ do not take for more than 14 days or more often than every 4 months  
■ unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

8U130 FA CB-J 100422

**Lansoprazole**  
Delayed Release  
Orally Disintegrating Tablets 15 mg

**Safety Feature – Do not use if printed tablet blister unit is open or torn.**

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

NDC 64239-181-02

**Treats Frequent Heartburn**

MELTech™  
Melts In Your Mouth

Compare to the active ingredient in Prevacid® 24 HR\*

**Child-resistant**

**24 HR**  
**Lansoprazole**  
Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer

May take 1 to 4 days for full effect

**Dissolve Tabs**

**28 TABLETS** TWO 14-DAY COURSES OF TREATMENT

**Melts in your mouth**  
Dissolves without water

Strawberry Flavor

Actual size

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Avoid fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods.  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight.  
 ■ If you smoke, quit smoking.

■ This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®.  
 ■ Manufactured by:  
 Dexcel Pharma Technologies Ltd  
 10 Hakidma Street, Yokneam 2069200, Israel  
 Made in Israel

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

**Drug Facts** (continued)

**Active ingredient (in each tablet)**  
Lansoprazole 15 mg.....Acid Reducer

**Use**  
 ■ treats frequent heartburn (occurs 2 or more days a week)  
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
 ■ lansoprazole may cause severe skin reactions. Symptoms may include:  
 ■ skin redness ■ blisters ■ rash  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
 ■ heartburn with **lightheadness, sweating or dizziness**  
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

**Ask a doctor before use if you have:** ■ liver disease  
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.  
 ■ frequent wheezing, particularly with heartburn  
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
 Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:** ■ your heartburn continues or worsens  
 ■ you need to take this product for more than 14 days  
 ■ you need to take more than 1 course of treatment every 4 months  
 ■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

**Drug Facts**

**Directions** ■ adults 18 years of age and older  
 ■ this product is to be used once a day (every 24 hours), every day for 14 days  
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**  
 ■ take 1 tablet before eating in the morning  
 ■ do not crush or chew tablets  
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.  
 ■ take every day for 14 days. ■ do not take more than 1 tablet a day  
 ■ do not use for more than 14 days unless directed by your doctor  
 ■ do not take this medicine with alcohol

**Repeated 14-Day Course (if needed)**  
 ■ you may repeat a 14-day course every 4 months  
 ■ do not take for more than 14 days or more often than every 4 months  
**unless directed by a doctor**

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other information**  
 ■ read the directions and warnings before use  
 ■ keep the carton. It contains important information.  
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
 ■ protect product from moisture  
 ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

8U130 FA CB-K 100422



**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed  
 tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
 IMPORTANT INFORMATION.

NDC 64239-181-02  
**Treats Frequent Heartburn**

  
 Melts In Your Mouth

Compare to the active ingredient  
 in Prevacid® 24 HR\*

**Buy One 14 Count,  
 Get One 14 Count Free!**

  
**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg  
 Acid Reducer

May take 1 to 4 days for full effect

**Dissolve Tabs**

**28 TABLETS** TWO 14-DAY COURSES OF TREATMENT

**Melts in your mouth**  
 Dissolves without water

  
 Strawberry Flavor

  
 Actual size

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Avoid fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods.  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight.  
 ■ If you smoke, quit smoking.

■ This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®.  
 ■ Manufactured by:  
 Dexcel Pharma Technologies Ltd  
 10 Hakidma Street, Yokneam 2069200, Israel  
 Made in Israel

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

**Drug Facts** (continued)

**Active ingredient (in each tablet)**  
Lansoprazole 15 mg.....Acid Reducer

**Use**  
 ■ treats frequent heartburn (occurs 2 or more days a week)  
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
 ■ lansoprazole may cause severe skin reactions. Symptoms may include:  
 ■ skin redness ■ blisters ■ rash  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
 ■ heartburn with lightheadedness, sweating or dizziness  
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

**Ask a doctor before use if you have:** ■ liver disease  
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.  
 ■ frequent wheezing, particularly with heartburn  
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
 Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:** ■ your heartburn continues or worsens  
 ■ you need to take this product for more than 14 days  
 ■ you need to take more than 1 course of treatment every 4 months  
 ■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

**Drug Facts**

**14-Day Course of Treatment**  
 ■ take 1 tablet before eating in the morning  
 ■ do not crush or chew tablets  
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.  
 ■ take every day for 14 days. ■ do not take more than 1 tablet a day  
 ■ do not use for more than 14 days unless directed by your doctor  
 ■ do not take this medicine with alcohol

**Repeated 14-Day Course (if needed)**  
 ■ you may repeat a 14-day course every 4 months  
 ■ do not take for more than 14 days or more often than every 4 months  
**unless directed by a doctor**

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other information**  
 ■ read the directions and warnings before use  
 ■ keep the carton. It contains important information.  
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
 ■ protect product from moisture  
 ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

8U130 FA CB-L 100422



**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed  
 tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
 IMPORTANT INFORMATION.

NDC 64239-181-02  
**Treats Frequent Heartburn**

  
 Melts In Your Mouth

Compare to the active ingredient  
 in Prevacid® 24 HR\*

**100% More Free!**  
**14 Free Tablets!**

  
**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg  
 Acid Reducer

May take 1 to 4 days for full effect

**Dissolve Tabs**

**28 TABLETS** TWO 14-DAY COURSES OF TREATMENT

**Melts in your mouth**  
 Dissolves without water

  
 Strawberry Flavor

  
 Actual size

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Avoid fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods.  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight.  
 ■ If you smoke, quit smoking.

■ This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®.  
 10 Hakidma Street, Yokneam 2069200, Israel  
 Manufactured by:  
**Dexcel Pharma Technologies Ltd**

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

**Drug Facts** (continued)

**Active ingredient (in each tablet)**  
Lansoprazole 15 mg.....Acid Reducer

**Use**  
 ■ treats frequent heartburn (occurs 2 or more days a week)  
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
 ■ lansoprazole may cause severe skin reactions. Symptoms may include:  
 ■ skin redness ■ blisters ■ rash  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
 ■ heartburn with **lightheadness, sweating or dizziness**  
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

**Ask a doctor before use if you have:** ■ liver disease  
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.  
 ■ frequent wheezing, particularly with heartburn  
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
 Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:** ■ your heartburn continues or worsens  
 ■ you need to take this product for more than 14 days  
 ■ you need to take more than 1 course of treatment every 4 months  
 ■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

**Drug Facts**

**Directions** ■ adults 18 years of age and older  
 ■ this product is to be used once a day (every 24 hours), every day for 14 days  
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours


**14-Day Course of Treatment**  
 ■ take 1 tablet before eating in the morning  
 ■ do not crush or chew tablets  
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.  
 ■ take every day for 14 days. ■ do not take more than 1 tablet a day  
 ■ do not use for more than 14 days unless directed by your doctor  
 ■ do not take this medicine with alcohol

**Repeated 14-Day Course (if needed)**  
 ■ you may repeat a 14-day course every 4 months  
 ■ do not take for more than 14 days or more often than every 4 months  
**unless directed by a doctor**

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other information**  
 ■ read the directions and warnings before use  
 ■ keep the carton. It contains important information.  
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
 ■ protect product from moisture  
 ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate



**8U130 FA CB-N 100422**

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg



**Safety Feature – Do not use if printed tablet blister unit is open or torn.**

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

NDC 64239-181-02

**Treats Frequent Heartburn**



Compare to the active ingredient in Prevacid® 24 HR\*

**Bonus Size!**  
**14 Tablets Free**


**24 HR**

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg  
 Acid Reducer

May take 1 to 4 days for full effect

**Dissolve Tabs**

**28 TABLETS** TWO 14-DAY COURSES OF TREATMENT

**Melts in your mouth**  
 Dissolves without water

 Strawberry Flavor  
 Actual size

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Avoid fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods.  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight.  
 ■ If you smoke, quit smoking.

■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Avoid fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods.  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight.  
 ■ If you smoke, quit smoking.

■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Avoid fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods.  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight.  
 ■ If you smoke, quit smoking.

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

**Drug Facts**  
 Active ingredient (in each tablet) Lansoprazole 15 mg.....Acid Reducer  
**Use**  
 ■ treats frequent heartburn (occurs 2 or more days a week)  
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect  
**Warnings**  
 ■ do not use if you are allergic to lansoprazole  
 ■ lansoprazole may cause severe skin reactions. Symptoms may include:  
 ■ skin redness ■ blisters ■ rash  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Do not use if you have:**  
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
 ■ heartburn with lightheadedness, sweating or dizziness  
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness  
**Ask a doctor before use if you have:** ■ liver disease  
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.  
 ■ frequent wheezing, particularly with heartburn  
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain  
**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
 Acid reducers may interact with certain prescription drugs.  
**Stop use and ask a doctor if:** ■ your heartburn continues or worsens  
 ■ you need to take this product for more than 14 days  
 ■ you need to take more than 1 course of treatment every 4 months  
 ■ you get diarrhea ■ you develop a rash or joint pain  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).  
**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

**Drug Facts (continued)**  
**Directions** ■ adults 18 years of age and older  
 ■ this product is to be used once a day (every 24 hours), every day for 14 days  
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours  
**14-Day Course of Treatment**  
 ■ take 1 tablet before eating in the morning  
 ■ do not crush or chew tablets  
 ■ place the tablet on tongue; tablet disintegrates, with or without water.  
 The tablets can also be swallowed whole with water.  
 ■ do not take more than 1 tablet a day  
 ■ take every day for 14 days ■ do not take more than 1 tablet a day  
 ■ do not use for more than 14 days unless directed by your doctor  
 ■ do not take this medicine with alcohol  
**Repeated 14-Day Course (if needed)**  
 ■ you may repeat a 14-day course every 4 months  
 ■ do not take for more than 14 days or more often than every 4 months  
**unless directed by a doctor**  
 ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.  
**Other information**  
 ■ read the directions and warnings before use  
 ■ keep the carton. It contains important information.  
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
 ■ protect product from moisture  
 ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

8U130 FA CB-P 100422

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

NDC 64239-181-02

Treats Frequent Heartburn

MELTech™  
 Melts In Your Mouth

Compare to the active ingredient in Prevacid® 24 HR\*

**Bonus Size! 14 Free**

**24 HR**  
**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg  
 Acid Reducer

May take 1 to 4 days for full effect

**Dissolve Tabs**

**28 TABLETS** TWO 14-DAY COURSES OF TREATMENT

**Melts in your mouth**  
 Dissolves without water

Strawberry Flavor

Actual size

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Avoid fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods.  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight.  
 ■ If you smoke, quit smoking.

■ This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®.  
 ■ Manufactured by:  
 Dexcel Pharma Technologies Ltd  
 10 Hakdima Street, Yokneam 2069200, Israel  
 Made in Israel

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

**Drug Facts** (continued)

**Active ingredient (in each tablet)**  
Lansoprazole 15 mg.....Acid Reducer

**Use**  
 ■ treats frequent heartburn (occurs 2 or more days a week)  
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
 ■ lansoprazole may cause severe skin reactions. Symptoms may include:  
 ■ skin redness ■ blisters ■ rash  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
 ■ heartburn with lightheadedness, sweating or dizziness  
 ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

**Ask a doctor before use if you have:** ■ liver disease  
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.  
 ■ frequent wheezing, particularly with heartburn  
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
 Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:** ■ your heartburn continues or worsens  
 ■ you need to take this product for more than 14 days  
 ■ you need to take more than 1 course of treatment every 4 months  
 ■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

**Drug Facts**

**Directions** ■ adults 18 years of age and older  
 ■ this product is to be used once a day (every 24 hours), every day for 14 days  
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours


**14-Day Course of Treatment**  
 ■ take 1 tablet before eating in the morning  
 ■ do not crush or chew tablets  
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.  
 ■ take every day for 14 days. ■ do not take more than 1 tablet a day  
 ■ do not use for more than 14 days unless directed by your doctor  
 ■ do not take this medicine with alcohol

**Repeated 14-Day Course (if needed)**  
 ■ you may repeat a 14-day course every 4 months  
 ■ do not take for more than 14 days or more often than every 4 months  
**unless directed by a doctor**

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other information**  
 ■ read the directions and warnings before use  
 ■ keep the carton. It contains important information.  
 ■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
 ■ protect product from moisture  
 ■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.


**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate



**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed  
 tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
 IMPORTANT INFORMATION.



8U130 FA CB-R 100422

**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed  
 tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
 IMPORTANT INFORMATION.



NDC 64239-181-02

**Treats Frequent Heartburn**



Compare to the active ingredient  
 in Prevacid® 24 HR\*

**Bonus! + 14 Free**



**Lansoprazole**  
 Delayed Release  
 Orally Disintegrating Tablets 15 mg  
 Acid Reducer

May take 1 to 4 days for full effect

**Dissolve Tabs**

**28 TABLETS** TWO 14-DAY COURSES OF TREATMENT

**Melts in your mouth**  
 Dissolves without water


  
 Strawberry Flavor


  
 Actual size

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Manufactured by:  
**Dexcel Pharma Technologies Ltd**  
10 Hakdima Street, Yokneam 2069200, Israel

\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc., owner of the registered trademark Prevacid®.

**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- Raise the head of your bed.
- Do not lie flat or bend over soon after eating.

**Lansoprazole**  
Delayed Release  
Orally Disintegrating Tablets 15 mg

**Drug Facts**

**Active ingredient (in each tablet)**  
Lansoprazole 15 mg.....Acid Reducer

**Use**  
■ treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
■ lansoprazole may cause severe skin reactions. Symptoms may include:  
■ skin redness ■ blisters ■ rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
■ heartburn with lightheadedness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

**Ask a doctor before use if you have:** ■ liver disease  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Questions or comments?** 1-800-719-9260; weekdays 7:30 AM to 5:00 PM EST

**Other information**  
■ read the directions and warnings before use  
■ keep the carton. It contains important information.  
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
■ protect product from moisture  
■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, crospovidone, flavor, hypromellose, hypromellose phthalate, maize polyorbate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

**Directions** ■ adults 18 years of age and older  
■ this product is to be used once a day (every 24 hours), every day for 14 days  
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**  
■ take 1 tablet before eating in the morning  
■ do not crush or chew tablets  
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.  
■ take every day for 14 days. ■ do not take more than 1 tablet a day  
■ do not use for more than 14 days unless directed by your doctor  
■ do not take this medicine with alcohol

**Repeated 14-Day Course (if needed)**  
■ you may repeat a 14-day course every 4 months  
■ do not take for more than 14 days or more often than every 4 months  
■ unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Location for lot number and expiration date

8U130 FA CB-S 100422

**Lansoprazole**  
Delayed Release  
Orally Disintegrating Tablets 15 mg

**Safety Feature – Do not use if printed tablet blister unit is open or torn.**

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

NDC 64239-181-02

**Treats Frequent Heartburn**

MELTech™  
Melts In Your Mouth

Compare to the active ingredient in Prevacid® 24 HR\*

**Bonus! 14 Tablets FREE**

**24 HR**  
**Lansoprazole**  
Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer

May take 1 to 4 days for full effect

**Dissolve Tabs**

**28 TABLETS** TWO 14-DAY COURSES OF TREATMENT

**Melts in your mouth**  
Dissolves without water

Strawberry Flavor

Actual size

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight. ■ If you smoke, quit smoking.

\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc.,  
 owner of the registered trademark Prevacid®.  
 10 Hakdima Street, Yokneam 2069200, Israel  
 Manufactured by:  
**Decel Pharma Technologies Ltd**  
 Made in Israel

Location for  
lot number and  
expiration date

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Drug Facts

**Active Ingredient (in each tablet)**  
Lansoprazole 15 mg, Acid Reducer

**Use**  
■ Treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn, this drug may take 1 to 4 days for full effect

**Warnings**  
do not use if you are allergic to lansoprazole  
lansoprazole may cause severe skin reactions. Symptoms may include:  
■ rash ■ skin redness ■ blisters ■ stomach pain  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or body or black stools  
■ heartburn with tightness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
■ frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have** ■ liver disease  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain

**If pregnant or breast-feeding**, ask a health care professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

5:00 PM EST

**Questions or comments? 1-800-719-9260, weekdays 7:30 AM to**

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed  
tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
IMPORTANT INFORMATION.

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Drug Facts

**Active Ingredient (in each tablet)**  
Lansoprazole 15 mg, Acid Reducer

**Use**  
■ Treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn, this drug may take 1 to 4 days for full effect

**Warnings**  
do not use if you are allergic to lansoprazole  
lansoprazole may cause severe skin reactions. Symptoms may include:  
■ rash ■ skin redness ■ blisters ■ stomach pain  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or body or black stools  
■ heartburn with tightness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
■ frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have** ■ liver disease  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain

**If pregnant or breast-feeding**, ask a health care professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

5:00 PM EST

**Questions or comments? 1-800-719-9260, weekdays 7:30 AM to**

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed  
tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
IMPORTANT INFORMATION.

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Drug Facts

**Active Ingredient (in each tablet)**  
Lansoprazole 15 mg, Acid Reducer

**Use**  
■ Treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn, this drug may take 1 to 4 days for full effect

**Warnings**  
do not use if you are allergic to lansoprazole  
lansoprazole may cause severe skin reactions. Symptoms may include:  
■ rash ■ skin redness ■ blisters ■ stomach pain  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or body or black stools  
■ heartburn with tightness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
■ frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have** ■ liver disease  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain

**If pregnant or breast-feeding**, ask a health care professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

5:00 PM EST

**Questions or comments? 1-800-719-9260, weekdays 7:30 AM to**

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed  
tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
IMPORTANT INFORMATION.

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight. ■ If you smoke, quit smoking.

\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc.,  
 owner of the registered trademark Prevacid®.  
 10 Hakdima Street, Yokneam 2069200, Israel  
 Manufactured by:  
 Dexcel Pharma Technologies Ltd  
 Made in Israel

Location for  
lot number and  
expiration date

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Drug Facts

**Active Ingredient (in each tablet)**  
Lansoprazole 15 mg, Acid Reducer

**Purpose**  
■ Treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn, this drug may take 1 to 4 days for full effect

**Warnings**  
do not use if you are allergic to lansoprazole  
lansoprazole may cause severe skin reactions. Symptoms may include:  
■ rash ■ skin redness ■ blisters ■ stomach pain  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or body or black stools  
■ heartburn with tightness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
■ frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have** ■ liver disease  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain

**If pregnant or breast-feeding**, ask a health care professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

5:00 PM EST

**Questions or comments? 1-800-719-9260, weekdays 7:30 AM to**

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize polydoxtrin, methylcellulose, microcrystalline cellulose, polyethylene glycol, polyethylene glycol, sodium lauryl sulfate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

**Other information**  
■ read the directions and warnings before use  
■ keep the carton. It contains important information.  
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**Drug Facts (continued)**

**Directions** ■ adults 18 years of age and older  
■ this product is to be used once a day (every 24 hours), every day for 14 days  
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**  
■ take 1 tablet before eating in the morning  
■ do not crush or chew tablets  
■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.

**Repeated 14-Day Courses (if needed)**  
■ you may repeat a 14-day course every 4 months  
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

**Other information**  
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.  
■ do not take this medicine with alcohol

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed  
tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
IMPORTANT INFORMATION.

0000000000

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer

May take 1 to 4 days for full effect

Dissolve Tabs

42 TABLETS

THREE 14-DAY COURSES  
OF TREATMENT

  
 Melts In Your Mouth

Compare to the active ingredient  
in Prevacid® 24 HR\*

24 HR

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer

May take 1 to 4 days for full effect

Dissolve Tabs

42 TABLETS

THREE 14-DAY COURSES  
OF TREATMENT

Melts in your mouth  
Dissolves without water

Orally Disintegrating Tablets

Strawberry Flavor

Actual size

0000000000

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight.  
 ■ If you smoke, quit smoking.

\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc.,  
 owner of the registered trademark Prevacid®.  
 10 Hakdima Street, Yokneam 2069200, Israel  
 Manufactured by:  
 Dexcel Pharma Technologies Ltd

Location for  
lot number and  
expiration date

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Drug Facts

**Active Ingredient (in each tablet)**  
Lansoprazole 15 mg, Acid Reducer

**Purpose**  
Treats frequent heartburn (occurs 2 or more days a week)  
not intended for immediate relief of heartburn, this drug may take 1 to 4 days for full effect

**Warnings**  
do not use if you are allergic to lansoprazole  
lansoprazole may cause severe skin reactions. Symptoms may include:  
rash  
skin redness  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
trouble or pain swallowing food, vomiting with blood, or body or black stools  
heartburn with tightness, sweating or dizziness  
chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have** ■ liver disease  
had heartburn over 3 months. This may be a sign of a more serious condition.  
frequent wheezing, particularly with heartburn  
nausea or vomiting ■ stomach pain  
unexplained weight loss

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
you need to take this product for more than 14 days  
you need to take more than 1 course of treatment every 4 months  
you get diarrhea ■ you develop a rash or joint pain

**If pregnant or breast-feeding,** ask a health care professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

5:00 PM EST

**Questions or comments? 1-800-719-9260, weekdays 7:30 AM to**

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg



Safety Feature – Do not use if printed  
tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
IMPORTANT INFORMATION.

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer

Dissolve Tabs

42 TABLETS

THREE 14-DAY COURSES  
OF TREATMENT

MELTech™

Melts In Your Mouth

Compare to the active ingredient  
in Prevacid® 24 HR\*

Value 3 Pack



Strawberry  
Flavor



Actual  
size

Melts in your mouth

Dissolves without water

24 HR

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer

May take 1 to 4 days for full effect

FORMAT: STANDARD

TYPE SIZES:

<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

Treats Frequent Heartburn

NDC 64239-181-03

8U155 FA CS-B 100422



■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are rich, spicy, fatty and  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight. ■ If you smoke, quit smoking.

\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc.,  
 owner of the registered trademark Prevacid®.  
 10 Hakdima Street, Yokneam 2069200, Israel  
 Manufactured by:  
 Dexcel Pharma Technologies Ltd  
 Made in Israel

Location for  
lot number and  
expiration date

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

**Drug Facts**

**Active Ingredient (in each tablet)**  
Lansoprazole 15 mg, Acid Reducer

**Purpose**  
■ treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn, this drug may take 1 to 4 days for full effect

**Warnings**  
do not use if you are allergic to lansoprazole  
lansoprazole may cause severe skin reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or body or black stools  
■ heartburn with tightness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
■ frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have** ■ liver disease  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain

**If pregnant or breast-feeding**, ask a health care professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

5:00 PM EST

**Questions or comments? 1-800-719-9260, weekdays 7:30 AM to**

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize polyorbital, mannitol, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

**Other information**  
■ read the directions and warnings before use  
■ keep the carton. It contains important information.  
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

**Safety Feature – Do not use if printed  
tablet blister unit is open or torn.**

KEEP THE CARTON. IT CONTAINS  
IMPORTANT INFORMATION.



Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

NDC 64239-181-03

Treats Frequent Heartburn

24 HR

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer

May take 1 to 4 days for full effect

Dissolve Tabs

42 TABLETS

THREE 14-DAY COURSES  
OF TREATMENT

Melts in your mouth  
*Dissolves without water*



Strawberry  
Flavor



15  
Actual  
size

Compare to the active ingredient  
in Prevacid® 24 HR\*

Child-resistant

Melts In Your Mouth



8U155 FA CS-J 100422

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
(continued)	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box







■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight. ■ If you smoke, quit smoking.

\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc.,  
 owner of the registered trademark Prevacid®.  
 10 Hakdima Street, Yokneam 2069200, Israel  
 Manufactured by:  
 Dexcel Pharma Technologies Ltd  
 Made in Israel

Location for  
lot number and  
expiration date

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Drug Facts

**Active Ingredient (in each tablet)**  
Lansoprazole 15 mg, Acid Reducer

**Purpose**  
Treats frequent heartburn (occurs 2 or more days a week)  
not intended for immediate relief of heartburn, this drug may take 1 to 4 days for full effect

**Warnings**  
do not use if you are allergic to lansoprazole  
lansoprazole may cause severe skin reactions. Symptoms may include:  
rash ■ skin redness ■ blisters ■ if an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
touble or pain swallowing food, vomiting with blood, or body or black stools  
heartburn with tightness, sweating or dizziness  
chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have** ■ liver disease  
had heartburn over 3 months. This may be a sign of a more serious condition.  
frequent wheezing, particularly with heartburn  
unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
you need to take this product for more than 14 days  
you need to take more than 1 course of treatment every 4 months  
you get diarrhea ■ you develop a rash or joint pain

**If pregnant or breast-feeding**, ask a health care professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

5:00 PM EST

**Questions or comments? 1-800-719-9260, weekdays 7:30 AM to**

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopolone, flavor, hypromellose, hypromellose phthalate, maize polydoxtrin, methylcellulose, microcrystalline cellulose, polyethylene glycol, polyethylene glycol, sodium lauryl sulfate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

**Other information**  
read the directions and warnings before use  
keep the carton. It contains important information.  
store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**14-Day Course of Treatment**  
relief of symptoms within 24 hours  
it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**  
take 1 tablet before eating in the morning  
do not crush or chew tablets  
place the tablet on tongue; tablet disintegrates, with or without water.  
The tablets can also be swallowed whole with water.  
take every day for 14 days ■ do not take more than 1 tablet a day  
do not use for more than 14 days unless directed by your doctor  
do not take this medicine with alcohol

**Repeated 14-Day Courses (if needed)**  
you may repeat a 14-day course every 4 months  
do not take for more than 14 days or more often than every 4 months  
unless directed by a doctor

children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed  
tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
IMPORTANT INFORMATION.



NDC 64239-181-03

Treats Frequent Heartburn



Melts In Your Mouth

Compare to the active ingredient  
in Prevacid® 24 HR\*

Bonus Size! 14 Free



24 HR

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer

May take 1 to 4 days for full effect

Dissolve Tabs

42 TABLETS

THREE 14-DAY COURSES  
OF TREATMENT



Strawberry  
Flavor



Actual  
size

Melts in your mouth

Dissolves without water



FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box



■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight.  
 ■ If you smoke, quit smoking.

\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc.,  
 owner of the registered trademark Prevacid®.  
 10 Hakdima Street, Yokneam 2069200, Israel  
 Manufactured by:  
 Dexcel Pharma Technologies Ltd  
 Made in Israel

Location for  
lot number and  
expiration date

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Drug Facts

**Active Ingredient (in each tablet)**  
Lansoprazole 15 mg, Acid Reducer

**Use**  
■ Treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn, this drug may take 1 to 4 days for full effect

**Warnings**  
do not use if you are allergic to lansoprazole  
lansoprazole may cause severe skin reactions. Symptoms may include:  
■ rash ■ skin redness ■ blisters ■ rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or body or black stools  
■ heartburn with **tightness, sweating or dizziness**  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
■ frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have** ■ liver disease  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain

**If pregnant or breast-feeding**, ask a health care professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

5:00 PM EST

**Questions or comments? 1-800-719-9260, weekdays 7:30 AM to**

**Inactive ingredients** ascorbic acid, cetyl alcohol, colloidal silicon dioxide, copovidone, croscopollose, fumed, hypromellose, hypromellose phthalate, maize polyborate 80, polyethylene glycol, silicon dioxide, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

**Other information**  
■ read the directions and warnings before use  
■ keep the carton. It contains important information.  
■ store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity  
■ take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

**Drug Facts (continued)**

**Directions** ■ adults 18 years of age and older  
■ this product is to be used once a day (every 24 hours), every day for 14 days  
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**  
■ take 1 tablet before eating in the morning  
■ do not crush or chew tablets  
■ place the tablet on tongue; tablet disintegrates, with or without water.  
The tablets can also be swallowed whole with water.  
■ take every day for 14 days ■ do not take more than 1 tablet a day  
■ do not use for more than 14 days unless directed by your doctor  
■ do not take this medicine with alcohol

**Repeated 14-Day Course (if needed)**  
■ you may repeat a 14-day course every 4 months  
■ do not take for more than 14 days or more often than every 4 months  
unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed  
tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
IMPORTANT INFORMATION.



NDC 64239-181-03

Treats Frequent Heartburn



Melts In Your Mouth

Compare to the active ingredient  
in Prevacid® 24 HR\*

Bonus! + 14 Free



24 HR

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer

May take 1 to 4 days for full effect

Dissolve Tabs

42 TABLETS

THREE 14-DAY COURSES  
OF TREATMENT



Strawberry  
Flavor



Actual  
size

Melts in your mouth

Dissolves without water



FORMAT: STANDARD

**TYPE SIZES:**

<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

8U155 FA CS-R 100422



■ Do not eat late at night or just before bedtime.  
 ■ Eat slowly and do not eat big meals.  
 ■ Fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
 ■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and  
 ■ Do not lie flat or bend over soon after eating.  
 ■ Raise the head of your bed.  
 ■ Wear loose-fitting clothing around your stomach.  
 ■ If you are overweight, lose weight. ■ If you smoke, quit smoking.

\*This product is not manufactured or distributed by Takeda Pharmaceuticals U.S.A., Inc.,  
 owner of the registered trademark Prevacid®.  
 10 Hakdima Street, Yokneam 2069200, Israel  
 Manufactured by:  
**Decel Pharma Technologies Ltd**  
 Made in Israel

Location for  
lot number and  
expiration date

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Drug Facts

**Active Ingredient (in each tablet)**  
Lansoprazole 15 mg, Acid Reducer

**Purpose**  
Treats frequent heartburn (occurs 2 or more days a week)  
not intended for immediate relief of heartburn, this drug may take 1 to 4 days for full effect

**Warnings**  
do not use if you are allergic to lansoprazole  
lansoprazole may cause severe skin reactions. Symptoms may include:  
rash ■ skin redness ■ blisters ■ if an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
trouble or pain swallowing food, vomiting with blood, or body or black stools  
heartburn with tightness, sweating or dizziness  
chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have** ■ liver disease  
had heartburn over 3 months. This may be a sign of a more serious condition.  
frequent wheezing, particularly with heartburn  
unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**  
Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if** ■ your heartburn continues or worsens  
you need to take this product for more than 14 days  
you need to take more than 1 course of treatment every 4 months  
you get diarrhea ■ you develop a rash or joint pain

**If pregnant or breast-feeding**, ask a health care professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

5:00 PM EST

**Questions or comments? 1-800-719-9260, weekdays 7:30 AM to**

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

Safety Feature – Do not use if printed  
tablet blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS  
IMPORTANT INFORMATION.



Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg

NDC 64239-181-03

Treats Frequent Heartburn

Compare to the active ingredient  
in Prevacid® 24 HR\*

Bonus! 14 Tablets FREE

24 HR

Lansoprazole

Delayed Release  
Orally Disintegrating Tablets 15 mg  
Acid Reducer

May take 1 to 4 days for full effect

Dissolve Tabs

42 TABLETS

THREE 14-DAY COURSES  
OF TREATMENT

Melts in your mouth

Dissolves without water



Strawberry  
Flavor



15  
Actual  
size

8U155 FA CS-S 100422

FORMAT: STANDARD	
<b>TYPE SIZES:</b>	
<b>TITLE</b>	9 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>TITLE (continued)</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>(continued)</b>	8 pt Helvetica Neue LT Std 57 Condensed
<b>HEADINGS</b>	8 pt Helvetica Neue LT Std 77 Bold Condensed Oblique
<b>SUBHEADINGS</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>TEXT</b>	6 pt Helvetica Neue LT Std 57 Condensed
<b>LEADING</b>	6.5 pt
<b>BULLETS</b>	5 pt ITC Zapf Dingbats Std Medium (square) with 2 ems (MM) spacing between statements
<b>TELEPHONE #</b>	6 pt Helvetica Neue LT Std 77 Bold Condensed
<b>HEAVY LINES</b>	1.5 pt
<b>HAIRLINES</b>	0.5 pt Extend to within 2 spaces of the "Drug Facts" box

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

NUSHIN F TODD  
11/01/2022 01:46:06 PM