


Emergency Treatment of Opioid Overdose

Important:

- For use in the nose only
- Do not test nasal spray device before use
- 1 nasal spray device contains 1 dose of medicine
- Each device sprays 1 time only



Drug Facts

Active ingredient (in each spray)	Purpose
Naloxone hydrochloride 4 mg.....	Emergency treatment of opioid overdose

Uses

- to “revive” someone during an overdose from many **prescription pain medications** or **street drugs such as heroin**
- this medicine can save a life

Directions

1 CHECK



Step 1: CHECK if you suspect an overdose:

- **CHECK** for a suspected overdose: the person will not wake up or is very sleepy or not breathing well
 - yell “Wake up!”
 - shake the person gently
 - if the person is not awake, go to Step 2

2 GIVE



Step 2: Give 1st dose in the nose

- **HOLD** the nasal spray device with your thumb on the bottom of the plunger
- **INSERT** the nozzle into either NOSTRIL
- **PRESS** the plunger firmly to give the 1st dose
- 1 nasal spray device contains 1 dose

3 CALL



Step 3: Call 911

- **CALL 911** immediately after giving the 1st dose

4 WATCH/GIVE



Step 4: WATCH & GIVE


- **WAIT** 2-3 minutes after the 1st dose to give the medicine time to work
- if the person wakes up: Go to Step 5
- if the person does not wake up:
 - **CONTINUE TO GIVE** doses every 2-3 minutes until the person wakes up
 - it is safe to keep giving doses

5 STAY




Step 5: STAY

- **STAY** until ambulance arrives: even if the person wakes up
- **GIVE** another dose if the person becomes very sleepy again
- You may need to give all the doses in the pack



Unfold for Drug Facts

Unfold for Drug Facts



Drug Facts Continued on Other Side

Drug Facts (continued)

Warning

When using this product some people may experience symptoms when they wake up, such as shaking, sweating, nausea, or feeling angry. This is to be expected.

Other information

- store at room temperature or refrigerator, between 2°C to 25°C (36°F to 77°F)
- do not freeze
- avoid excessive heat above 40°C (104°F)
- protect from light
- the product is packaged in individually-sealed blisters. Do not use if the blister is open or torn, or if the device appears damaged.

Inactive ingredients

benzalkonium chloride, edetate disodium, hydrochloric acid, sodium chloride, water

Questions?

call 1-844-4NARCAN (1-844-462-7226) or go to www.narcan.com

EMERGENT[®]

For opioid emergencies, call 911. For questions on NARCAN, call 1-844-4NARCAN (1-844-462-7226) or go to www.narcan.com.
©2025 Emergent Devices Inc. EMERGENT[®] and NARCAN[®] are registered trademarks of Emergent BioSolutions Inc, or its subsidiaries.

LBL040711

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
11/21/2025 04:05:34 PM