

Xyzal Allergy 24 Hour Tablet Blister Carton

01.06.17

Xyzal_Allergy 10ct_CTN_2016_r5_CLEAN

| Drug Facts | Purpose |
|---|---------|
| Active ingredient (in each tablet) Levocetirizine dihydrochloride 5 mg....Antihistamine | |
| Uses temporarily relieves these symptoms due to hay fever or other respiratory allergies: <ul style="list-style-type: none"> ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat | |
| Warnings Do not use <ul style="list-style-type: none"> ■ if you have kidney disease ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine | |
| Ask a doctor before use if you have <ul style="list-style-type: none"> ■ ever had trouble urinating or emptying your bladder | |
| When using this product <ul style="list-style-type: none"> ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery | |
| Stop use and ask a doctor if <ul style="list-style-type: none"> ■ you have trouble urinating or emptying your bladder ■ an allergic reaction to this product occurs. Seek medical help right away. | |
| If pregnant or breast-feeding: <ul style="list-style-type: none"> ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use | |
| Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). | |

Read directions and warnings before use. Keep this carton. It has important information.

| Drug Facts (continued) | |
|---|--|
| Directions adults 65 years of age and older | <ul style="list-style-type: none"> ■ ask a doctor |
| adults and children 12-64 years of age | <ul style="list-style-type: none"> ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms |
| children 6-11 years of age | <ul style="list-style-type: none"> ■ take ½ tablet (2.5 mg) once daily in the evening ■ do not take more than ½ tablet (2.5 mg) in 24 hours |
| children under 6 years of age | <ul style="list-style-type: none"> ■ do not use |
| consumers with kidney disease | <ul style="list-style-type: none"> ■ do not use |
| Other information | <ul style="list-style-type: none"> ■ store between 20° and 25°C (68° and 77°F) ■ safety sealed: do not use if carton was opened or if individual blister unit is open or torn |
| Inactive ingredients | colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide |
| Questions or comments? | call 1-800-630-1610 |

NDC 41167-3510-0

CLINICALLY PROVEN RELIEF FROM INDOOR AND OUTDOOR ALLERGIES

CLINICALLY PROVEN RELIEF IN 60 MINUTES

CLINICALLY PROVEN RELIEF FOR 24 HOURS

ORIGINAL PRESCRIPTION STRENGTH

XYZAL[®]

ALLERGY 24HR

levocetirizine dihydrochloride tablet
5 mg / antihistamine

10 TABLETS

24 HOUR RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

UPC FP0

LOT:

EXP:

The trade dress of this Xyzal[®] package is subject to trademark protection.

Dist. By: Chatten, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2017

Origin: Switzerland

50122365

Xyzal Allergy 24 Hour Tablet Blister

1.03.17

Xyzal_Allergy 10ct_PeelPush Blister_2016_r1_CLEAN



Xyzal Allergy 24 Hour Tablet Bottle Carton

01.06.17

Xyzal_Allergy_35ct_CTN_2016_r3_CLEAN



CLINICALLY PROVEN RELIEF
IN 60 MINUTES

CLINICALLY PROVEN RELIEF
FOR 24 HOURS

UPC
FPO

Drug Facts

Active ingredient (in each tablet)
Levocetirizine dihydrochloride 5 mg.....Antihistamine

Purpose

Uses
temporarily relieves these symptoms due to hay fever or other respiratory allergies:
 ■ runny nose ■ itchy, watery eyes
 ■ sneezing ■ itching of the nose or throat

Warnings

Do not use
 ■ if you have kidney disease
 ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have
 ■ ever had trouble urinating or emptying your bladder

When using this product
 ■ drowsiness may occur
 ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if
 ■ you have trouble urinating or emptying your bladder
 ■ an allergic reaction to this product occurs. Seek medical help right away.

Drug Facts (continued)

If pregnant or breast-feeding:
 ■ if breast-feeding: not recommended
 ■ if pregnant: ask a health professional before use

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

adults 65 years of age and older

■ ask a doctor

adults and children 12-64 years of age

■ take 1 tablet (5 mg) once daily in the evening
 ■ do not take more than 1 tablet (5 mg) in 24 hours
 ■ ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms

children 6-11 years of age

■ take ½ tablet (2.5 mg) once daily in the evening
 ■ do not take more than ½ tablet (2.5 mg) in 24 hours

children under 6 years of age

■ do not use

consumers with kidney disease

■ do not use

Drug Facts (continued)

Other information
 ■ store between 20° and 25°C (68° and 77°F)
 ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients
colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide

Questions or comments?
call 1-800-633-1610

Read directions and warnings before use. Keep this carton. It has important information.

The trade dress of this Xyzal® package is subject to trademark protection.

Dist. By: Chatterin, Inc., a Sareff Company, Chattanooga, TN 37409-0219

©2017 Origin: Switzerland

0100066-01

NDC 41167-3510-1

ORIGINAL
PRESCRIPTION STRENGTH

XYZAL
ALLERGY 24HR

levocetirizine dihydrochloride tablet
5 mg / antihistamine

**24 HOUR
RELIEF OF**

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

**35
TABLETS**

ORIGINAL
PRESCRIPTION STRENGTH



XYZAL
ALLERGY 24HR

Clinically Proven 24 Hour Relief of
Indoor and Outdoor Allergies

Xyzal Allergy 24 Hour Tablet Bottle Label

01.06.17


Xyzal_Allergy_35ct_Label_2016_r5_CLEAN



Xyzal Allergy 24 Hour Tablet Bottle Bonus Carton

01.06.17

Xyzal_Allergy_45ct_Bonus_CTN_2016_r4_CLEAN



CLINICALLY PROVEN RELIEF
IN 60 MINUTES

CLINICALLY PROVEN RELIEF
FOR 24 HOURS

Drug Facts
Active ingredient (in each tablet) Purpose
Levocetirizine dihydrochloride 5 mg.....Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other respiratory allergies:
 ■ runny nose ■ itchy, watery eyes
 ■ sneezing ■ itching of the nose or throat

Warnings
Do not use
 ■ if you have kidney disease
 ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have
 ■ ever had trouble urinating or emptying your bladder

When using this product
 ■ drowsiness may occur
 ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if
 ■ you have trouble urinating or emptying your bladder
 ■ an allergic reaction to this product occurs. Seek medical help right away.

Drug Facts (continued)
If pregnant or breast-feeding:
 ■ if breast-feeding: not recommended
 ■ if pregnant: ask a health professional before use

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

| | |
|--|--|
| adults 65 years of age and older | ask a doctor |
| adults and children 12-64 years of age | <ul style="list-style-type: none"> take 1 tablet (5 mg) once daily in the evening do not take more than 1 tablet (5 mg) in 24 hours ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms |
| children 6-11 years of age | <ul style="list-style-type: none"> take ½ tablet (2.5 mg) once daily in the evening do not take more than ½ tablet (2.5 mg) in 24 hours |
| children under 6 years of age | do not use |
| consumers with kidney disease | do not use |

Drug Facts (continued)
Other information
 ■ store between 20° and 25°C (68° and 77°F)
 ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients
colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide

Questions or comments?
call 1-800-633-1610

Read directions and warnings before use. Keep this carton. It has important information.

The trade dress of this Xyzal® package is subject to trademark protection.
Dist. By: Chatterin, Inc., a Sandoz Company, Chattanooga, TN 37409-0219
©2017 Origin: Switzerland
XXXXXXXX-XX

BONUS! 10 TABLETS FREE

ORIGINAL PRESCRIPTION STRENGTH

XYZAL
ALLERGY 24HR

levocetirizine dihydrochloride tablet
5 mg / antihistamine

45
TABLETS

24 HOUR RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

ORIGINAL PRESCRIPTION STRENGTH

XYZAL
ALLERGY 24HR

Clinically Proven 24 Hour Relief of
Indoor and Outdoor Allergies




UPC
FPO

LOT:
EXP:

Xyzal Allergy 24 Hour Tablet Bottle Label

01.06.17

Xyzal_Allergy_45ct_Label_2016_r4_CLEAN



Xyzal Allergy 24 Hour Tablet Bottle Carton

01.06.17

Xyzal_Allergy_55ct_CTN_2016_r3_CLEAN



CLINICALLY PROVEN RELIEF
IN 60 MINUTES

CLINICALLY PROVEN RELIEF
FOR 24 HOURS

UPC
FPO

LOT:
EXP:

Drug Facts
Active ingredient (in each tablet) Purpose
Levocetirizine dihydrochloride 5 mg.....Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other respiratory allergies:
 ■ runny nose ■ itchy, watery eyes
 ■ sneezing ■ itching of the nose or throat

Warnings
Do not use
 ■ if you have kidney disease
 ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have
 ■ ever had trouble urinating or emptying your bladder

When using this product
 ■ drowsiness may occur
 ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if
 ■ you have trouble urinating or emptying your bladder
 ■ an allergic reaction to this product occurs. Seek medical help right away.

Drug Facts (continued)
If pregnant or breast-feeding:
 ■ if breast-feeding: not recommended
 ■ if pregnant: ask a health professional before use

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

| | |
|--|--|
| adults 65 years of age and older | ■ ask a doctor |
| adults and children 12-64 years of age | ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours |
| children 6-11 years of age | ■ ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms ■ do not take more than ½ tablet (2.5 mg) in 24 hours |
| children under 6 years of age | ■ do not use |
| consumers with kidney disease | ■ do not use |

Drug Facts (continued)
Other information
 ■ store between 20° and 25°C (68° and 77°F)
 ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients
colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide

Questions or comments?
call 1-800-633-1610

Read directions and warnings before use. Keep this carton. It has important information.

The trade dress of this Xyzal® package is subject to trademark protection.
Dist. By: Chatterin, Inc., a Sareff Company, Chattanooga, TN 37409-0219
©2017 Origin: Switzerland
0100067-01

NDC 41167-3510-2

ORIGINAL
PRESCRIPTION STRENGTH

XYZAL®


ALLERGY 24HR

levocetirizine dihydrochloride tablet
5 mg / antihistamine

55
TABLETS

24 HOUR RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES






Clinically Proven 24 Hour Relief of
Indoor and Outdoor Allergies

Xyzal Allergy 24 Hour Tablet Bottle Label

01.06.17

Xyzal_Allergy_55ct_Label_2016_r4_CLEAN



Xyzal Allergy 24 Hour Tablet Bottle Carton

01.06.17

Xyzal_Allergy_80ct_CTN_2016_r3_CLEAN



CLINICALLY PROVEN RELIEF
IN 60 MINUTES

CLINICALLY PROVEN RELIEF
FOR 24 HOURS

Drug Facts

Active ingredient (in each tablet) Purpose
Levocetirizine dihydrochloride 5 mg.....Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other respiratory allergies:
 ■ runny nose ■ itchy, watery eyes
 ■ sneezing ■ itching of the nose or throat

Warnings
Do not use
 ■ if you have kidney disease
 ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have
 ■ ever had trouble urinating or emptying your bladder

When using this product
 ■ drowsiness may occur
 ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if
 ■ you have trouble urinating or emptying your bladder
 ■ an allergic reaction to this product occurs. Seek medical help right away.

Drug Facts (continued)
If pregnant or breast-feeding:
 ■ if breast-feeding: not recommended
 ■ if pregnant: ask a health professional before use

Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

| | |
|--|--|
| adults 65 years of age and older | ■ ask a doctor |
| adults and children 12-64 years of age | ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours |
| children 6-11 years of age | ■ ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms ■ do not take more than ½ tablet (2.5 mg) in 24 hours |
| children under 6 years of age | ■ do not use |
| consumers with kidney disease | ■ do not use |

Drug Facts (continued)
Other information
 ■ store between 20° and 25°C (68° and 77°F)
 ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients
 colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide

Questions or comments?
 call 1-800-633-1610

Read directions and warnings before use. Keep this carton. It has important information.

The trade dress of this Xyzal® package is subject to trademark protection.
 Dist. By: Chatterin, Inc., a Sandoz Company, Chattanooga, TN 37409-0219
 ©2017 Origin: Switzerland
 0100068-01

NDC 41167-3510-3

ORIGINAL PRESCRIPTION STRENGTH

XYZAL
ALLERGY 24HR

levocetirizine dihydrochloride tablet
5 mg / antihistamine

80 TABLETS

24 HOUR RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

ORIGINAL PRESCRIPTION STRENGTH

XYZAL
ALLERGY 24HR

Clinically Proven 24 Hour Relief of
Indoor and Outdoor Allergies




UPC
FPO

LOT:
EXP:

Xyzal Allergy 24 Hour Tablet Bottle Label

01.06.17

Xyzal_Allergy_80oct_Label_2016_r4_CLEAN



Xyzal Allergy 24 Hour Club Card

01.06.17

Xyzal_Allergy_110ct_ClubCard_2016_r3_CLEAN

NDC 41167-3510-4

ORIGINAL
PRESCRIPTION STRENGTH

XYZAL[®]

ALLERGY 24HR

levocetirizine dihydrochloride tablet
5 mg / antihistamine

24 HOUR RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

110 TABLETS

0051550-01

LOT
EXP

XYZAL[®]

ALLERGY 24HR

CLINICALLY PROVEN RELIEF IN 60 MINUTES

CLINICALLY PROVEN RELIEF FOR 24 HOURS

| Drug Facts | | Drug Facts (continued) | |
|---|--|---|--|
| Active ingredient (in each tablet) Levocetirizine dihydrochloride 5 mg..... | | Purpose Antihistamine | |
| Uses temporarily relieves these symptoms due to hay fever or other respiratory allergies: ■ runny nose ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat | | Directions adults 65 years of age and older ■ ask a doctor adults and children 12-64 years of age ■ take 1 tablet (5 mg) once daily in the evening ■ do not take more than 1 tablet (5 mg) in 24 hours ■ ½ tablet (2.5 mg) once daily in the evening may be appropriate for less severe symptoms | |
| Warnings Do not use ■ if you have kidney disease ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine | | children 6-11 years of age ■ take ½ tablet (2.5 mg) once daily in the evening ■ do not take more than ½ tablet (2.5 mg) in 24 hours | |
| Ask a doctor before use if you have ■ ever had trouble urinating or emptying your bladder | | children under 6 years of age ■ do not use | |
| When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery | | consumers with kidney disease ■ do not use | |
| Stop use and ask a doctor if ■ you have trouble urinating or emptying your bladder ■ an allergic reaction to this product occurs. Seek medical help right away. | | Other information ■ store between 20° and 25°C (68° and 77°F) ■ safety sealed: do not use if sealed package was torn or opened, or if printed foil inner seal on bottle is torn or missing | |
| If pregnant or breast-feeding: ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use | | Inactive ingredients colloidal anhydrous silica, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, titanium dioxide | |
| Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). | | Questions or comments? call 1-800-633-1610 | |

Read directions and warnings before use. Keep this carton. It has important information.

The trade dress of this Xyzal[®] package is subject to trademark protection.
Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
©2017 Origin: Switzerland

0051552-01

DATA MATRIX FPO

UPC FPO

Xyzal Allergy 24 Hour Tablet Bottle Induction Seal

1.22.16

Xyzal_Allergy Tablet_Bottle Safety Seal_2016

SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION ----- 5 point Helvetica
SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION Neue Bold Cond Italic
SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION
SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION
SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION
SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION
SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION

Children's Xyzal Allergy 24 Hour Oral Solution Carton

1.04.17

Xyzal_Childrens_Allergy 5oz_CANADA_CTN_2016_r3_CLEAN

Drug Facts (continued)

Directions
 ■ use only with enclosed dosing cup

| | |
|--|---|
| adults 65 years of age and older | ■ ask a doctor |
| adults and children 12-64 years of age | ■ take 5 mL or 10 mL once daily in the evening depending upon severity of symptoms ■ do not take more than 10 mL in 24 hours |
| children 6-11 years of age | ■ take 5 mL once daily in the evening ■ do not take more than 5 mL in 24 hours |
| children 2-5 years of age | ■ take 2.5 mL once daily in the evening ■ do not take more than 2.5 mL in 24 hours |
| children under 2 years of age | ■ do not use |
| consumers with kidney disease | ■ do not use |

Note: mL=milliliters

Other information

- each 5 mL contains: sodium 3 mg
- store between 20° and 25°C (68° and 77°F)
- safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients
 flavor, glacial acetic acid, glycerin, maltitol solution, methylparaben, propylparaben, purified water, saccharin sodium, sodium acetate trihydrate

Questions or comments?
 call 1-800-633-1610

The trade dress of this Xyzal® package is subject to trademark protection.
 Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219
 ©2017 Origin: Canada

NDC 41167-3530-1

CLINICALLY PROVEN RELIEF FOR 24 HOURS

CLINICALLY PROVEN RELIEF FROM INDOOR AND OUTDOOR ALLERGIES

Children's XYZAL ALLERGY 24HR

Children's XYZAL ALLERGY 24HR
 levocetirizine dihydrochloride oral solution
 2.5 mg / 5mL antihistamine

Dye Free - Alcohol Free

24 HOUR RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

Use Only With Enclosed Dosing Cup
 Wash and let air dry after each use

5 fl. oz. (148 mL)
 Tutti Frutti Flavor

Drug Facts

| Active ingredient (in each 5 mL) | Purpose |
|--|---------------|
| Levocetirizine dihydrochloride 2.5 mg..... | Antihistamine |

Uses
 temporarily relieves these symptoms due to hay fever or other respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use

- if you have kidney disease
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine

Ask a doctor before use if you have

- ever had trouble urinating or emptying your bladder

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you have trouble urinating or emptying your bladder
- an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use

Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Read directions and warnings before use. Keep this carton. It has important information.

0100070-01

UPC FPO

Area for LOT & EXP

Children's Xyzal Allergy 24 Hour Oral Solution Label

1.04.17

Xyzal_Childrens_Allergy_5oz_CANADA_Label_2016_r3_CLEAN

NDC 41167-3530-1

Ages 2 Years & Older

Children's XYZAL[®] ALLERGY 24HR
levocetirizine dihydrochloride oral solution
2.5 mg/5mL antihistamine

24 HOUR RELIEF OF

- SNEEZING
- RUNNY NOSE
- ITCHY NOSE OR THROAT
- ITCHY, WATERY EYES

Tutti Frutti Flavor 5 fl. oz. (148 mL)

Active ingredient (in each 5 mL)
Levocetirizine dihydrochloride 2.5 mg.....Antihistamine

Purpose
Levocetirizine dihydrochloride 2.5 mg.....Antihistamine
Uses temporarily relieves these symptoms due to hay fever or other respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

Warnings Do not use ■ if you have kidney disease ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing cetirizine Ask a doctor before use if you have ■ ever had trouble urinating or emptying your bladder When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery Stop use and ask a doctor if ■ you have trouble urinating or emptying your bladder ■ an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding: ■ if breast-feeding: not recommended ■ if pregnant: ask a health professional before use Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ use only with enclosed dosing cup adults 65 years of age and older ■ ask a doctor adults and children 12-64 years of age ■ take 5 mL or 10 mL once daily in the evening depending upon severity of symptoms ■ do not take more than 10 mL in 24 hours children 6-11 years of age ■ take 5 mL once daily in the evening ■ do not take more than 5 mL in 24 hours children 2-5 years of age ■ take 2.5 mL once daily in the evening ■ do not take more than 2.5 mL in 24 hours children under 2 years of age ■ do not use consumers with kidney disease ■ do not use Note: mL = milliliters

Other information ■ each 5 mL contains: sodium 3 mg ■ store between 20° and 25°C (68° and 77°F) ■ safety sealed: do not use if carton was opened or if printed foil inner seal on bottle is torn or missing

Inactive ingredients flavor, glacial acetic acid, glycerin, maltitol solution, methylparaben, propylparaben, purified water, saccharin sodium, sodium acetate trihydrate

Questions or comments?
call 1-800-633-1610

The trade dress of this Xyzal[®] package is subject to trademark protection.
Dist. By: Chattem, Inc., a Sanofi Company, Chattanooga, TN 37409-0219 ©2017
Origin: Canada 0100144-01


DATA MATRIX FPO

Lot:
Exp.:

Xyzal Allergy 24 Hour Oral Solution Dosing Cup

1.03.17

Xyzal_Allergy_Dosing Cup_2016_r2_CLEAN



Detailed rendering of dosing cup on next page

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

Xyzal Allergy 24 Hour Oral Solution Dosing Cup
Digital Rendering
1.03.17



Xyzal Allergy 24 Hour Oral Solution Induction Seal

1.22.16

Xyzal_Allergy Liquid_Bottle Safety Seal_2016

SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION ----- 5 point Helvetica
SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION Neue Bold Cond Italic

SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION

SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION

SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION

SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION

SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION SEALD for YOUR PROTECTION