

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Drug Facts (continued)
Questions or comments? 1-800-719-9580
 Visit us at www.meltech.com or call 1-800-719-9580

Made in Israel
 Manufactured by:
Dexcel Pharma Technologies Ltd
 10 Hakdima Street, Yokneam 2089200, Israel

Drug Facts

Active ingredient (in each tablet)
 Omeprazole 20 mg

Purpose
 Acid reducer

Use
 ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:
 ■ Trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ Heartburn with **lightheadedness, sweating or dizziness**
 ■ Chest or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness

Ask a doctor before use if you have:
 ■ Had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ Frequent wheezing, particularly with heartburn
 ■ Unexplained weight loss
 ■ Nausea or vomiting
 ■ Stomach pain

Ask a doctor or pharmacist before use if you are taking:
 ■ Warfarin, clopidogrel or clostazol (blood-thinning medicines)
 ■ Prescription antifungal or anti-yeast medicines
 ■ Digoxin (heart medicine)
 ■ Dexamethasone (steroid medicine)
 ■ Calcium or magnesium supplements (immune system medicines)
 ■ Prescription antibiotics (medicines for HIV infection)
 ■ Methotrexate (arthritis medicine)

Stop use and ask a doctor if:
 ■ Your heartburn continues or worsens
 ■ You need to take this product for more than 14 days
 ■ You need to take more than 1 course of treatment every 4 months
 ■ You get diarrhea

Other information
 ■ Store at 20-25°C (68-77°F); keep product out of high heat and moisture
 ■ Keep the carton. It contains important information.
 ■ Read the directions and warnings before use

Inactive ingredients
 amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolone, ferric oxide, flavor, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium steryl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Directions
 ■ For adults 18 years of age and older
 ■ This product is to be used once a day (every 24 hours), every day for 14 days
 ■ It may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ Take 1 tablet before eating in the morning

Do not crush or chew tablets
 ■ Place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.

Take every day for 14 days
 ■ Do not take more than 1 tablet a day
 ■ Do not use for more than 14 days unless directed by your doctor
 ■ Do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ You may repeat 14-day course every 4 months
 ■ Do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Location for lot number and expiration date

1Y554 FA C3-AK

NOT FOR SALE

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Safety Feature – Do not use if printed tablet blister unit is open or torn.

Read Drug Facts label on back before trying NDC 64239-XXX-07

FREE SAMPLE INSIDE

NEW DOSAGE FORM!

MELTech™
 Melts In Your Mouth

Compare to Prilosec OTC®*

FDA Approved

Omeprazole
 Delayed Release
 Orally Disintegrating Tablets 20 mg
 Acid Reducer

Treats Frequent Heartburn!
 Occurring 2 or More Days a Week

2 TABLETS

Follow samples with a 14-day course of treatment

Melts in your mouth
 Dissolves without water

Strawberry Flavor

Orally Disintegrating Tablets

Actual size



Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Drug Facts (continued)

Stop use and ask a doctor if:

- you get diarrhea
- you need to take more than 1 course of treatment every 4 months
- you do not take this product for more than 14 days
- your heartburn continues or worsens

Warnings

Do not use if you have:

- stomach pain
- nausea or vomiting
- unexplained weight loss
- frequent wheezing, particularly with heartburn
- had heartburn over 3 months. This may be a sign of a more serious condition.

Do not use if you have:

- chest or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- heartburn with lightheadedness, sweating or dizziness
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools

Warnings

Acid reducer

Use

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Do not use if you are allergic to omeprazole

Do not use if you have:

- frequent chest pain
- stomach pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking:

- warfarin, clopidogrel or clostazolol (blood-thinning medicines)
- prescription antifungals (medicines for HIV infection)
- digoxin (heart medicine)
- diazepam (anxiety medicine)
- prescription antifungal or anti-yeast medicines

Ask a doctor before use if you are taking:

- methotrexate (arthritis medicine)

Other Information

- keep the carton, it contains important information.
- store at 20-25°C (68-77°F); keep product out of high heat and moisture

Other Information

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- you may repeat a 14-day course every 4 months
- do not take this medicine with alcohol
- do not use for more than 14 days unless directed by your doctor
- do not take more than 1 tablet a day
- take every day for 14 days
- The tablets can also be swallowed whole with water.
- Place the tablet on tongue; tablet disintegrates, with or without water.
- do not crush or chew tablets
- take 1 tablet before eating in the morning

14-Day Course of Treatment

- symptoms within 24 hours
- if may take 1 to 4 days for full effect; some people get complete relief of

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Active ingredient (in each tablet)

Purpose

Ingredients

Other Ingredients

cellyl alcohol, colloidal silicon dioxide, croscopolone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize meal, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Drug Facts (continued)

Questions or comments? 1-800-719-9386

Weekdays 7:30 AM to 5:00 PM EST

Made in Israel

Manufactured by:
Cescael Pharma Technologies Ltd
 10 Hakhina Street, Yokneam 2069200, Israel

Location for lot number and expiration date

1Y574 FA C3



This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC 64239-XXX-01

MELTech™
Melts In Your Mouth

Compare to Prilosec OTC®*

NEW DOSAGE FORM!

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer



Strawberry Flavor

Treats Frequent Heartburn!
Occurring 2 or More Days a Week

14 TABLETS

ONE 14-DAY COURSE OF TREATMENT

Melts in your mouth

Dissolves without water



Actual size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Drug Facts (continued)
 Questions or comments? 1-800-719-9380.
 Weekdays 7:30 AM to 5:00 PM EST

Made in Israel
 Manufactured by:
 Dexel Pharma Technologies Ltd
 10 Hakhdma Street, Yoknein 2099200, Israel

Drug Facts (continued)

Active ingredient (in each tablet)
 Omeprazole 20 mg, Acid reducer

Purpose
 ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest or shoulder pain with shortness of breath; sweating; pain spreading to arm, neck or shoulders; or lightheadedness

requent chest pain
 These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.

Ask a doctor or pharmacist before use if you are taking:
 ■ warfarin, clopidogrel or clostazol (blood-thinning medicines)
 ■ prescription antifungal or anti-yeast medicines
 ■ dazepam (anxiety medicine)
 ■ digoxin (heart medicine)
 ■ calcium or myophosphate (medicines for HIV infection)
 ■ prescription antidiarrheals (medicines for HIV infection)

Stop use and ask a doctor if: ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea

■ you develop a rash or joint pain
 ■ methoxyacetate (antibiotics medicine)

Other information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize starch, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Directions
 ■ for adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning

do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.

take every day for 14 days
 ■ do not take more than 1 tablet a day
 ■ do not use for more than 14 days unless directed by your doctor

do not take this medicine with alcohol
 ■ do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months

do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture

Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscopolone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize starch, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Location for lot number and expiration date

1Y574 FA C3-K



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC 64239-XXX-01

MELTech™
 Melts In Your Mouth

Compare to Prilosec OTC®*

NEW DOSAGE FORM!

Omeprazole
 Delayed Release
 Orally Disintegrating Tablets 20 mg
 Acid Reducer

Strawberry Flavor

Treats Frequent Heartburn!
 Occurring 2 or More Days a Week

14 TABLETS ONE 14-DAY COURSE OF TREATMENT

Melts in your mouth
 Dissolves without water

Orally Disintegrating Tablets

Actual size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Drug Facts (continued)
Questions or comments? 1-800-719-9280
 weekdays 7:30AM to 5:00 PM EST

Made in Israel
 Manufactured by:
Chelexel Pharma Technologies Ltd
 10 Herodias Street, Yokneam 2069200, Israel

<p>Drug Facts (continued)</p> <p>Warnings Allergy alert: do not use if you are allergic to omeprazole</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with shortness of breath, sweating, pain spreading to arm, neck or shoulders, or lightheadedness ■ frequent chest pain ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ Ask a doctor before use if you have: ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are taking: ■ warfarin, clopidogrel or ciprofloxacin (blood-thinning medicines) ■ prescription antifungal or anti-yeast medicines ■ dicyclanil (antxiety medicine) ■ digoxin (heart medicine) ■ calcium or myophosphate medrel (immune system medicines) ■ prescription antiretrovirals (medicines for HIV infection) ■ methoclopramide (gastrointestinal medicine)</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p>	<p>Drug Facts</p> <p>Active ingredient (in each tablet) Omeprazole 20 mg Acid reducer</p> <p>Purpose ■ Treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: do not use if you are allergic to omeprazole</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with shortness of breath, sweating, pain spreading to arm, neck or shoulders, or lightheadedness ■ frequent chest pain ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ Ask a doctor before use if you have: ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are taking: ■ warfarin, clopidogrel or ciprofloxacin (blood-thinning medicines) ■ prescription antifungal or anti-yeast medicines ■ dicyclanil (antxiety medicine) ■ digoxin (heart medicine) ■ calcium or myophosphate medrel (immune system medicines) ■ prescription antiretrovirals (medicines for HIV infection) ■ methoclopramide (gastrointestinal medicine)</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p>
<p>Drug Facts (continued)</p> <p>Directions ■ For adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>Other information ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture</p> <p>Inactive ingredients cetyl alcohol, colloidal silicon dioxide, croscopolone, ferric oxide, flavor, hypromellose, hydroxypropyl methylcellulose, hypromellose, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p>	<p>Drug Facts (continued)</p> <p>Directions ■ For adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</p> <p>14-Day Course of Treatment ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>Other information ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture</p> <p>Inactive ingredients cetyl alcohol, colloidal silicon dioxide, croscopolone, ferric oxide, flavor, hypromellose, hydroxypropyl methylcellulose, hypromellose, mannitol, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate</p>

Location for lot number and expiration date

1Y574 FA C3-B



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC 64239-XXX-01

MELTech™
 Melts In Your Mouth

Compare to Prilosec OTC®*

Child-resistant

Omeprazole
 Delayed Release
 Orally Disintegrating Tablets 20 mg
 Acid Reducer



Strawberry Flavor

Treats Frequent Heartburn!
 Occurring 2 or More Days a Week

14 TABLETS ONE 14-DAY COURSE OF TREATMENT

Melts in your mouth
 Dissolves without water

Actual size



Omeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg

Drug Facts (continued)

Warnings

- Do not use if you have: trouble or pain swallowing food, vomiting w/ blood, or bloody or black stools
- Do not use if you have: heartburn with nighttime, sweating or dizziness
- Do not use if you have: chest or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or back, dizziness, or lightheadedness
- Do not use if you have: frequent chest pain
- Do not use if you have: frequent heartburn over 3 months. This may be a sign of a more serious condition.
- Do not use if you have: frequent wheezing, particularly w/ heartburn
- Do not use if you have: unexplained weight loss
- Do not use if you have: nausea or vomiting
- Do not use if you have: contact pain

Use

Heartburn (heartburn) occurs 2 or more days a week

Heartburn is a moderate relief of heartburn; this drug may take 1 to 4 days for full effect

Directions

For adults 18 years of age and older

Take 1 tablet before eating in the morning

Do not crush or chew tablets

The tablets can also be swallowed whole with water.

Take every day for 14 days

Do not take more than 1 tablet a day

Do not use for more than 14 days unless directed by your doctor

Repeat 14-day courses (if needed)

You may repeat a 14-day course every 4 months


Do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- Keep the carton and warnings before use
- Keep the carton, it contains important information
- Store at 20°-25° C (68°-77° F); keep product out of high heat and moisture

1 Y955 FA C3-B



0 000000 000000 0


Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –
Do not use if printed tablet blister unit is open or torn.

NDC 64239-XXX-03


NEW DOSAGE FORM!



Melts In Your Mouth

Compare to Prilosec OTC*

Child-resistant



Strawberry
Flavor

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer


Treats Frequent Heartburn!
Occurring 2 or More Days a Week

42 TABLETS

THREE 14-DAY COURSES
OF TREATMENT

Melts in your mouth

Disolves without water



Actual size

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Drug Facts (continued)
Questions or comments? 1-800-719-9206
 weekdays 7:30 AM to 5:00 PM EST

Made in Israel
 Manufactured by:
Cexcel Pharma Technologies Ltd
 10 Hakdama Street, Yokneam 2069200, Israel

Drug Facts

Active ingredient (in each tablet)
 Omeprazole 20 mg, Acid reducer

Purpose
 Acid reducer

Use
 ■ Treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
 ■ Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness

Treat chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ stomach pain

Ask a doctor before use if you are taking:
 ■ warfarin, clopidogrel or clostazolol (blood-thinning medicines)
 ■ prescription antifungal or anti-yeast medicines
 ■ diazepam (anxiety medicine)
 ■ digoxin (heart medicine)
 ■ ecchinus or myopiplanate methyl (immune system medicines)
 ■ prescription antidiarrheals (medicines for HIV infection)
 ■ methionine (amino acid medicine)

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea
 ■ you develop a rash or joint pain

Other information
 ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture
 ■ keep the carton, it contains important information
 ■ read the directions and warnings before use

Inactive ingredients
 cetyl alcohol, colloidal silicon dioxide, croscarmellose, ferrous oxide, flavor, hypromellose, hydroxypropylcellulose, polyethylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Directions
 ■ For adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
 ■ take 1 tablet before eating in the morning
 ■ do not crush or chew tablets
 ■ place the tablet on tongue; tablet disintegrates, with or without water.

Repeated 14-Day Course (if needed)
 ■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more than every 4 months

do not take for more than 14 days or more than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)

Location for lot number and expiration date

1Y574 FA C3-A



This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC.

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC 64239-XXX-01

MELTech™
 Melts In Your Mouth

Compare to Prilosec OTC®*

FDA Approved

Omeprazole
 Delayed Release
 Orally Disintegrating Tablets 20 mg
 Acid Reducer

Strawberry Flavor

Treats **Frequent Heartburn!**
 Occurring 2 or More Days a Week

14 TABLETS ONE 14-DAY COURSE OF TREATMENT

Melts in your mouth
 Dissolves without water

Actual size

Omeprazole
 Delayed Release
 Orally Disintegrating Tablets 20 mg

11930 FA C3-C

Drug Facts (continued)
Active ingredient (in each tablet)
 Omeprazole 20 mg, Acid reducer

Use
 Treats frequent heartburn (occurs 2 or more days a week) for full effect.

Warnings
 Allergy alert: do not use if you are allergic to omeprazole.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- requires chest pain

 These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- require wheezing, particularly with heartburn
- unexplained weight loss
- stomach pain

Ask a doctor or pharmacist before use if you are taking:

- warfarin, clopidogrel or clostazol (blood-thinning medicines)
- prescription antifungal or anti-worm medicines
- digoxin (heart medicine)
- baclofen or mycophenolate mofetil (immune system medicines)
- prescription antiretroviral medicines for HIV infection)
- methotrexate (arthritis medicine)

Drug Facts (continued)
Inactive ingredients amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, croscarmellose, ferric oxide, flavor, hypromellose, hydroxypropylmethylcellulose, magnesium stearate, mannitol, microcrystalline cellulose, polyethylene glycol, sodium bicarbonate, sodium lauryl sulfate, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments? 1-800-719-9266; weekdays 7:30 AM to 5:00 PM EST

Manufactured by:
 Dexcel Pharma Technologies Ltd
 10 Haidimna Street, Yokneam 2069200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC™.

Directions
 Take 1 to 4 tablets 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours.

1-4 Day Course of Treatment
 Take 1 tablet before eating in the morning.

Do not crush or chew tablets
 The tablets can also be swallowed whole with water.

Repeat 1-4 Day Course (if needed)
 Do not take more than 1 tablet a day.

Other information
 Children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you get diarrhea
- you get dizziness

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature –
 Do not use if printed tablet blister unit is open or torn.

NDC 642239-XX-02

NEW DOSAGE FORM!

Compare to Prilosec OTC®*

Buy One 14 Count, Get One 14 Count Free!

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Strawberry
Flavor

Treats Frequent Heartburn!
Occurring 2 or More Days a Week

28 TABLETS

TWO 14-DAY COURSES
OF TREATMENT

Melts in your mouth

Dissolves without water

Actual size

Omeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg

11980 FA C3-F

Drug Facts (continued)

Active Ingredient (in each tablet)
Omeprazole 20 mg; Acid reducer

Purpose
Acid reducer

Use
● treats frequent heartburn (occurs 2 or more days a week)
● not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:
● heartburn w/ thin highheadness, sweating or dizziness
● chest or shoulder pain with shortness of breath, sweating; pain spreading to arm, neck or shoulder or highheadness
● frequent chest pain
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
● had heartburn over 3 months. This may be a sign of a more serious condition.
● frequent wheezing, particularly with heartburn
● unexplained weight loss
● nausea or vomit

Ask a doctor or pharmacist before use if you are taking:
● warfarin, clopidogrel or clozapine (blood-thinning medicines)
● prescription antifungal or anti-yeast medicines
● digoxin (heart medicine)
● bacitrimin or mupirocin (immune system medicines)
● prescription antiretroviral (medicines for HIV infection)
● metformin (diabetes medicine)

Other information
● keep the carton. It contains important information.
● read the Medication and Warnings before use.
● store at 20-25°C (68-77°F); keep product out of high heat and moisture.

Directions
● for adults 18 years of age and older
● this product is to be used once a day (every 24 hours), every day for 14 days
● if may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
● take 1 tablet before eating in the morning
● do not crush or chew tablets
● place the tablet on tongue; tablet disintegrates, with or without water.
The tablets can also be swallowed whole with water.

14-Day Course (if needed)
● take every day for 14 days
● do not use for more than 14 days unless directed by your doctor
● do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
● do not take for more than 14 days or more often than every 4 months
● you may repeat 14-day course every 4 months

Other information
● children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Stop use and ask a doctor if:
● your heartburn continues or worsens
● you need to take more than 1 course of treatment every 4 months

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Use if pregnant or breast-feeding. Ask a health professional before use.
● you get diarrhea ● you develop a rash or joint pain

Directions
● for adults 18 years of age and older
● this product is to be used once a day (every 24 hours), every day for 14 days
● if may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment
● take 1 tablet before eating in the morning
● do not crush or chew tablets
● place the tablet on tongue; tablet disintegrates, with or without water.
The tablets can also be swallowed whole with water.

14-Day Course (if needed)
● take every day for 14 days
● do not use for more than 14 days unless directed by your doctor
● do not take this medicine with alcohol

Repeated 14-Day Course (if needed)
● do not take for more than 14 days or more often than every 4 months
● you may repeat 14-day course every 4 months

Other information
● children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Drug Facts (continued)

Active ingredients
Omeprazole 20 mg; Acid reducer

Warnings
Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:
● heartburn w/ thin highheadness, sweating or dizziness
● chest or shoulder pain with shortness of breath, sweating; pain spreading to arm, neck or shoulder or highheadness
● frequent chest pain
These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
● had heartburn over 3 months. This may be a sign of a more serious condition.
● frequent wheezing, particularly with heartburn
● unexplained weight loss
● nausea or vomit

Ask a doctor or pharmacist before use if you are taking:
● warfarin, clopidogrel or clozapine (blood-thinning medicines)
● prescription antifungal or anti-yeast medicines
● digoxin (heart medicine)
● bacitrimin or mupirocin (immune system medicines)
● prescription antiretroviral (medicines for HIV infection)
● metformin (diabetes medicine)

Other information
● keep the carton. It contains important information.
● read the Medication and Warnings before use.
● store at 20-25°C (68-77°F); keep product out of high heat and moisture.

Questions or comments? 1-800-719-9200; weekdays 7:30 AM to 5:00 PM EST

Manufactured by:
Dexel Pharma Technologies Ltd
10 Hahadina Street, Yokneam 209200, Israel

This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC™.

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Safety Feature – Do not use if printed tablet blister unit is open or torn.

NDC 64239-63-XX-02

NEW DOSAGE FORM!

Compare to Prilosec OTC®*

Bonus Size!
14 Tablets Free

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Treats Frequent Heartburn!
Occurring 2 or More Days a Week

28 TABLETS TWO 14-DAY COURSES OF TREATMENT

Melts in your mouth
Dissolves without water

Actual size

Strawberry Flavor

Reference ID: 4288423

need off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!
On This Purchase

MELTech™
Melts In Your Mouth

Compare to Prilosec OTC®*
NDC 64239-XXX-01

NEW DOSAGE FORM!

Omeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Treats Frequent Heartburn!
Occurring 2 or More Days a Week

14 TABLETS ONE 14-DAY COURSE OF TREATMENT

Melts in your mouth
Dissolves without water

Strawberry Flavor

Actual size

1Y974 FA CP2

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX} on your purchase of
14 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Orally Disintegrating Tablets 20 mg

Omeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg

14 TABLETS

CONSUMER: Only one coupon is redeemable per purchase. RETAILER: This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. Do Not Double. Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001

COUPON NOT SUBJECT TO DOUBLING
Customer pays any sales tax.



*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX
SAVE \$X NOW!
 On This Purchase



MELTech™
Melts In Your Mouth

Compare to Prilosec OTC®*
 NDC 64239-XXX-02

NEW DOSAGE FORM!

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Treats **Frequent Heartburn!**
Occurring **2 or More Days a Week**

28

TABLETS

TWO 14-DAY COURSES
OF TREATMENT

Melts in your mouth

Dissolves without water



Strawber***
Flavor



Actual
size

1Y930 FA CP2

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX}

on your purchase of
28 COUNT SIZE
"Store Brand" Omeprazole
Delayed Release Orally Disintegrating Tablets 20 mg

Omeprazole

Delayed Release
Orally Disintegrating Tablets 20 mg

28 TABLETS

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. **Do Not Double.** Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Perrigo Company (#1565), NCH Marketing, PO Box 880001, El Paso, TX 88588-0001

COUPON NOT SUBJECT TO DOUBLING

Customer pays any sales tax.



0 00000 00000 0

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

peel off coupon

Instant Redeemable Coupon - Expires XX-XX-XX

SAVE \$X NOW!
 On This Purchase

MELTech™
 Melts In Your Mouth

Compare to Prilosec OTC®*
 NDC 64239-XXX-03

NEW DOSAGE FORM!

Omeprazole
 Delayed Release
 Orally Disintegrating Tablets 20 mg
 Acid Reducer

Treats Frequent Heartburn!
 Occurring 2 or More Days a Week

42 TABLETS THREE 14-DAY COURSES
 OF TREATMENT

Melts in your mouth
 Dissolves without water

Strawberry
 Flavor

Actual size

1Y955 FA CP2

SAVE \$X NOW!

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XXXX

SAVE \$X^{XX} on your purchase of
 42 COUNT SIZE
 "Store Brand" Omeprazole
 Delayed Release Orally Disintegrating Tablets 20 mg

CONSUMER: Only one coupon is redeemable per purchase. **RETAILER:** This coupon will be redeemable for face value on specified products (plus a handling fee) if submitted in compliance with the terms of this offer. Cash value 1/100th of one cent. Limit one coupon per customer. **Do Not Double.** Coupon is non-reproducible. Coupons cannot be combined. Mail properly handled coupons to: L. Penigo Company (#1565), WCH Marketing, PO Box 880001, El Paso, TX 88568-0001

COUPON NOT SUBJECT TO DOUBLING
 Customer pays any sales tax.

Omeprazole
 Delayed Release
 Orally Disintegrating Tablets 20 mg

42 TABLETS

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

KAREN M MAHONEY
07/12/2018