

Rx only

400 mg daily dose
(two 200 mg tablets)

Contents: 3 individual weekly blister packs. Each blister pack contains 14 tablets (200 mg per tablet).

42 Film-coated tablets

Kisqali®
(ribociclib)
tablets

Physician Sample - Not For Sale

Usual Dosage: Take two 200 mg tablets at the same time once daily, with or without food. Swallow tablets whole.

DO NOT chew, crush, or split tablets.

- Take Kisqali® exactly as directed by your prescriber.
- Use the dose calendar located on the back of each weekly pack to keep track of your treatment.

Store refrigerated at 2°C to 8°C (36°F to 46°F). Excursions permitted between 2°C and 15°C (36°F to 59°F). After dispensing, patients may store at room temperature 20°C to 25°C (68°F to 77°F) for up to 2 months.

Store tablets in the original blister pack.

Each tablet contains 200 mg ribociclib equivalent to 254.40 mg of the succinate salt of ribociclib.

Keep all drugs out of the reach of children.

For more information, visit www.KISQALI.com or call 1-844-KISQALI (1-844-547-7254)

Product of Slovenia.

Manufactured by:
Novartis d.o.o.
Ljubljana, Slovenia

Distributed by:
Novartis Pharmaceuticals Corp.
East Hanover, NJ 07936
©Novartis

0078-0867-93

Discard after / /

Kisqali®
(ribociclib)
tablets

400 mg
daily dose
(two 200 mg tablets)

42 Film-coated tablets

Physician Sample - Not For Sale

ATTENTION:
Store refrigerated until
dispensed to patient

Rx only

Kisqali®
(ribociclib)
tablets

400 mg daily dose
(two 200 mg tablets)

Contents: 3 individual weekly blister packs.
Each blister pack contains 14 tablets (200 mg per tablet).

Storage: Store refrigerated at 2°C to 8°C (36°F to 46°F). Excursions permitted between 2°C and 15°C (36°F to 59°F). After dispensing, patients may store at room temperature 20°C to 25°C (68°F to 77°F) for up to 2 months.

42 Film-coated tablets

NOVARTIS

199x45x84_L


400 mg
daily dose
(two 200 mg tablets)

Kisqali®
(ribociclib)
tablets

42 Film-coated tablets

46359570 US





Contents: 3 individual weekly blister packs. Each blister pack contains 21 tablets (200 mg per tablet).

600 mg daily dose
(three 200 mg tablets)

Rx only

Physician Sample - Not For Sale

Kisqali[®]
(ribociclib)
tablets

63 Film-coated tablets

Usual Dosage: Take three 200 mg tablets at the same time once daily, with or without food. Swallow tablets whole.

DO NOT chew, crush, or split tablets.

- Take Kisqali[®] exactly as directed by your prescriber.
- Use the dose calendar located on the back of each weekly pack to keep track of your treatment.

Store refrigerated at 2°C to 8°C (36°F to 46°F). Excursions permitted between 2°C and 15°C (36°F to 59°F). After dispensing, patients may store at room temperature 20°C to 25°C (68°F to 77°F) for up to 2 months.

Store tablets in the original blister pack.

Each tablet contains 200 mg ribociclib equivalent to 254.40 mg of the succinate salt of ribociclib.

Keep all drugs out of the reach of children.

For more information, visit www.KISQALI.com or call 1-844-KISQALI (1-844-547-7254)

Product of China.

0078-9874-63

Discard after

___/___/___

Manufactured by:
Novartis d.o.o.
Ljubljana, Slovenia

Distributed by:
Novartis Pharmaceuticals Corp.
East Hanover, NJ 07936
©Novartis

Kisqali[®]
(ribociclib)
tablets

600 mg
daily dose
(three 200 mg tablets)

63 Film-coated tablets

NOVARTIS

Kisqali[®]
(ribociclib)
tablets

63 Film-coated tablets

ATTENTION:
Store refrigerated until
dispensed to patient

Physician Sample - Not For Sale

Rx only

600 mg daily dose
(three 200 mg tablets)

Contents: 3 individual weekly blister packs.
Each blister pack contains 21 tablets (200 mg per tablet).

Storage: Store refrigerated at 2°C to 8°C (36°F to 46°F). Excursions permitted between 2°C and 15°C (36°F to 59°F). After dispensing, patients may store at room temperature 20°C to 25°C (68°F to 77°F) for up to 2 months.

199x45x84_I


600 mg
daily dose
(three 200 mg tablets)

63 Film-coated tablets

Kisqali[®]
(ribociclib)
tablets

46359567 US





Rx only

Physician Sample - Not For Sale

600 mg daily dose
(three 200 mg tablets)

Contents: 3 individual weekly blister packs. Each blister pack contains 21 tablets (200 mg per tablet).

63 Film-coated tablets

Kisqali[®]
(ribociclib)
tablets

46359568 US

Usual Dosage: Take three 200 mg tablets at the same time once daily, with or without food. Swallow tablets whole. **DO NOT** chew, crush, or split tablets.

- Take Kisqali[®] exactly as directed by your prescriber.
- Use the dose calendar located on the back of each weekly pack to keep track of your treatment.

Store refrigerated at 2°C to 8°C (36°F to 46°F). Excursions permitted between 2°C and 15°C (36°F to 59°F). After dispensing, patients may store at room temperature 20°C to 25°C (68°F to 77°F) for up to 2 months.

Store tablets in the original blister pack.

Each tablet contains 200 mg ribociclib equivalent to 254.40 mg of the succinate salt of ribociclib.

Keep all drugs out of the reach of children.

For more information, visit www.KISQALI.com or call 1-844-KISQALI (1-844-547-7254)

Product of Slovenia.

Manufactured by:
Novartis d.o.o.
Ljubljana, Slovenia

Distributed by:
Novartis Pharmaceuticals Corp.
East Hanover, NJ 07936
©Novartis

0078-9874-63

Discard after

___/___/___

Kisqali[®]
(ribociclib)
tablets

600 mg
daily dose
(three 200 mg tablets)

63 Film-coated tablets

199x45x84_J

ATTENTION:
Store refrigerated until
dispensed to patient

Physician Sample - Not For Sale

Rx only

Kisqali[®]
(ribociclib)
tablets

600 mg daily dose
(three 200 mg tablets)

Contents: 3 individual weekly blister packs.
Each blister pack contains 21 tablets (200 mg per tablet).

Storage: Store refrigerated at 2°C to 8°C (36°F to 46°F). Excursions permitted between 2°C and 15°C (36°F to 59°F). After dispensing, patients may store at room temperature 20°C to 25°C (68°F to 77°F) for up to 2 months.

63 Film-coated tablets

NOVARTIS

600 mg
daily dose
(three 200 mg tablets)

Kisqali[®]
(ribociclib)
tablets

63 Film-coated tablets



Physician Sample – Not For Sale

Rx only

28-Day Supply

Kisqali® Femara® CO-PACK

(ribociclib tablets; letrozole tablets)

600 mg daily dose
(three 200 mg tablets)

63 Film-coated tablets

2.5 mg per tablet

28 Film-coated tablets

Storage: Store refrigerated at 2°C to 8°C (36°F to 46°F). Excursions permitted between 2°C and 15°C (36°F to 59°F). After dispensing, patients may store at room temperature 20°C to 25°C (68°F to 77°F) for up to 2 months.

ATTENTION:
Store refrigerated until dispensed to patient



Variable data printing

LOT XXXXXX

EXP YYYY-MMM-DD

B S 009002

Kisqali® Femara® CO-PACK
(ribociclib tablets; letrozole tablets)

600 mg daily dose
(three 200 mg tablets)

63 Film-coated tablets

2.5 mg per tablet

28 Film-coated tablets



Kisqali®
(ribociclib) tablets
Product of China.
Manufactured by:
Novartis d.o.o.,
Ljubljana, 1000, Slovenia

Femara®
(letrozole) tablets
Product of France.
Manufactured by:
Novartis Pharma Stein
Stein, Switzerland

0078-9923-61

Distributed by:
Novartis Pharmaceuticals Corp.
East Hanover, NJ 07936

©Novartis

Rx only

Physician Sample – Not For Sale

This carton contains:
• Kisqali® (ribociclib) tablets: 3 individual weekly blister packs.
Each blister pack contains 21 tablets (200 mg per tablet).
• Femara® (letrozole) tablets: 1 bottle containing 28 tablets.
(0078-9881-50)
(0078-9902-21)
• Prescribing Information for Kisqali® Femara® CO-PACK.
• Dosing Calendar

Kisqali® Femara® CO-PACK
(ribociclib tablets; letrozole tablets)

2.5 mg per tablet

600 mg daily dose
(three 200 mg tablets)

28 Film-coated tablets

Usual Dosage: Take three 200 mg Kisqali® tablets once daily for 3 weeks followed by a one-week break, and one 2.5 mg Femara® tablet once daily for 28 days. Take with or without food. Swallow tablets whole.

DO NOT chew, crush, or split tablets.

- Take Kisqali® Femara® CO-PACK exactly as directed by your prescriber.
- Use the dosing calendar inside to keep track of your 28-day treatment.

Store refrigerated at 2°C to 8°C (36°F to 46°F). Excursions permitted between 2°C and 15°C (36°F to 59°F).

After dispensing, patients may store at room temperature 20°C to 25°C (68°F to 77°F) for up to 2 months. Store tablets in the original blister pack.

Keep all drugs out of the reach of children. For more information, visit www.KISQALI.com or call 1-844-KISQALI (1-844-547-7254)

Each Kisqali® tablet contains 200 mg ribociclib equivalent to 254.40 mg of the succinate salt of ribociclib. Each Femara® tablet contains 2.5 mg letrozole.

Discard after

__/__/__

Kisqali® Femara® CO-PACK
(ribociclib tablets; letrozole tablets)

600 mg daily dose
(three 200 mg tablets)

63 Film-coated tablets

2.5 mg per tablet

28 Film-coated tablets

247x47x85



Physician Sample – Not For Sale

Rx only

28-Day Supply

Kisqali® Femara® CO-PACK

(ribociclib tablets; letrozole tablets)

600 mg daily dose
(three 200 mg tablets)

63 Film-coated tablets

2.5 mg per tablet

28 Film-coated tablets

Storage: Store refrigerated at 2°C to 8°C (36°F to 46°F). Excursions permitted between 2°C and 15°C (36°F to 59°F). After dispensing, patients may store at room temperature 20°C to 25°C (68°F to 77°F) for up to 2 months.

ATTENTION:
Store refrigerated until dispensed to patient



Variable data printing

LOT XXXXXX

EXP YYYY-MMM-DD

200600 B S 009002

Kisqali® Femara® CO-PACK
(ribociclib tablets; letrozole tablets)

600 mg daily dose
(three 200 mg tablets)

63 Film-coated tablets

2.5 mg per tablet

28 Film-coated tablets



Kisqali®
(ribociclib) tablets
Product of Slovenia.
Manufactured by:
Novartis d.o.o.,
Ljubljana, 1000, Slovenia

Femara®
(letrozole) tablets
Product of France.
Manufactured by:
Novartis Pharma Stein
Stein, Switzerland

0078-9923-61

Distributed by:
Novartis Pharmaceuticals Corp.
East Hanover, NJ 07936

©Novartis

Rx only

Physician Sample – Not For Sale

This carton contains:
• Kisqali® (ribociclib) tablets: 3 individual weekly blister packs.
Each blister pack contains 21 tablets (200 mg per tablet).
• Femara® (letrozole) tablets: 1 bottle containing 28 tablets.
(0078-9881-50)
(0078-9902-21)
• Prescribing Information for Kisqali® Femara® CO-PACK.
• Dosing Calendar

Kisqali® Femara® CO-PACK

(ribociclib tablets; letrozole tablets)

600 mg daily dose
(three 200 mg tablets)

63 Film-coated tablets

2.5 mg per tablet

28 Film-coated tablets

Usual Dosage: Take three 200 mg Kisqali® tablets once daily for 3 weeks followed by a one-week break, and one 2.5 mg Femara® tablet once daily for 28 days. Take with or without food. Swallow tablets whole.

DO NOT chew, crush, or split tablets.

- Take Kisqali® Femara® CO-PACK exactly as directed by your prescriber.
- Use the dosing calendar inside to keep track of your 28-day treatment.

Store refrigerated at 2°C to 8°C (36°F to 46°F). Excursions permitted between 2°C and 15°C (36°F to 59°F).

After dispensing, patients may store at room temperature 20°C to 25°C (68°F to 77°F) for up to 2 months. Store tablets in the original blister pack.

Keep all drugs out of the reach of children. For more information, visit www.KISQALI.com or call 1-844-KISQALI (1-844-547-7254)

Each Kisqali® tablet contains 200 mg ribociclib equivalent to 254.40 mg of the succinate salt of ribociclib. Each Femara® tablet contains 2.5 mg letrozole.

Discard after

Kisqali® Femara® CO-PACK
(ribociclib tablets; letrozole tablets)

600 mg daily dose
(three 200 mg tablets)

63 Film-coated tablets

2.5 mg per tablet

28 Film-coated tablets

247x47x85

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

CHRISTY L OSGOOD
02/11/2025 01:07:19 PM