


BACK




SEAL AREA FOR LOT & EXP  
25 mm x 4.5 mm

AREA FOR LOT AND EXP  
TO BE PRINTED IN DARK COLOR  
26 mm x 5.5 mm


000067291

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt






PANEL 1




PANEL 2




PANEL 3



PANEL 4




BASE PANEL




000067292

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



UNVARNISHED AREA  
FOR LOT & EXP. DATE



000067293

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

PANEL 2

<p><b>Drug Facts (continued)</b></p> <p><b>NSAID allergy alert:</b> ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> <li>hives</li> <li>facial swelling</li> <li>asthma (wheezing)</li> <li>shock</li> <li>skin reddening</li> <li>rash</li> <li>blisters</li> </ul> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>NSAID stomach bleeding warning:</b></p> <p>This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>are age 60 or older</li> <li>have had stomach ulcers or bleeding problems</li> <li>take a blood thinning (anticoagulant) or steroid drug</li> <li>take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li> <li>have 3 or more alcoholic drinks every day while using this product</li> <li>take more or for a longer time than directed</li> </ul> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p>
---	--

PANEL 3

<p><b>Drug Facts (continued)</b></p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist</li> <li>if you have ever had an allergic reaction to acetaminophen or any other pain reliever</li> <li>right before or after heart surgery</li> </ul> <p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>you have liver disease</li> <li>stomach bleeding warning applies to you</li> <li>you have problems or serious side effects from taking pain relievers</li> <li>you have a history of stomach problems, such as heartburn</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke</li> <li>you are taking a diuretic</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>under a doctor's care for any serious condition</li> <li>taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin</li> <li>taking any other drug</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>take with food or milk if stomach upset occurs</li> </ul>
---	---

PANEL 4

<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> <li>feel faint</li> <li>vomit blood</li> <li>have bloody or black stools</li> <li>have stomach pain that does not get better</li> </ul> </li> <li>you have symptoms of heart problems or stroke: <ul style="list-style-type: none"> <li>chest pain</li> <li>trouble breathing</li> <li>weakness in one part or side of body</li> <li>slurred speech</li> <li>leg swelling</li> <li>pain gets worse or lasts more than 10 days</li> </ul> </li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>redness or swelling is present in the painful area</li> <li>any new symptoms appear</li> </ul> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>
---	--

PANEL 1



<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each caplet)</b></p> <p>Acetaminophen 250 mg ..... Pain reliever</p> <p>Ibuprofen 125 mg (NSAID*) ..... Pain reliever *nonsteroidal anti-inflammatory drug</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> <li>headache</li> <li>toothache</li> <li>backache</li> <li>menstrual cramps</li> <li>muscular aches</li> <li>minor pain of arthritis</li> </ul> </li> </ul>	<p><b>Drug Facts (continued)</b></p> <p><b>Warnings</b></p> <p><b>Acetaminophen liver damage warning:</b></p> <p>This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> <li>with other drugs containing acetaminophen</li> <li>more than 6 caplets in 24 hours, which is the maximum daily amount for this product</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> <p><b>Acetaminophen allergy alert:</b> may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>rash</li> </ul> <p>If skin reaction occurs, stop use and seek medical help right away.</p>
--	--

<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not take more than directed</li> </ul> <table border="1"> <tr> <td>adults and children 12 years and over</td> <td>take 2 caplets every 8 hours while symptoms persist</td> </tr> <tr> <td>children under 12 years</td> <td>ask a doctor</td> </tr> </table> <ul style="list-style-type: none"> <li>do not take more than 6 caplets in 24 hours, unless directed by a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>read all warnings and directions before use.</li> <li>store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F)</li> </ul> <p><b>Inactive ingredients</b> carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide</p> <p><b>Questions or comments?</b> call weekdays 9 AM to 5 PM at 1-800-88-ADVIL</p> <p>000067296</p>	adults and children 12 years and over	take 2 caplets every 8 hours while symptoms persist	children under 12 years	ask a doctor
adults and children 12 years and over	take 2 caplets every 8 hours while symptoms persist			
children under 12 years	ask a doctor			

BASE PANEL

000067296

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces




LOT & EXP AREA

000067295

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces




BASE PANEL



PANEL #4

PANEL #1



PANEL #2

PANEL #3

00067298

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	N/A
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	N/A
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP  
PANEL 1

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ if you have ever had an allergic reaction to acetaminophen or any other pain reliever</li> <li>■ right before or after heart surgery</li> </ul> <p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>■ you have liver disease</li> <li>■ stomach bleeding warning applies to you</li> <li>■ you have problems or serious side effects from taking pain relievers</li> <li>■ you have a history of stomach problems, such as heartburn</li> <li>■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke</li> <li>■ you are taking a diuretic</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>■ under a doctor's care for any serious condition</li> <li>■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin</li> <li>■ taking any other drug</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>■ take with food or milk if stomach upset occurs</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ you experience any of the following signs of stomach bleeding:             <ul style="list-style-type: none"> <li>■ feel faint</li> <li>■ vomit blood</li> <li>■ have bloody or black stools</li> <li>■ have stomach pain that does not get better</li> </ul> </li> </ul>
---	--

000070384  
LIFT HERE

BACK  
PANEL 2

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ you have symptoms of heart problems or stroke:             <ul style="list-style-type: none"> <li>■ chest pain</li> <li>■ trouble breathing</li> </ul> </li> <li>■ weakness in one part or side of body</li> <li>■ slurred speech</li> <li>■ leg swelling</li> <li>■ pain gets worse or lasts more than 10 days</li> <li>■ redness or swelling is present in the painful area</li> <li>■ any new symptoms appear</li> </ul> <p><b>If pregnant or breast-feeding</b>, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <p><b>do not take more than directed</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="padding: 2px;">adults and children 12 years and over</td> <td style="padding: 2px;">■ take 2 caplets every 8 hours while symptoms persist</td> </tr> <tr> <td style="padding: 2px;">children under 12 years</td> <td style="padding: 2px;">■ ask a doctor</td> </tr> </table> <p>■ do not take more than 6 caplets in 24 hours, unless directed by a doctor</p>	adults and children 12 years and over	■ take 2 caplets every 8 hours while symptoms persist	children under 12 years	■ ask a doctor
adults and children 12 years and over	■ take 2 caplets every 8 hours while symptoms persist				
children under 12 years	■ ask a doctor				

BASE  
PANEL 3

<p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ read all warnings and directions before use. Keep carton.</li> <li>■ store at 20-25°C (68-77°F)</li> <li>■ avoid excessive heat above 40°C (104°F)</li> </ul> <p><b>Inactive ingredients</b> carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide</p> <p><b>Questions or comments?</b> call weekdays 9 AM to 5 PM at 1-800-88-ADVIL</p>	
---	--

000070384

	TYPE SIZE
• DRUG FACTS TITLE	N/A
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



TOP  
PANEL 1

FOR LOT & EXPIRATION AREA.  
UNVARNISHED AND PRINTED  
IN BLACK OR DARK BLUE.

BASE  
PANEL 3

RELEASED BACK  
PANEL 2

000067299

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



TOP  
PANEL 1

FOR LOT & EXPIRATION AREA. UNVARNISHED AND PRINTED IN BLACK OR DARK BLUE.

BASE  
PANEL 3

RELEASED BACK  
PANEL 2

000070389

	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP  
PANEL 1

**CONTAINS 2 MEDICINES**  
Acetaminophen + Ibuprofen

**Advil**  
**DUAL ACTION**  
WITH Acetaminophen

Acetaminophen 250 mg and  
Ibuprofen (NSAID) 125 mg Tablets  
Pain Reliever

**180**  
Caplets\*  
\*Capsule-Shaped Tablets

Do Not Use if seal under bottle cap imprinted with  
"SEALED for YOUR PROTECTION" is broken or missing.

Do not use with other medicines containing  
**ACETAMINOPHEN**; can cause liver damage.

**Drug Facts**

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Ibuprofen 125 mg (NSAID)*	Pain reliever *nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - muscular aches
  - minor pain of arthritis

000070390

LIFT HERE  
for More Drug Facts

81% For Position Only

FOR LOT & EXPIRATION AREA,  
UNVARNISHED AND PRINTED  
IN BLACK OR DARK BLUE.

BASE  
PANEL 3

<p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>under a doctor's care for any serious condition</li> <li>taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin</li> <li>taking any other drug</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>take with food or milk if stomach upset occurs</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>you experience any of the following signs of stomach bleeding:                     <ul style="list-style-type: none"> <li>feel faint</li> <li>vomit blood</li> <li>have bloody or black stools</li> <li>have stomach pain that does not get better</li> </ul> </li> <li>you have symptoms of heart problems or stroke:                     <ul style="list-style-type: none"> <li>chest pain</li> <li>trouble breathing</li> <li>weakness in one part or side of body</li> <li>slurred speech</li> <li>leg swelling</li> </ul> </li> <li>pain gets worse or lasts more than 10 days</li> <li>redness or swelling is present in the painful area</li> <li>any new symptoms appear</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p><b>If pregnant or breast-feeding</b>, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not take more than directed</li> </ul> <table border="1"> <tr> <td>adults and children 12 years and over</td> <td>take 2 caplets every 8 hours while symptoms persist</td> </tr> <tr> <td>children under 12 years</td> <td>ask a doctor</td> </tr> </table> <ul style="list-style-type: none"> <li>do not take more than 6 caplets in 24 hours, unless directed by a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>read all warnings and directions before use.</li> <li>store at 20-25°C (68-77°F)</li> <li>avoid excessive heat above 40°C (104°F)</li> </ul> <p><b>Inactive ingredients</b> carnauba wax, colloidal silicon dioxide, croscarmellose sodium, ferric oxides, glyceryl dibehenate, hypromellose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, titanium dioxide</p> <p><b>Questions or comments?</b> call weekdays 9 AM to 5 PM at 1-800-88-ADVIL</p> <p>For most recent product information, visit <a href="http://www.Advil.com">www.Advil.com</a> For US Patent or Application status see <a href="http://www.productpats.com">www.productpats.com</a></p> <p>Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or licensor Trademarks owned or licensed by GSK</p>	adults and children 12 years and over	take 2 caplets every 8 hours while symptoms persist	children under 12 years	ask a doctor
adults and children 12 years and over	take 2 caplets every 8 hours while symptoms persist				
children under 12 years	ask a doctor				

RELEASED BACK  
PANEL 2

<p><b>Drug Facts (continued)</b></p> <p><b>Warnings</b></p> <p><b>Acetaminophen liver damage warning:</b></p> <p>This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> <li>with other drugs containing acetaminophen</li> <li>more than 6 caplets in 24 hours, which is the maximum daily amount for this product</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> <p><b>Acetaminophen allergy alert:</b> may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>rash</li> </ul> <p>If skin reaction occurs, stop use and seek medical help right away.</p> <p><b>NSAID allergy alert:</b> ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> <li>hives</li> <li>facial swelling</li> <li>asthma (wheezing)</li> <li>shock</li> <li>skin reddening</li> <li>rash</li> <li>blisters</li> </ul> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>NSAID stomach bleeding warning:</b></p> <p>This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> <li>are age 60 or older</li> <li>have had stomach ulcers or bleeding problems</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>take a blood thinning (anticoagulant) or steroid drug</li> <li>take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]</li> <li>have 3 or more alcoholic drinks every day while using this product</li> <li>take more or for a longer time than directed</li> </ul> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription).</li> <li>if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist</li> <li>if you have ever had an allergic reaction to acetaminophen or any other pain reliever</li> <li>right before or after heart surgery</li> </ul> <p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>you have liver disease</li> <li>stomach bleeding warning applies to you</li> <li>you have problems or serious side effects from taking pain relievers</li> <li>you have a history of stomach problems, such as heartburn</li> <li>you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke</li> <li>you are taking a diuretic</li> </ul>
---	---

000070390

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



TOP  
PANEL 1

BASE  
PANEL 3

RELEASED BACK  
PANEL 2

000075078

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



**Drug Facts (continued)**

**When using this product:**

- Take with food or milk if stomach upset occurs.
- Stop use and ask a doctor if:
  - You experience any of the following signs of stomach bleeding:
    - feel faint
    - womit blood
    - have bloody or black stools
    - have stomach pain that does not get better
    - you have symptoms of heart problems or stroke:
      - weakness in one part or side of body
      - chest pain
      - trouble breathing
  - Pain gets worse or lasts more than 10 days
  - stirred speech
  - leg swelling
  - padres or swelling is present in the painful area

If pregnant or breast-feeding, ask a health professional before use. It is especially important to use ibuprofen at 20 weeks or later in pregnancy, unless directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for overdose.

**Questions or comments?** Call weekdays 9 AM to 5 PM at 1-800-88-ADVIL.

**Drug Facts (continued)**

**Directions:**

- adults and children: take 2 caplets every 8 hours while symptoms persist
- 12 years and over: ask a doctor
- children under 12 years: ask a doctor

**Do not take more than directed.**

**Other information:**

- read all warnings and directions before use. keep carton, store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Active ingredients (in each caplet):**

- acetaminophen, ask a doctor or pharmacist if you have ever had an allergic reaction to acetaminophen or other pain reliever
- if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- have had stomach ulcers or bleeding problems
- are age 65 or older

**Warnings:**

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - muscular aches
  - minor pain of arthritis

**Uses:**

- ibuprofen 125 mg (NSAID), Pain reliever
- acetaminophen 250 mg, Pain reliever

**Drug Facts (continued)**

**Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.**

**Read and keep carton for complete warnings and information.**



**Do Not Use:** If seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

For most recent product information, visit [www.Advil.com](http://www.Advil.com). For US Patent or Application status see [www.products.com](http://www.products.com). Dist. by: GSK, CK, Warren, NJ 07099

©2021 GSK or licensor. Trademarks owned or licensed by GSK. 000070388

100% For Position Only

3 0573 0147 85 9

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)

000070388

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP LAYER FACE

UNVARNISHED & PRINTED AREA FOR LOT & EXP. (LINES DO NOT PRINT)

TOP LAYER BACK

BASE LAYER

adults and children 12 years and over	take 2 caplets every 8 hours while symptoms persist
children under 12 years	ask a doctor

000070392


NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP LAYER FACE




UNVARNISHED & PRINTED AREA FOR LOT & EXP. (LINES DO NOT PRINT)

TOP LAYER BACK



BASE LAYER



000070391

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

VALERIE S PRATT  
01/25/2022 02:36:23 PM