This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Reference ID: 4813202

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda



This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Drug Facts (continued)

Other information
- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

Inactive ingredients
benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?
1-800-317-2165
(Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.
The Astepro mark is owned by Viatris. Used under license.
© 2021 Bayer.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981
Patents: patents.livewell.bayer.com
Product of Germany.
000000 Bayer

INDOOR & OUTDOOR
Allergy Relief

NDC 0280-0065-05

ASTEPRO® ALLERGY

First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

SAMPLE - NOT FOR SALE

NEW! ANTIHISTAMINE NASAL SPRAY

ASTEPRO® ALLERGY

Azelastine Hydrochloride
ANTIHISTAMINE NASAL SPRAY

FULL PRESCRIPTION STRENGTH

STEROID FREE

UP TO 24 HOUR RELIEF OF
- Nasal congestion
- Runny nose
- Sneezing
- Itchy nose

22 METERED SPRAYS
0.14 fl oz (4 mL)

Alcohol-Free
Fragrance-Free

THIS WAY TO RELIEF

Drug Facts

Active ingredient (in each spray) — ***Purpose***
Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine) ..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- nasal congestion
- runny nose
- sneezing
- itchy nose

Warnings
Only for use in the nose. Do not spray in eyes or mouth.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you
- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

When using this product
- drowsiness may occur
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if
- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- Read the User Guide for how to:
  - prime the bottle before first use
  - prime bottle again if not used for 3 or more days
  - use the spray
  - clean the spray nozzle if it gets clogged

adults and children 12 years and older	This product may be used either once or twice a day: ■ once daily: use 2 sprays in each nostril; **OR** ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours ■ **do not use more than 4 sprays** in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ **do not use more than 2 sprays** in each nostril in a 24 hour period
children under 6 years	do not use

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

THIS WAY TO RELIEF

ASTEPRO® ALLERGY

First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

ASTEPRO® ALLERGY

Azelast 2 .

ANTIHISTAMINE NASAL SPRAY

NEW! ANTIHISTAMINE NASAL SPRAY

FULL PRESCRIPTION STRENGTH

STEROID FREE

UP TO 24 HOUR RELIEF OF

- Nasal congestion
- Runny nose
- Sneezing
- Itchy nose

60 METERED SPRAYS
0.37 fl oz (11 mL)

Alcohol-Free
Fragrance-Free

INDOOR & OUTDOOR
Allergy Relief

NDC 0280-0065-01

Drug Facts (continued)

Other information
- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

Inactive ingredients
benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?
1-800-317-2165
(Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.
The Astepro mark is owned by Viatris. Used under license.
© 2021 Bayer.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981

Patents: patents.livewell.bayer.com

Product of Germany.

000000 Bayer

Drug Facts

Active ingredient (in each spray) — ***Purpose***
Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine) ..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- nasal congestion
- runny nose
- sneezing
- itchy nose

Warnings
Only for use in the nose. Do not spray in eyes or mouth.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you
- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

When using this product
- drowsiness may occur
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if
- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- Read the User Guide for how to:
  - prime the bottle before first use
  - prime bottle again if not used for 3 or more days
  - use the spray
  - clean the spray nozzle if it gets clogged

adults and children 12 years and older	This product may be used either once or twice a day: ■ once daily: use 2 sprays in each nostril; **OR** ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours ■ **do not use more than 4 sprays** in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ **do not use more than 2 sprays** in each nostril in a 24 hour period
children under 6 years	do not use

0 00000 00000 0

LO

Reference ID: 4813202

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda



This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

THIS WAY TO RELIEF

ASTEPRO® ALLERGY

First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

NEW! ANTIHISTAMINE NASAL SPRAY

ASTEPRO® ALLERGY

Azelastine HCl 0.1%
ANTIHISTAMINE NASAL SPRAY

FULL PRESCRIPTION STRENGTH

STEROID FREE

UP TO 24 HOUR RELIEF OF

- Nasal congestion
- Runny nose
- Sneezing
- Itchy nose

INDOOR & OUTDOOR Allergy Relief

BAYER

120 METERED SPRAYS
0.78 fl oz (23 mL)

Alcohol-Free
Fragrance-Free

Drug Facts (continued)

Other information
- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

Inactive ingredients
benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?
1-800-317-2165
(Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.
The Astepro mark is owned by Viatris. Used under license.
© 2021 Bayer.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981

Patents: patents.livewell.bayer.com

Product of Germany.

000000

Bayer

NDC 0280-0065-02

0 00000 00000 0

LOT:
EXP: YYYY-MM

Drug Facts

Active ingredient (in each spray)	***Purpose***
Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)	Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- nasal congestion
- runny nose
- sneezing
- itchy nose

Warnings
Only for use in the nose. Do not spray in eyes or mouth.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you
- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

When using this product
- drowsiness may occur
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if
- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- Read the User Guide for how to:
  - prime the bottle before first use
  - prime bottle again if not used for 3 or more days
  - use the spray
  - clean the spray nozzle if it gets clogged

adults and children 12 years and older	This product may be used either once or twice a day: - once daily: use 2 sprays in each nostril; **OR** - twice daily: use 1 or 2 sprays in each nostril every 12 hours - **do not use more than 4 sprays** in each nostril in a 24 hour period
children 6 years to 11 years	- an adult should supervise use - 1 spray in each nostril every 12 hours - **do not use more than 2 sprays** in each nostril in a 24 hour period
children under 6 years	do not use

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

THIS WAY TO RELIEF

ASTEPRO® ALLERGY

First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

Drug Facts (continued)

Other information
- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

Inactive ingredients
benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?
1-800-317-2165
(Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.
The Astepro mark is owned by Viatris. Used under license.
© 2021 Bayer.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981
Patents: patents.livewell.bayer.com
Product of Germany.
000000

Bayer

INDOOR & OUTDOOR
Allergy Relief

NDC 0280-0065-04

ASTEPRO® ALLERGY

Azelastine HCl
ANTIHISTAMINE NASAL SPRAY

NEW! ANTIHISTAMINE NASAL SPRAY

FULL PRESCRIPTION STRENGTH

STEROID FREE

UP TO 24 HOUR RELIEF OF
- Nasal congestion
- Runny nose
- Sneezing
- Itchy nose

BAYER

200 METERED SPRAYS
1.01 fl oz (30 mL)

Alcohol-Free
Fragrance-Free

0 00000 00000 0

LOT:
EXP: YYYY-MM

Drug Facts

Active ingredient (in each spray) — ***Purpose***
Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine) ..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- nasal congestion
- runny nose
- sneezing
- itchy nose

Warnings
Only for use in the nose. Do not spray in eyes or mouth.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you
- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

When using this product
- drowsiness may occur
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if
- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- Read the User Guide for how to:
  - prime the bottle before first use
  - prime bottle again if not used for 3 or more days
  - use the spray
  - clean the spray nozzle if it gets clogged

adults and children 12 years and older	This product may be used either once or twice a day: ■ once daily: use 2 sprays in each nostril; **OR** ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours ■ **do not use more than 4 sprays** in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ **do not use more than 2 sprays** in each nostril in a 24 hour period
children under 6 years	do not use

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

THIS WAY TO RELIEF

Children's
ASTEPRO® ALLERGY
First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

NEW! ANTIHISTAMINE NASAL SPRAY

Children's
ASTEPRO®
ALLERGY
Azelastine HCl 205.5 mcg per spray
ANTIHISTAMINE NASAL SPRAY

FULL PRESCRIPTION STRENGTH

UP TO 24 HOUR RELIEF OF
- Nasal congestion
- Runny nose
- Sneezing
- Itchy nose

60 METERED SPRAYS
0.37 fl oz (11 mL)

Alcohol-Free
Fragrance-Free

AGE 6+

INDOOR & OUTDOOR
Allergy Relief

NDC 0280-0066-01

Drug Facts (continued)

Other information
- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

Inactive ingredients
benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?
1-800-317-2165
(Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.
The Astepro mark is owned by Viatris. Used under license.
© 2021 Bayer.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981

Patents: patents.livewell.bayer.com

Product of Germany.

000000 Bayer

Drug Facts

Active ingredient (in each spray)	***Purpose***
Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)	Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- nasal congestion
- runny nose
- sneezing
- itchy nose

Warnings
Only for use in the nose. Do not spray in eyes or mouth.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you
- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

When using this product
- drowsiness may occur
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if
- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- Read the User Guide for how to:
  - prime the bottle before first use
  - prime bottle again if not used for 3 or more days
  - use the spray
  - clean the spray nozzle if it gets clogged

adults and children 12 years and older	This product may be used either once or twice a day: - once daily: use 2 sprays in each nostril; **OR** - twice daily: use 1 or 2 sprays in each nostril every 12 hours - **do not use more than 4 sprays** in each nostril in a 24 hour period
children 6 years to 11 years	- an adult should supervise use - 1 spray in each nostril every 12 hours - **do not use more than 2 sprays** in each nostril in a 24 hour period
children under 6 years	do not use

LOT :

Reference ID: 4813202

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda



This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

THIS WAY TO RELIEF

Children's
ASTEPRO® ALLERGY
First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

NEW! ANTIHISTAMINE NASAL SPRAY

Children's
ASTEPRO®
ALLERGY
Azelastine HCl 0.1% ANTIHISTAMINE NASAL SPRAY

FULL PRESCRIPTION STRENGTH

STEROID FREE

UP TO
24 HOUR
RELIEF OF
- Nasal congestion
- Runny nose
- Sneezing
- Itchy nose

INDOOR & OUTDOOR
Allergy Relief

BAYER

120 METERED SPRAYS
0.78 fl oz (23 mL)

Alcohol-Free
Fragrance-Free

AGE 6+

NDC 0280-066-02

Drug Facts (continued)

Other information
- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

Inactive ingredients
benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?
1-800-317-2165
(Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.
The Astepro mark is owned by Viatris. Used under license.
© 2021 Bayer.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981
Patents: patents.livewell.bayer.com
Product of Germany.
000000

Bayer

0 00000 00000 0

LOT:
EXP: YYYY-MM

Drug Facts

Active ingredient (in each spray) — ***Purpose***

Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)..... Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- nasal congestion
- runny nose
- sneezing
- itchy nose

Warnings

Only for use in the nose. Do not spray in eyes or mouth.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you
- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

When using this product
- drowsiness may occur
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if
- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- Read the User Guide for how to:
  - prime the bottle before first use
  - prime bottle again if not used for 3 or more days
  - use the spray
  - clean the spray nozzle if it gets clogged

adults and children 12 years and older	This product may be used either once or twice a day: ■ <u>once daily:</u> use 2 sprays in each nostril; **OR** ■ <u>twice daily:</u> use 1 or 2 sprays in each nostril every 12 hours ■ **do not use more than 4 sprays** in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ **do not use more than 2 sprays** in each nostril in a 24 hour period
children under 6 years	do not use

Reference ID: 4813202

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda



This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

THIS WAY TO RELIEF

ASTEPRO® ALLERGY

First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

Drug Facts (continued)

Other information
- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

Inactive ingredients
benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?
1-800-317-2165
(Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.
The Astepro mark is owned by Viatris. Used under license.
© 2021 Bayer.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981

Patents: patents.livewell.bayer.com

Product of Germany.

000000

Bayer

INDOOR & OUTDOOR
Allergy Relief

NDC 0280-0065-09

ASTEPRO® ALLERGY

Azelastine HCl 205.5 mcg per spray

ANTIHISTAMINE NASAL SPRAY

NEW! ANTIHISTAMINE NASAL SPRAY

FULL PRESCRIPTION STRENGTH

UP TO 24 HOUR RELIEF OF
- Nasal congestion
- Runny nose
- Sneezing
- Itchy nose

BAYER

2x120 METERED SPRAY BOTTLES
0.78 fl oz (23 mL) each
1.56 fl oz (46 mL) total

Alcohol-Free
Fragrance-Free

0 00000 00000 0

LOT:
EXP: YYYY-MM

Drug Facts

Active ingredient (in each spray) — ***Purpose***
Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine) ..... Antihistamine

Uses
temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- nasal congestion
- runny nose
- sneezing
- itchy nose

Warnings
Only for use in the nose. Do not spray in eyes or mouth.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you
- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

When using this product
- drowsiness may occur
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if
- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- Read the User Guide for how to:
  - prime the bottle before first use
  - prime bottle again if not used for 3 or more days
  - use the spray
  - clean the spray nozzle if it gets clogged

adults and children 12 years and older	This product may be used either once or twice a day: ■ once daily: use 2 sprays in each nostril; **OR** ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours ■ **do not use more than 4 sprays** in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ **do not use more than 2 sprays** in each nostril in a 24 hour period
children under 6 years	do not use

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

ASTEPRO® ALLERGY

STEROID FREE

First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

Drug Facts

Active ingredient (in each spray) — **Purpose**

Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)..... Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- nasal congestion
- runny nose
- sneezing
- itchy nose

Warnings

Only for use in the nose. Do not spray in eyes or mouth.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

When using this product

- drowsiness may occur
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if

- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- Read the User Guide for how to:
  - prime the bottle before first use
  - use the spray
  - prime bottle again if not used for 3 or more days
  - clean the spray nozzle if it gets clogged

adults and children 12 years and older	This product may be used either once or twice a day: ■ once daily: use 2 sprays in each nostril; **OR** ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours ■ **do not use more than 4 sprays** in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ **do not use more than 2 sprays** in each nostril in a 24 hour period
children under 6 years	do not use

Other information

- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this label and the enclosed User Guide for important information
- do not use if sealed package is torn or opened

Inactive ingredients

benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?

1-800-317-2165 (Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.
The Astepro mark is owned by Viatris. Used under license.
© 2021 Bayer.

Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981
Patents: patents.livewell.bayer.com
Product of Germany.
000000

Bayer

NDC 0280-0065-03

LOT:
EXP: YYYY-MM

0 00000 00000 0

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Actual Size 100%

FDA00342-1

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

ASTEPRO® ALLERGY

FPO

HOW TO START GETTING STEROID-FREE ALLERGY RELIEF RIGHT NOW

BAYER

This User Guide will give you detailed directions on how to prime the bottle, use the spray, and how to clean a clogged nozzle.

Keep this User Guide as it contains important information.

GET THE BOTTLE READY

Remove the blue cap and blue clip before using.

PRIME THE BOTTLE

WHEN TO PRIME

- Before you use Astepro Allergy for the first time
- If you have not used Astepro Allergy for 3 or more days
- After you clean a clogged nozzle

HOW TO PRIME

- Aim bottle away from face.
- Hold bottle as shown and pump until a fine mist appears.
- If a fine mist does not appear after 6 sprays, your nozzle may be clogged. Please see "How to Clean a Clogged Nozzle" section.

HOW TO USE

Important: For use in your nose only

STEP 1
Blow your nose to clear your nostrils.

STEP 2
Tilt your head downward toward your toes.

TILT HEAD DOWNWARD

STEP 3
Hold bottle with thumb under bottle and spray nozzle between fingers.

STEP 4
Close one nostril.

HOLD ONE NOSTRIL SHUT

STEP 5
Put the spray tip about ¼ inch to ½ inch into the other nostril. Hold bottle upright and aim the spray tip toward the back of your nose.

STEP 6
While sniffing gently, press the pump once or twice (according to dosing directions found in Table 1).

Repeat Steps 4, 5 and 6 in other nostril.

STEP 7
Do not tilt your head back immediately after using Astepro Allergy. This will help to keep the medicine from going into your throat, which may cause a bitter taste.

STEP 8
When you finish using your Astepro Allergy, wipe the spray tip with a clean tissue or cloth. Put the clip and cap back on the bottle.

FOR USE IN CHILDREN
An adult should supervise use.
(See Steps 1 through 8).

TURN OVER →

SELECT THE DOSE

DIRECTIONS
Do not use more than directed

TABLE 1

adults & children 12 years and older	This product may be used either once or twice a day: ■ once daily: use 2 sprays in each nostril; **OR** ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours ■ **do not use more than 4 sprays** in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ **do not use more than 2 sprays** in each nostril in a 24 hour period
children under 6 years	do not use

IMPORTANT INFORMATION FOR USE
Remember, when using this product:

- Drowsiness may occur*
  - › Avoid alcoholic drinks.
  - › Alcohol, sedatives, and tranquilizers may increase drowsiness.
  - › Be careful when driving a motor vehicle or operating machinery.

*In clinical trials, drowsiness was observed in less than 4% of patients taking Astepro Allergy.

HOW TO CLEAN A CLOGGED NOZZLE

If you cannot get your Astepro Allergy to spray a fine mist, the nozzle may be clogged. Don't try to unclog the nozzle with a pin or sharp object—that can damage it.

- Unscrew the spray pump unit from the bottle by turning it to the left (counter-clockwise).
- **Keep open bottle out of reach of children.**
- Fill a bowl or container with warm water. Soak only the spray pump unit in the warm water. Pump the nozzle several times while holding it under the water to clear the clog.
- Let the spray pump unit air dry before putting it back on.
- Tightly screw pump unit back on the open bottle by turning clockwise (to the right).
- After cleaning, follow the instructions for priming.

UNSCREW PUMP UNIT

PUMP NOZZLE WHILE HOLDING IT UNDERWATER

QUESTIONS OR COMMENTS?
1-800-317-2165
(Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.

The Astepro mark is owned by Viatris.
Used under license.

© 2021 Bayer.

Distributed by:
Bayer HealthCare LLC
Whippany, New Jersey 07981

Product of Germany.

000000

Bayer

Reference ID: 4813202

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

SELECT THE DOSE

DIRECTIONS

Do not use more than directed

TABLE 1

adults & children 12 years and older	This product may be used either once or twice a day: • once daily: use 2 sprays in each nostril; **OR** • twice daily: use 1 or 2 sprays in each nostril every 12 hours • **do not use more than 4 sprays** in each nostril in a 24 hour period
children 6 years to 11 years	• an adult should supervise use • 1 spray in each nostril every 12 hours • **do not use more than 2 sprays** in each nostril in a 24 hour period
children under 6 years	do not use

IMPORTANT INFORMATION FOR USE

Remember, when using this product:

- Drowsiness may occur*
  - Avoid alcoholic drinks.
  - Alcohol, sedatives, and tranquilizers may increase drowsiness.
  - Be careful when driving a motor vehicle or operating machinery.
- In clinical trials, drowsiness was observed in less than 4% of patients taking Astepro Allergy.

HOW TO CLEAN A CLOGGED NOZZLE

If you cannot get your Astepro Allergy to spray a fine mist, the nozzle may be clogged. Don't try to unclog the nozzle with a pin or sharp object—that can damage it.

- Unscrew the spray pump unit from the bottle by turning it to the left (counter-clockwise).

HOW TO CLEAN A CLOGGED NOZZLE (CONT.)

- **Keep open bottle out of reach of children.**
- Fill a bowl or container with warm water. Soak only the spray pump unit in the warm water. Pump the nozzle several times while holding it under the water to clear the clog.
- Let the spray pump unit air dry before putting it back on.
- Tightly screw pump unit back on the open bottle by turning clockwise (to the right).
- After cleaning, follow the instructions for priming.

QUESTIONS OR COMMENTS?

1-800-317-2165
(Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.

The Astepro mark is owned by Viatris. Used under license.

© 2021 Bayer.

Distributed by:
Bayer Healthcare LLC
Whippany, New Jersey 07981

Product of Germany.

CONSUMER: LIMIT ONE COUPON PER PURCHASE as specified product(s), size(s) and quantity stated. LIMIT OF 4 EXACT SAME COUPONS PER HOUSEHOLD PER DAY. You pay any sales tax. Void if copied, sold, transferred, altered, or auctioned.

RETAILER: You are authorized to act as our agent and redeem this coupon at face value on the specified product(s). We will reimburse you for the face value of this coupon plus 8¢ if submitted in compliance with the Bayer HealthCare LLC Coupon Redemption Policy, incorporated herein by reference. **USE NOT CONSISTENT WITH THESE TERMS MAY CONSTITUTE FRAUD AND MAY VOID ALL COUPONS SUBMITTED.** Send coupons to: Bayer HealthCare LLC 1555 P.O. Box 880001, El Paso, TX 88588-0001. Cash value 1/20th of 1¢.

Good only in the U.S.A. XXXXXX

MANUFACTURER'S COUPON EXPIRES XX/XX/XXXX

Save $X on
(120 spray or higher)

ASTEPRO® ALLERGY

Bayer and the Bayer Cross are registered trademarks of Bayer.
The Astepro mark is owned by Viatris. Used under license.
© 2021 Bayer. Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981

ASTEPRO® ALLERGY

HOW TO START GETTING STEROID-FREE ALLERGY RELIEF RIGHT NOW

This User Guide will give you detailed directions on how to prime the bottle, use the spray, and how to clean a clogged nozzle.

Keep this User Guide as it contains important information.

FPO

GET THE BOTTLE READY

Remove the blue cap and blue clip before using.

PRIME THE BOTTLE

WHEN TO PRIME

- Before you use Astepro Allergy for the first time
- If you have not used Astepro Allergy for 3 or more days
- After you clean a clogged nozzle

HOW TO PRIME

- Aim bottle away from face.
- Hold bottle as shown and pump until a fine mist appears.
- If a fine mist does not appear after 6 sprays, your nozzle may be clogged. Please see "How to Clean a Clogged Nozzle" section.

HOW TO USE

Important: For use in your nose only

STEP 1
Blow your nose to clear your nostrils.

STEP 2
Tilt your head downward toward your toes.

STEP 3
Hold bottle with thumb under bottle and spray nozzle between fingers.

STEP 4
Close one nostril.

STEP 5
Put the spray tip about ¼ inch to ½ inch into the other nostril. Hold bottle upright and aim the spray tip toward the back of your nose.

HOW TO USE (CONT.)

STEP 6
While sniffing gently, press the pump once or twice (according to dosing directions found in Table 1).

Repeat Steps 4, 5 and 6 in other nostril.

STEP 7
Do not tilt your head back immediately after using Astepro Allergy. This will help to keep the medicine from going into your throat, which may cause a bitter taste.

STEP 8
When you finish using your Astepro Allergy, wipe the spray tip with a clean tissue or cloth. Put the clip and cap back on the bottle.

FOR USE IN CHILDREN
An adult should supervise use.

(See Steps 1 through 8).

This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

---

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
06/17/2021 02:41:15 PM