



CHILDREN'S ASTEPRO[®] ALLERGY
 First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

CHILDREN'S ASTEPRO[®] ALLERGY
 Azelastine HCl 205.5 mcg per spray
 ANTIHISTAMINE NASAL SPRAY

FULL PRESCRIPTION STRENGTH

UP TO 24 HOUR RELIEF OF

- ◀ Nasal congestion
- ◀ Runny nose
- ◀ Sneezing
- ◀ Itchy nose

22 METERED SPRAYS
 0.14 fl oz (4mL)

Alcohol-Free
 Fragrance-Free
 AGE 6+



Drug Facts (continued)
Other information
 ■ store between 20° to 25°C (68° to 77°F). Protect from freezing.
 ■ keep this carton and the enclosed User Guide for important information
 ■ do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

Inactive ingredients
 benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?
 1-800-317-2165 (Mon-Fri, 9AM-5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer. The Astepro mark is owned by Viatris. Used under license. © 2023 Bayer. Dist. by: Bayer HealthCare LLC Whippany, NJ 07981 Patents: patents.livewell.bayer.com Product of Germany. 00000000 Bayer

INDOOR & OUTDOOR Allergy Relief

NDC 0280-0066-03



Drug Facts
Active ingredient (in each spray)
 Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine).....
 Purpose
 Antihistamine

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
 Only for use in the nose. Do not spray in eyes or mouth.
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed

When using this product
 ■ drowsiness may occur
 ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery.

Stop use and ask a doctor if
 ■ you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
 ■ nasal discomfort or sneezing may occur right after use
 ■ do not share this bottle with anyone else as this may spread germs

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ Read the User Guide for how to:
 ■ prime the bottle before first use ■ use the spray
 ■ prime bottle again if not used for 3 or more days ■ clean the spray nozzle if it gets clogged

adults and children 12 years and older This product may be used either once or twice a day:
 ■ once daily: use 2 sprays in each nostril; OR
 ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours
 ■ do not use more than 4 sprays in each nostril in a 24 hour period

children 6 years to 11 years ■ an adult should supervise use
 ■ 1 spray in each nostril every 12 hours
 ■ do not use more than 2 sprays in each nostril in a 24 hour period

children under 6 years do not use



CHILDREN'S ASTEPRO[®] ALLERGY
 First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

Drug Facts (continued)

Other information

- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

Inactive ingredients
 benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?
 1-800-317-2165 (Mon-Fri, 9AM-5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer. The Astepro mark is owned by Viatris. Used under license. © 2023 Bayer. Dist. by: Bayer HealthCare LLC Whippany, NJ 07981 Patents: patents.livewell.bayer.com Product of Germany. 00000000 Bayer

INDOOR & OUTDOOR Allergy Relief

NDC 0280-0066-01

CHILDREN'S ASTEPRO[®] ALLERGY
 Azelastine HCl 205.5 mcg per spray
 ANTIHISTAMINE NASAL SPRAY

FULL PRESCRIPTION STRENGTH

UP TO 24 HOUR RELIEF OF

- ◀ Nasal congestion
- ◀ Runny nose
- ◀ Sneezing
- ◀ Itchy nose

STERIOD FREE

60 METERED SPRAYS
 0.37 fl oz (11 mL)

Alcohol-Free **AGE 6+**
Fragrance-Free





Drug Facts

Active ingredient (in each spray)	Purpose
Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)	Antihistamine

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
 Only for use in the nose. Do not spray in eyes or mouth.
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
When using this product
 ■ drowsiness may occur
 ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery.
 ■ you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
 ■ nasal discomfort or sneezing may occur right after use
 ■ do not share this bottle with anyone else as this may spread germs
Stop use and ask a doctor if
 ■ an allergic reaction, such as a skin rash, to this product occurs
 ■ you have severe or frequent nosebleeds
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ Read the User Guide for how to: ■ prime the bottle before first use ■ prime bottle again if not used for 3 or more days adults and children 12 years and older	■ use the spray ■ clean the spray nozzle if it gets clogged This product may be used either once or twice a day: ■ once daily: use 2 sprays in each nostril; OR ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours ■ do not use more than 4 sprays in each nostril in a 24 hour period children 6 years to 11 years ■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ do not use more than 2 sprays in each nostril in a 24 hour period children under 6 years do not use
---	--









CHILDREN'S ASTEPRO ALLERGY
 First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

Drug Facts (continued)
Other information
 ■ store between 20° to 25°C (68° to 77°F). Protect from freezing.
 ■ keep this carton and the enclosed User Guide for important information
 ■ do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

Inactive ingredients
 benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

Questions or comments?
 1-800-317-2165 (Mon-Fri, 9AM-5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.
 The Astepro mark is owned by Viartis. Used under license.
 © 2023 Bayer.
 Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981
 Patents: patents.livonell.bayer.com
 Product of Germany. 00000000 Bayer

INDOOR & OUTDOOR Allergy Relief

NDC 0280-0066-02

CHILDREN'S ASTEPRO ALLERGY
 Azelastine HCl 205.5 mcg per spray
 ANTIHISTAMINE NASAL SPRAY

FULL PRESCRIPTION STRENGTH

UP TO 24 HOUR RELIEF OF
 ■ Nasal congestion
 ■ Runny nose
 ■ Sneezing
 ■ Itchy nose

STERIOD FREE

120 METERED SPRAYS
 0.78 fl oz (23 mL) Alcohol-Free Fragrance-Free AGE 6+



Drug Facts
Active ingredient (in each spray)
 Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine) Purpose Antihistamine

Uses
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
 Only for use in the nose. Do not spray in eyes or mouth.
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed

When using this product
 ■ drowsiness may occur
 ■ avoid alcoholic drinks
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery
 ■ you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
 ■ nasal discomfort or sneezing may occur right after use
 ■ do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if
 ■ an allergic reaction, such as a skin rash, to this product occurs
 ■ you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ Read the User Guide for how to:
 ■ prime the bottle before first use ■ use the spray
 ■ prime bottle again if not used for 3 or more days ■ clean the spray nozzle if it gets clogged

adults and children 12 years and older	This product may be used either once or twice a day: ■ once daily: use 2 sprays in each nostril; OR ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours ■ do not use more than 4 sprays in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ do not use more than 2 sprays in each nostril in a 24 hour period
children under 6 years	do not use





This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

MARTHA K LENHART
11/20/2023 02:58:22 PM