



VARIABLE PRINT  
AREA  
Printing Process: Ink Jet  
- LOT & EXP: printed on-line  
- EXP Format: YYYY/MM

**Drug Facts (continued)**

**Other information**  
 ■ store between 20° to 25°C (68° to 77°F). Protect from freezing.  
 ■ keep this carton and the enclosed User Guide for important information.  
 ■ do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact.

**Inactive ingredients**  
 benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

**Questions or comments?**  
 1-800-317-2165 (Mon-Fri, 9AM-5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer. The Astepro mark is owned by Viartis. Used under license.  
 © 2025 Bayer.  
 Dist. by:  
 Bayer HealthCare LLC  
 Whippany, NJ 07981  
 Patents: patents.livewell.bayer.com  
 Product of Germany.  
 00000000 Bayer

**INDOOR & OUTDOOR**  
 Allergy Relief

NDC 0280-0065-10



**ASTEPRO<sup>®</sup> ALLERGY**

First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

FULL PRESCRIPTION STRENGTH

**ASTEPRO<sup>®</sup> ALLERGY**

Azelastine HCl 205.5 mcg per spray  
 ANTIHISTAMINE NASAL SPRAY

**ALLERGY + CONGESTION**


UP TO 24 HOUR RELIEF OF

- ◀ Nasal congestion
- ◀ Runny nose
- ◀ Sneezing
- ◀ Itchy nose



**22** METERED SPRAYS  
 0.14 fl oz (4 mL)

Alcohol-Free  
 Fragrance-Free



**Drug Facts**

**Active ingredient (in each spray)**  
 Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)..... Antihistamine **Purpose**

**Uses**  
 temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:  
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

**Warnings**  
 Only for use in the nose. Do not spray in eyes or mouth.  
 Do not use if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you**  
 ■ have had recent nose ulcers or nose surgery  
 ■ have had a nose injury that has not healed

**When using this product**  
 ■ drowsiness may occur  
 ■ avoid alcoholic drinks  
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness  
 ■ be careful when driving a motor vehicle or operating machinery

■ you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.  
 ■ nasal discomfort or sneezing may occur right after use  
 ■ do not share this bottle with anyone else as this may spread germs

**Stop use and ask a doctor if**  
 ■ an allergic reaction, such as a skin rash, to this product occurs  
 ■ you have severe or frequent nosebleeds

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ Read the User Guide for how to:  
 ■ prime the bottle before first use ■ use the spray  
 ■ prime bottle again if not used for 3 or more days ■ clean the spray nozzle if it gets clogged

adults and children 12 years and older This product may be used either once or twice a day:  
 ■ once daily: use 2 sprays in each nostril; OR  
 ■ twice daily: use 1 or 2 sprays in each nostril every 12 hours  
 ■ do not use more than 4 sprays in each nostril in a 24 hour period

children 6 years to 11 years ■ an adult should supervise use  
 ■ 1 spray in each nostril every 12 hours  
 ■ do not use more than 2 sprays in each nostril in a 24 hour period

children under 6 years do not use



VARIABLE PRINT  
AREA  
Printing Process: Ink Jet  
- LOT & EXP: printed on-line  
- EXP Format: YYYY/MM

**Drug Facts (continued)**

**Other information**

- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

**Inactive ingredients**

- benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

**Questions or comments?**

1-800-317-2165 (Mon-Fri, 9AM-5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer. The Astepro mark is owned by Viartis. Used under license. © 2025 Bayer. Dist. by: Bayer HealthCare LLC Whippany, NJ 07981 Patents: patents.livewell.bayer.com Product of Germany. 00000000 Bayer

**INDOOR & OUTDOOR Allergy Relief**

NDC 0280-0065-05



SAMPLE - NOT FOR SALE

**ASTEPRO ALLERGY**

Azelastine HCl 205.5 mcg per spray ANTIHISTAMINE NASAL SPRAY

**ALLERGY + CONGESTION**

UP TO 24 HOUR RELIEF OF

- ◀ Nasal congestion
- ◀ Runny nose
- ◀ Sneezing
- ◀ Itchy nose



22 METERED SPRAYS  
0.14 fl oz (4 mL)

Alcohol-Free  
Fragrance-Free

FULL PRESCRIPTION STRENGTH

STEROID FREE



ACTUAL SIZE

**Drug Facts**

**Active ingredient (in each spray)** Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)..... Antihistamine

**Purpose** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- nasal congestion
- runny nose
- sneezing
- itchy nose

**Warnings**

Only for use in the nose. Do not spray in eyes or mouth.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if

- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

Read the User Guide for how to:

- prime the bottle before first use
- prime bottle again if not used for 3 or more days
- use the spray
- clean the spray nozzle if it gets clogged

This product may be used either once or twice a day:

- adults and children 12 years and older:
  - once daily: use 2 sprays in each nostril; OR
  - twice daily: use 1 or 2 sprays in each nostril every 12 hours
  - do not use more than 4 sprays in each nostril in a 24 hour period
- children 6 years to 11 years:
  - an adult should supervise use
  - 1 spray in each nostril every 12 hours
  - do not use more than 2 sprays in each nostril in a 24 hour period
- children under 6 years: do not use





VARIABLE PRINT  
 AREA  
 Printing Process: Ink Jet  
 - LOT & EXP: printed on-line  
 - EXP Format: YYYY/MM

**Drug Facts (continued)**

**Other information**

- store between 20° to 25° C (68° to 77° F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

**Inactive ingredients**

benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

**Questions or comments?**

1-800-317-2185 (Mon-Fri, 9AM-5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer. The Astepro mark is owned by Viatris. Used under license. © 2025 Bayer.

Dist. by: Bayer HealthCare LLC Whippany, NJ 07981  
 Patents: patents.bayer.com  
 Product of Germany. Bayer 00000000

**INDOOR & OUTDOOR Allergy Relief**

NDC 0280-0065-01



**ASTEPRO ALLERGY**

First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

**ASTEPRO ALLERGY**

Azelaestine HCl 205.5 mcg per spray  
 ANTIHISTAMINE NASAL SPRAY

**ALLERGY + CONGESTION**

**UP TO 24 HOUR RELIEF OF**

- ▲ Nasal congestion
- ▲ Runny nose
- ▲ Sneezing
- ▲ Itchy nose



**60 METERED SPRAYS**  
 0.37 fl oz (11 mL)

Alcohol-Free  
 Fragrance-Free

**FULL PRESCRIPTION STRENGTH**

**STEROID FREE**



**THIS WAY TO RELIEF**

**Drug Facts**

**Active ingredient (in each spray)**

Azelaestine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)..... Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- nasal congestion
- runny nose
- sneezing
- itchy nose

**Warnings**

Only for use in the nose. Do not spray in eyes or mouth.

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

Stop use and ask a doctor if

- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

Read the User Guide for how to:

- prime the bottle before first use
- use the spray
- prime bottle again if not used for 3 or more days
- clean the spray nozzle if it gets clogged

adults and children 12 years and older This product may be used either once or twice a day:

- once daily, use 2 sprays in each nostril; OR
- twice daily, use 1 or 2 sprays in each nostril every 12 hours
- do not use more than 4 sprays in each nostril in a 24 hour period

children 6 years to 11 years ■ an adult should supervise use

- 1 spray in each nostril every 12 hours
- do not use more than 2 sprays in each nostril in a 24 hour period

children under 6 years do not use

ACTUAL SIZE

# ASTEPRO<sup>®</sup> ALLERGY

First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

### Drug Facts (continued)

**Other information**  
■ store between 20° to 25°C (68° to 77°F). Protect from freezing.  
■ keep this carton and the enclosed User Guide for important information  
■ do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

**Inactive ingredients**  
benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

**Questions or comments?**  
1-800-317-2165 (Mon-Fri, 9AM-5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer. The Astepro mark is owned by Viatrix. Used under license. © 2025 Bayer. Dist. by: Bayer HealthCare LLC Whippany, NJ 07981 Patents: patents.livestell.bayer.com Product of Germany. Bayer 00000000

**INDOOR & OUTDOOR Allergy Relief**

NDC 0280-0065-02

# ASTEPRO<sup>®</sup> ALLERGY

Azelastine HCl 205.5 mcg per spray  
ANTIHISTAMINE NASAL SPRAY

## ALLERGY + CONGESTION

UP TO 24 HOUR RELIEF OF

- ◀ Nasal congestion
- ◀ Runny nose
- ◀ Sneezing
- ◀ Itchy nose



120 METERED SPRAYS  
0.78 fl oz (23 mL)

Alcohol-Free  
Fragrance-Free

FULL PRESCRIPTION STRENGTH

STEROID FREE



<b>Drug Facts</b>	<b>Purpose</b>
<b>Active ingredient (in each spray)</b> Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)	Antihistamine
<b>Uses</b> temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose	
<b>Warnings</b> Only for use in the nose. Do not spray in eyes or mouth. <b>Do not use</b> if you have ever had an allergic reaction to this product or any of its ingredients ■ have had recent nose ulcers or nose surgery ■ have had a nose injury that has not healed	
<b>When using this product</b> ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying. ■ nasal discomfort or sneezing may occur right after use ■ do not share this bottle with anyone else as this may spread germs	
<b>Stop use and ask a doctor if</b> ■ an allergic reaction, such as a skin rash, to this product occurs ■ you have severe or frequent nosebleeds	
<b>If pregnant or breast-feeding</b> , ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> ■ Read the User Guide for how to: ■ prime the bottle before first use ■ use the spray ■ prime bottle again if not used for 3 or more days ■ clean the spray nozzle if it gets clogged	
adults and children 12 years and older	This product may be used either once or twice a day: ■ <b>once daily:</b> use 2 sprays in each nostril. <b>OR</b> ■ <b>twice daily:</b> use 1 or 2 sprays in each nostril every 12 hours ■ <b>do not use more than 4 sprays</b> in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ <b>do not use more than 2 sprays</b> in each nostril in a 24 hour period
children under 6 years	do not use



THIS WAY TO RELIEF

VARIABLE PRINT AREA  
Printing Process: Ink Jet  
- Lot & EXP: Printed On-line  
- EXP Format: YYYY/MM



**Drug Facts (continued)**

**Other information**

- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape protection\* on top and bottom carton flaps is not intact.

**Inactive ingredients**

benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

**Questions or comments?**

1-800-317-2165 (Mon-Fri, 9AM-5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer. The Astepro mark is owned by Viatis. Used under license. © 2025 Bayer. Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981. Patents: patents.livestell.bayer.com. Product of Germany. Bayer 00000000

**INDOOR & OUTDOOR Allergy Relief**

NDC 0280-0065-09

**ASTEPRO<sup>®</sup> ALLERGY**  
First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

**ASTEPRO<sup>®</sup> ALLERGY**  
Azelastine HCl 205.5 mcg per spray  
ANTIHISTAMINE NASAL SPRAY  
**ALLERGY + CONGESTION**

**FULL PRESCRIPTION STRENGTH**

**STERIOD FREE**

**UP TO 24 HOUR RELIEF OF**

- Nasal congestion
- Runny nose
- Sneezing
- Itchy nose

**2 BOTTLES x 120 SPRAYS PER BOTTLE**  
0.78 fl oz (23 mL) each  
1.56 fl oz (46 mL) total

Alcohol-Free  
Fragrance-Free



**Drug Facts**

Active ingredient (in each spray)	Purpose
Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)	Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- nasal congestion
- runny nose
- sneezing
- itchy nose

**Warnings**

Only for use in the nose. Do not spray in eyes or mouth.

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you**

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

**When using this product**

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

**Stop use and ask a doctor if**

- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

Read the User Guide for how to:

- prime the bottle before first use
- prime bottle again if not used for 3 or more days
- use the spray
- clean the spray nozzle if it gets clogged

This product may be used either once or twice a day:

- once daily: use 2 sprays in each nostril; OR
- twice daily: use 1 or 2 sprays in each nostril every 12 hours
- do not use more than 4 sprays in each nostril in a 24 hour period

children 6 years to 11 years	children under 6 years
an adult should supervise use	do not use
1 spray in each nostril every 12 hours	
do not use more than 2 sprays in each nostril in a 24 hour period	



**VARIABLE PRINT AREA**  
Printing Process: Ink Jet  
- Lot/EXP: Printed On-line  
- EXP Format: YYYY/MM

# FRONT CARD

## ASTEPRO<sup>®</sup> ALLERGY

Azelastrine HCl 205.5 mcg per spray  
ANTIHISTAMINE NASAL SPRAY

**ALLERGY +  
CONGESTION**

**STEROID  
FREE**

UP TO  
**24 HOUR  
RELIEF OF**

- ◀ Nasal congestion
- ◀ Runny nose
- ◀ Sneezing
- ◀ Itchy nose



**3** BOTTLES **x 120** SPRAYS PER BOTTLE

0.78 fl oz (23 mL) each  
2.34 fl oz (69 mL) total

Alcohol-Free  
Fragrance-Free

**FULL  
PRESCRIPTION  
STRENGTH**

00000000

# ASTEPRO<sup>®</sup> ALLERGY



**First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use**

<b>Drug Facts</b>		<b>Drug Facts (continued)</b>	
<b>Active ingredient (in each spray)</b> Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine).....	<b>Purpose</b> Antihistamine	<b>Directions</b> Read the User Guide for how to: <ul style="list-style-type: none"> <li>■ prime the bottle before first use</li> <li>■ use the spray</li> <li>■ prime bottle again if not used for 3 or more days</li> <li>■ clean the spray nozzle if it gets clogged</li> </ul>	
<b>Uses</b> temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: <ul style="list-style-type: none"> <li>■ nasal congestion</li> <li>■ runny nose</li> <li>■ sneezing</li> <li>■ itchy nose</li> </ul>		adults and children 12 years and older	This product may be used either once or twice a day: <ul style="list-style-type: none"> <li>■ <b>once daily:</b> use 2 sprays in each nostril; <b>OR</b></li> <li>■ <b>twice daily:</b> use 1 or 2 sprays in each nostril every 12 hours</li> <li>■ <b>do not use more than 4 sprays</b> in each nostril in a 24 hour period</li> </ul>
<b>Warnings</b> Only for use in the nose. Do not spray in eyes or mouth. <b>Do not use</b> if you have ever had an allergic reaction to this product or any of its ingredients <b>Ask a doctor before use if you</b> <ul style="list-style-type: none"> <li>■ have had recent nose ulcers or nose surgery</li> <li>■ have had a nose injury that has not healed</li> </ul> <b>When using this product</b> <ul style="list-style-type: none"> <li>■ drowsiness may occur</li> <li>■ avoid alcoholic drinks</li> <li>■ alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>■ be careful when driving a motor vehicle or operating machinery</li> <li>■ you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.</li> <li>■ nasal discomfort or sneezing may occur right after use</li> <li>■ do not share this bottle with anyone else as this may spread germs</li> </ul> <b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>■ an allergic reaction, such as a skin rash, to this product occurs</li> <li>■ you have severe or frequent nosebleeds</li> </ul> <b>If pregnant or breast-feeding,</b> ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.		children 6 years to 11 years	<ul style="list-style-type: none"> <li>■ an adult should supervise use</li> <li>■ 1 spray in each nostril every 12 hours</li> <li>■ <b>do not use more than 2 sprays</b> in each nostril in a 24 hour period</li> </ul>
		children under 6 years	do not use
		<b>Other information</b> <ul style="list-style-type: none"> <li>■ store between 20° to 25°C (68° to 77°F). Protect from freezing.</li> <li>■ keep this label and the enclosed User Guide for important information</li> <li>■ do not use if sealed package is torn or opened</li> </ul>	
		<b>Inactive ingredients</b> benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose	
		<b>Questions or comments?</b> 1-800-317-2165 (Mon-Fri, 9AM-5PM EST)	

Bayer and the Bayer Cross are registered trademarks of Bayer.  
 The Astepro mark is owned by Viartis. Used under license.  
 © 2021 Bayer.  
 Dist. by:  
 Bayer HealthCare LLC  
 Whippany, NJ 07981  
 Patents: patents.livewell.bayer.com  
 Product of Germany.  
 000000

NDC 0280-0065-03  
 LOT:  
 EXP: YYYY-MM  
 Bayer



# ASTEPRO<sup>®</sup> ALLERGY

First Antihistamine Nasal Spray FDA-Approved for Over-the-Counter Use

### Drug Facts (continued)

**Other information**  
■ store between 20° to 25°C (68° to 77°F). Protect from freezing.  
■ keep this carton and the enclosed User Guide for important information  
■ do not use if tape printed with "sealed for your protection" on top and bottom carton flaps is not intact

**Inactive ingredients**  
benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

**Questions or comments?**  
1-800-317-2165 (Mon-Fri, 9AM-5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.  
The Astepro mark is owned by Viatrix.  
Used under license.  
© 2025 Bayer.  
Dist. by:  
Bayer HealthCare LLC  
Whippany, NJ 07981  
Patents: patents.livewell.bayer.com  
Product of Germany. Bayer  
00000000

**INDOOR & OUTDOOR**  
Allergy Relief

NDC 0280-0065-04

# ASTEPRO<sup>®</sup> ALLERGY

Azelastine HCl 205.5 mcg per spray  
ANTIHISTAMINE NASAL SPRAY

## ALLERGY + CONGESTION

UP TO  
**24 HOUR RELIEF OF**

- ◀ Nasal congestion
- ◀ Runny nose
- ◀ Sneezing
- ◀ Itchy nose



**200 METERED SPRAYS**  
1.01 fl oz (30 mL)

Alcohol-Free  
Fragrance-Free

FULL  
PRESCRIPTION  
STRENGTH

STEROID  
FREE



Drug Facts	
<b>Active ingredient (in each spray)</b> Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)	<b>Purpose</b> Antihistamine
<b>Uses</b> temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose	
<b>Warnings</b> Only for use in the nose. Do not spray in eyes or mouth. <b>Do not use</b> if you have ever had an allergic reaction to this product or any of its ingredients	
<b>Ask a doctor before use if you</b> ■ have had recent nose ulcers or nose surgery ■ have had a nose injury that has not healed	
<b>When using this product</b> ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying. ■ nasal discomfort or sneezing may occur right after use ■ do not share this bottle with anyone else as this may spread germs	
<b>Stop use and ask a doctor if</b> ■ an allergic reaction, such as a skin rash, to this product occurs ■ you have severe or frequent nosebleeds	
<b>If pregnant or breast-feeding</b> , ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> ■ Read the User Guide for how to: ■ prime the bottle before first use ■ use the spray ■ clean the spray nozzle if it gets clogged	
adults and children 12 years and older	This product may be used either once or twice a day: ■ <b>once daily:</b> use 2 sprays in each nostril. <b>OR</b> ■ <b>twice daily:</b> use 1 or 2 sprays in each nostril every 12 hours ■ <b>do not use more than 4 sprays</b> in each nostril in a 24 hour period
children 6 years to 11 years	■ an adult should supervise use ■ 1 spray in each nostril every 12 hours ■ <b>do not use more than 2 sprays</b> in each nostril in a 24 hour period
children under 6 years	do not use



000000



0 000000 000000 0



000000

VARIABLE PRINT AREA  
Printing Process: Ink Jet  
- Lot & EXP: Printed On-Line  
- EXP Format: YYYY/MM

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

NUSHIN F TODD  
10/27/2025 03:43:30 PM