

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use KYZATREX® safely and effectively. See full prescribing information for KYZATREX.

**KYZATREX (testosterone undecanoate) capsules, for oral use, CIII, Initial U.S. approval: 1953**

### RECENT MAJOR CHANGES

Boxed Warnings, Blood Pressure Increases	Removed 07/2025
Warnings and Precautions, Venous Thromboembolism (5.2)	07/2025
Warnings and Precautions, Blood Pressure Increases (5.4)	07/2025
Warnings and Precautions, Cardiovascular Risk (5.4)	Removed 07/2025

### INDICATIONS AND USAGE

KYZATREX is an androgen indicated for testosterone replacement therapy in adult males for conditions associated with a deficiency or absence of endogenous testosterone (1).

#### Limitations of Use:

- Safety and efficacy of KYZATREX in males less than 18 years old have not been established (1, 8.4).
- Safety and efficacy of KYZATREX in men with “age-related hypogonadism” have not been established (1).

### DOSAGE AND ADMINISTRATION

- KYZATREX is not substitutable with other oral testosterone undecanoate products (2.1).
- Prior to initiating KYZATREX, confirm the diagnosis of hypogonadism by ensuring that serum testosterone concentrations have been measured in the morning on at least two separate days and that these concentrations are below the normal range (2.2).
- Take KYZATREX with food (2.3).
- Starting dosage: 200 mg orally once in the morning and once in the evening (2.3).
- Adjust the dosage to a minimum of 100 mg once in the morning and a maximum of 400 mg twice daily based on serum testosterone drawn 3 to 5 hours after the morning dose at least 7 days after starting treatment or following dose adjustment and periodically thereafter (2.3).

### DOSAGE FORMS AND STRENGTHS

Capsules: 100 mg, 150 mg, 200 mg (3).

### CONTRAINDICATIONS

- Carcinoma of the breast or known or suspected carcinoma of the prostate (5.4)
- Women who are pregnant. Testosterone may cause fetal harm (4, 5.7, 8.1)
- Hypersensitivity to KYZATREX or any of its ingredients (4)
- Hypogonadal conditions not associated with structural or genetic etiologies (4)

### WARNINGS AND PRECAUTIONS

- **Polycythemia:** Monitor hemoglobin or hematocrit approximately every 3 months to detect increased red blood cell mass and polycythemia. Discontinue KYZATREX if necessary (5.1).
- **Venous thromboembolism (VTE):** VTE, including deep vein thrombosis (DVT) and pulmonary embolism (PE) have been reported in patients using testosterone. Discontinue KYZATREX if VTE is suspected and initiate appropriate workup and management (5.2).
- **Worsening of Benign Prostatic Hyperplasia (BPH) and Potential Risk of Prostate Cancer:** Monitor patients for worsening of signs and symptoms of BPH. Evaluate patients for prostate cancer, including monitoring prostate specific antigen (PSA) prior to initiating and during treatment with androgens (5.3).
- **Blood Pressure Increases:** KYZATREX can increase blood pressure, which can increase cardiovascular risk over time. Measure blood pressure periodically. Not recommended for use in men with uncontrolled hypertension (5.4)
- **Abuse of Testosterone and Monitoring of Testosterone Concentrations:** If testosterone use at doses higher than recommended for the approved indication and in combination with other anabolic androgenic steroids is suspected, check serum testosterone concentration (5.5).
- **Potential for Adverse Effects on Spermatogenesis:** KYZATREX may cause azoospermia (5.7).
- **Edema:** Edema, with or without congestive heart failure (CHF), may occur in patients with pre-existing cardiac, renal, or hepatic disease. Discontinue KYZATREX and initiate appropriate workup (5.9).
- **Sleep Apnea:** KYZATREX may potentiate sleep apnea in those with risk factors (5.10)
- **Lipid Changes:** KYZATREX may affect serum lipid profile. Monitor patient lipid concentrations periodically; if necessary, adjust dosage of lipid lowering drug(s) or discontinue KYZATREX (5.12).

### ADVERSE REACTIONS

Most common adverse reactions (incidence ≥ 2%): hypertension (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Marius Pharmaceuticals at 1-833-949-5040 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### DRUG INTERACTIONS

- **Insulin:** In patients with diabetes, concomitant use with KYZATREX may decrease blood glucose and insulin requirements (7.1).
- **Oral Anticoagulants:** Concomitant use with KYZATREX may cause changes in anticoagulant activity. Monitor International Normalized Ratio (INR) and prothrombin time (PT) frequently (7.2).
- **Corticosteroids:** Concomitant use with KYZATREX may result in increased fluid retention. Use with caution, particularly in patients with cardiac, renal, or hepatic disease (7.3).
- **Drugs that May Also Increase Blood Pressure:** Concomitant use with KYZATREX may lead to additional increases in blood pressure (7.4).

### USE IN SPECIFIC POPULATIONS

**Geriatric Patients:** Geriatric patients treated with androgens may also be at risk for worsening of signs and symptoms of BPH and hypertension (8.5).

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling and Medication Guide.

Revised: 07/2025

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**WARNING: BLOOD PRESSURE INCREASES**

- 1. INDICATIONS AND USAGE**
- 2. DOSAGE AND ADMINISTRATION**
  - 2.1 Important Dosage Information
  - 2.2 Confirmation of Hypogonadism Before Initiation of KYZATREX
  - 2.3 Recommended Dosage and Administration
- 3. DOSAGE FORMS AND STRENGTHS**
- 4. CONTRAINDICATIONS**
- 5. WARNINGS AND PRECAUTIONS**
  - 5.1 Polycythemia
  - 5.2 Venous Thromboembolism
  - 5.3 Worsening of Benign Prostatic Hyperplasia (BPH) and Potential Risk of Prostate Cancer
  - 5.4 Blood Pressure Increases
  - 5.5 Abuse of Testosterone and Monitoring of Testosterone Concentrations
  - 5.6 Not for Use in Women
  - 5.7 Potential for Adverse Effects on Spermatogenesis
  - 5.8 Hepatic Adverse Events
  - 5.9 Edema
  - 5.10 Sleep Apnea
  - 5.11 Gynecomastia
  - 5.12 Lipid Changes
  - 5.13 Hypercalcemia
  - 5.14 Decreased Thyroxine-binding Globulin
- 6. ADVERSE REACTIONS**
  - 6.1 Clinical Trial Experience
  - 6.2 Postmarketing Experience

- 7. DRUG INTERACTIONS**
  - 7.1 Insulin
  - 7.2 Oral Vitamin K Antagonist Anticoagulants
  - 7.3 Corticosteroids
  - 7.4 Medications that May Also Increase Blood Pressure

- 8. USE IN SPECIFIC POPULATIONS**
  - 8.1 Pregnancy
  - 8.2 Lactation
  - 8.3 Females and Males of Reproductive Potential
  - 8.4 Pediatric Use
  - 8.5 Geriatric Use

- 9. DRUG ABUSE AND DEPENDENCE**
  - 9.1 Controlled Substance
  - 9.2 Abuse
  - 9.3 Dependence

**10. OVERDOSAGE**

**11. DESCRIPTION**

**12. CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

**13. NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

**14. CLINICAL STUDIES**

**16. HOW SUPPLIED/STORAGE AND HANDLING**

**17. PATIENT COUNSELING INFORMATION**

\*Sections or subsections omitted from the full prescribing information are not listed.

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

KYZATREX is indicated for testosterone replacement therapy in adult males for conditions associated with a deficiency or absence of endogenous testosterone:

- Primary hypogonadism (congenital or acquired): testicular failure due to conditions such as cryptorchidism, bilateral torsion, orchitis, vanishing testis syndrome, orchiectomy, Klinefelter syndrome, chemotherapy, or toxic damage from alcohol or heavy metals. These men usually have low serum testosterone concentrations and gonadotropins (follicle-stimulating hormone (FSH), luteinizing hormone (LH)) above the normal range.
- Hypogonadotropic hypogonadism (congenital or acquired): gonadotropin or luteinizing hormone-releasing hormone (LHRH) deficiency, pituitary-hypothalamic injury from tumors, trauma, or radiation. These men have low serum testosterone concentrations but have gonadotropins in the normal or low range.

#### **Limitations of Use**

- Safety and efficacy of KYZATREX in males less than 18 years old have not been established [*see Use in Specific Populations (8.4)*].
- Safety and efficacy of KYZATREX in men with “age-related hypogonadism” (also referred to as “late-onset hypogonadism”) have not been established.

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Important Dosage Information

KYZATREX is not substitutable with other oral testosterone undecanoate products.

#### 2.2 Confirmation of Hypogonadism Before Initiation of KYZATREX

Prior to initiating KYZATREX, confirm the diagnosis of hypogonadism by ensuring that serum testosterone concentrations have been measured in the morning on at least two separate days and that these testosterone concentrations are below the normal range.

#### 2.3 Recommended Dosage and Administration

Individualize the dosage of KYZATREX based on the patient’s serum testosterone concentration response to the drug.

The recommended starting dose is 200 mg orally twice daily, once in the morning and once in the evening. Take KYZATREX with food.

#### **Dosage Adjustment**

Check serum testosterone concentrations 7 days after starting treatment or after dosage adjustment, 3 to 5 hours after the morning dose. Adjust the KYZATREX dose as necessary as shown in [Table 1](#). Thereafter, periodically monitor serum testosterone concentrations.

The minimum recommended dose is 100 mg once daily in the morning. The maximum recommended dose is 400 mg twice daily. For total daily doses greater than 100 mg, administer the same dose in the morning and evening.

**Table 1: KYZATREX Dosage Adjustment Scheme**

Serum Testosterone Concentration	Current KYZATREX Dosage	New KYZATREX Dosage
Less than 460 ng/dL	100 mg with breakfast only	100 mg twice daily with meals
	100 mg twice daily with meals	200 mg twice daily with meals
	200 mg twice daily with meals	300 mg twice daily with meals
	300 mg twice daily with meals	400 mg twice daily with meals
460 to 971 ng/dL	No Dosage Change	
More than 971 ng/dL	400 mg twice daily with meals	300 mg twice daily with meals
	300 mg twice daily with meals	200 mg twice daily with meals
	200 mg twice daily with meals	100 mg twice daily with meals
	100 mg twice daily with meals	100 mg with breakfast only
	100 mg with breakfast only	Discontinue treatment

### 3 DOSAGE FORMS AND STRENGTHS

Capsules:

- 100 mg, oval, opaque, white, imprinted with “MP100” in red ink
- 150 mg, oblong, opaque, white, imprinted with “MP150” in red ink
- 200 mg, oblong, opaque, white, imprinted with “MP200” in red ink

### 4 CONTRAINDICATIONS

KYZATREX is contraindicated in:

- Patients with carcinoma of the breast or known or suspected carcinoma of the prostate [*see Warnings and Precautions (5.3)*].
- Women who are pregnant. Testosterone can cause virilization of the female fetus when administered to a pregnant woman [*see Use in Specific Populations (8.1)*].
- Patients with known hypersensitivity to KYZATREX or any of its ingredients [*see Description (11)*].

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Polycythemia

Androgens, including KYZATREX, can cause increase in hemoglobin or hematocrit, reflective of increase in red blood cell mass. Check hematocrit prior to initiating KYZATREX. An

increase in red blood cell mass may increase the risk of thromboembolic events [see *Warnings and Precautions* (5.2)]. Evaluate hematocrit approximately every 3 months while the patient is on KYZATREX. If hematocrit becomes elevated, stop KYZATREX until the hematocrit decreases to an acceptable concentration. If KYZATREX is restarted and again causes hematocrit to become elevated, permanently discontinue KYZATREX.

## 5.2 Venous Thromboembolism

There have been post-marketing reports of venous thromboembolic events, including deep vein thrombosis (DVT) and pulmonary embolism (PE), in patients using testosterone replacement products such as KYZATREX.

In the Testosterone Replacement therapy for Assessment of long-term Vascular Events and efficacy Response in hypogonadal men (TRAVERSE) Study, a randomized, double-blind, placebo-controlled, cardiovascular (CV) outcomes study, compared to placebo, topical testosterone gel was associated with a numerically higher incidence of VTE (1.7% vs 1.2%) which included DVT (0.6% vs 0.5%) and PE events (0.9% vs 0.5%) [see *Adverse Reactions* (6.1)].

Evaluate patients who report symptoms of pain, edema, warmth, and erythema in the lower extremity for DVT and those who present with acute shortness of breath for PE. If a venous thromboembolic event is suspected, discontinue KYZATREX and initiate appropriate workup and management [see *Adverse Reactions* (6.2)].

## 5.3 Worsening of Benign Prostatic Hyperplasia (BPH) and Potential Risk of Prostate Cancer

- Patients with BPH who are treated with androgens are at an increased risk for worsening of signs and symptoms of BPH. Monitor patients with BPH for worsening signs and symptoms.
- Patients treated with androgens may be at increased risk for prostate cancer. Evaluate patients for prostate cancer prior to initiating and during treatment with androgens [see *Contraindications* (4)].

## 5.4 Blood Pressure Increases

KYZATREX can increase blood pressure. Based on ambulatory blood monitoring in Study MRS-TU-2019EXT, KYZATREX increased mean systolic/diastolic blood pressure by 1.7/0.6 mm Hg from baseline after 4 months of treatment and 1.8/0.6 mm Hg from baseline after 6 months of treatment [see *Adverse Reactions* (6.1)]. In patients with hypertension on antihypertensive therapy, KYZATREX increased the mean systolic/diastolic BP by 3.4/0.7 mm Hg from baseline after 4 months of treatment and 3.1/1.0 mm Hg from baseline after 6 months of treatment [see *Adverse Reactions* (6.1)]. Blood pressure increases can increase cardiovascular (CV) risk over time.

The CV risk associated with topical testosterone gel was evaluated in TRAVERSE, a randomized, double-blind, placebo controlled, CV outcomes study in men with a history of CV disease or multiple CV risk factors. In TRAVERSE, topical testosterone gel increased mean systolic blood pressure by 1.0 mm Hg from baseline to 36 months, whereas a mean decrease from baseline of 0.5 mm Hg was observed in the placebo group at this timepoint, for a mean between-group difference of 1.5 mm Hg. However, the incidences of major adverse cardiovascular events (MACE), including cardiovascular death, non-fatal myocardial infarction [MI] and non-fatal stroke, were similar between treatment groups (7% for topical testosterone gel vs 7.3% for placebo) [See *Adverse Reactions* (6.1)].

Monitor BP periodically in men using KYZATREX, especially men with hypertension.

KYZATREX is not recommended for use in patients with uncontrolled hypertension.

### **5.5 Abuse of Testosterone and Monitoring of Testosterone Concentrations**

Testosterone has been subject to abuse, typically at doses higher than recommended for the approved indication and in combination with other anabolic androgenic steroids. Anabolic androgenic steroid abuse can lead to serious cardiovascular and psychiatric adverse reactions [see *Drug Abuse and Dependence (9)*].

If testosterone abuse is suspected, check testosterone concentrations to ensure they are within therapeutic range [see *Dosage and Administration (2.2)*]. Testosterone levels may remain in the normal or subnormal range in men abusing synthetic testosterone derivatives. Counsel patients concerning the serious adverse reactions associated with abuse of testosterone and anabolic androgenic steroids. Also consider the possibility of testosterone and anabolic androgenic steroid abuse in suspected patients who present with serious cardiovascular or psychiatric adverse events.

### **5.6 Not for Use in Women**

Due to lack of controlled studies in women and potential virilizing effects, KYZATREX is not indicated for use in women [see *Contraindications (4)* and *Use in Specific Populations (8.1, 8.2)*].

### **5.7 Potential for Adverse Effects on Spermatogenesis**

With large doses of exogenous androgens, including KYZATREX, spermatogenesis may be suppressed through feedback inhibition of pituitary FSH, possibly leading to adverse effects on semen parameters including sperm count [see *Use in Specific Populations (8.3)*]. Inform patients of this possible risk when deciding whether to use or to continue to use KYZATREX.

### **5.8 Hepatic Adverse Effects**

KYZATREX is not a 17-alpha-alkyl androgen and is not known to cause hepatic adverse effects. However, prolonged use of high doses of orally active 17-alpha-alkyl androgens (e.g., methyltestosterone) has been associated with serious hepatic adverse effects (peliosis hepatis, hepatic neoplasms, cholestatic hepatitis, and jaundice). Peliosis hepatis can be a life-threatening or fatal complication. Long-term therapy with intramuscular testosterone enanthate has produced multiple hepatic adenomas. Patients should be instructed to report any signs or symptoms of hepatic dysfunction (e.g., jaundice). If these occur, promptly discontinue KYZATREX while the cause is evaluated.

### **5.9 Edema**

Androgens, including KYZATREX, may promote retention of sodium and water. Edema, with or without congestive heart failure, may be a serious complication in patients with pre-existing cardiac, renal, or hepatic disease [see *Adverse Reactions (6.1)*]. In addition to discontinuation of the drug, diuretic therapy may be required.

### **5.10 Sleep Apnea**

The treatment of hypogonadal men with testosterone may potentiate sleep apnea in some patients, especially those with risk factors such as obesity or chronic lung disease.

### **5.11 Gynecomastia**

Gynecomastia may develop and persist in patients being treated for hypogonadism.

### 5.12 Lipid Changes

In clinical trials, patients receiving KYZATREX experienced reductions in lipid parameters, including total cholesterol, LDL cholesterol, HDL cholesterol, and triglycerides [see *Adverse Reactions (6.1)*]. Changes in the serum lipid profile may require dose adjustment of lipid lowering drugs or discontinuation of testosterone therapy. Monitor the lipid profile periodically, particularly after starting testosterone therapy.

### 5.13 Hypercalcemia

Androgens, including KYZATREX, should be used with caution in cancer patients at risk of hypercalcemia (and associated hypercalciuria). Monitor serum calcium concentrations periodically during treatment with KYZATREX in these patients.

### 5.14 Decreased Thyroxine-binding Globulin

Androgens, including KYZATREX, may decrease concentrations of thyroxin-binding globulin, resulting in decreased total T4 serum concentrations and increased resin uptake of T3 and T4. Free thyroid hormone concentrations remain unchanged, however, and there is no clinical evidence of thyroid dysfunction.

## 6 ADVERSE REACTIONS

The following clinically significant adverse reactions are discussed elsewhere in the labeling:

- Polycythemia [see *Warnings and Precautions (5.1)*]
- Venous Thromboembolism [see *Warnings and Precautions (5.2)*]
- Worsening of Benign Prostatic Hyperplasia (BPH) and Potential Risk of Prostate Cancer [see *Warnings and Precautions (5.3)*]
- Blood Pressure Increases [see *Warnings and Precautions (5.4)*]
- Hepatic Adverse Effects [see *Warnings and Precautions (5.8)*]
- Edema [see *Warnings and Precautions (5.9)*]
- Sleep Apnea [see *Warnings and Precautions (5.10)*]
- Gynecomastia [see *Warnings and Precautions (5.11)*]
- Lipid Changes [see *Warnings and Precautions (5.12)*]
- Hypercalcemia [see *Warnings and Precautions (5.13)*]
- Decreased Thyroxine-binding Globulin [see *Warnings and Precautions (5.14)*]

### 6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of KYZATREX was evaluated in Study MRS-TU-2019EXT in 155 hypogonadal males [see *Clinical Studies (14)*]. All patients initially received KYZATREX 200 mg orally twice daily. If needed, the dosage was titrated to 100 mg once daily in the morning or 100 mg, 300 mg, or 400 mg twice daily to achieve testosterone concentrations in the normal range. After the dosage titration period, patients continued their optimized dose for the remainder of the duration of the 6-month study. The mean duration of exposure was 168 days (range: 1 to 180 days). The median age was 52 years (range: 22 to 66 years); 77% were White, 19% were Black, 3% were Asian, and 2% were American Indian, Alaskan Native or Other.

**Table 2** summarizes adverse reactions reported in  $\geq 2\%$  of patients in this 6-month study.

**Table 2: Adverse Reactions in  $\geq 2\%$  of Patients Receiving KYZATREX in STUDY MRS-TU-2019EXT**

Adverse Reaction	N = 155 n (%)
Hypertension <sup>a</sup>	4 (2.6)

<sup>a</sup> Based upon blood pressure cuff measurements

One (0.8%) patient who received KYZATREX experienced an adverse reaction (acne) that led to premature discontinuation from the study.

In a 12-month, open-label study in hypogonadal adult males (N=212) who received KYZATREX 200 mg once daily to 400 mg twice daily (n=202) the following additional adverse reactions were reported: headache, arthralgia, diarrhea, hemoglobin increased, anxiety, constipation, peripheral edema, and PSA increased.

### Blood Pressure Increases

In Study MRS-TU-2019EXT, 24-hour ambulatory blood pressure monitoring (ABPM) was conducted in 155 male patients, 135 of whom completed the study. ABPM was conducted at 3 distinct 24-hour time periods: at baseline and following approximately 4 months and 6 months weeks of treatment with KYZATREX. A total of 151 patients had acceptable 24-hour ABPM recordings at both time periods. In that group, the mean change in systolic BP from Baseline to 4 months and 6 months was + 1.7 mm Hg (95% CI 0.3, 3.1) and 1.8 mm Hg (95% CI 0.3, 3.2), respectively. In that group, the mean change in diastolic BP from Baseline to 4 months and 6 months was 0.6 mm Hg (95% CI -0.3, 1.6) and 0.6 mm Hg (95% CI -0.4, 1.6), respectively. In patients with a history of hypertension on antihypertensive therapy at baseline, the mean ABPM systolic blood pressure increased from Baseline to 4 months and 6 months by 3.4 mm Hg (95% CI 1.0, 5.9) and 3.1 mm Hg (95% CI 0.6, 5.6), respectively (n=49).

In patients with no history of hypertension at baseline, the mean systolic blood pressure from Baseline increased by 0.7 mm Hg (95% CI -1.0, 2.4) at 4 months and 1.0 mm Hg (95% CI -0.7, 2.8), at six months respectively (n =90). Ambulatory (24-hour) and in-clinic (cuff) blood pressure. Changes from Baseline for study MRS- TU-2019EXT are presented in **Table 3** with 95% confidence intervals. No significant difference was observed between the 4-month and 6-month Changes from Baseline.

**Table 3: Blood Pressure Increases**

Blood Pressure	Change from Baseline (95% CI), mm Hg	
	Systolic	Diastolic
24-Hour Ambulatory		
4 Month	1.7 (0.3 to 3.1)	0.6 (-0.3 to 1.6)
6 Month	1.8 (0.3 to 3.2)	0.6 (-0.4 to 1.6)

A history of antihypertensive treatment and diabetes mellitus at baseline were significant factors related to ambulatory SBP increases.

Table 4 presents the Least Squares Mean estimates of Change from Baseline, with 95% CI's, for sub-populations of subjects at study start either with or without hypertensive treatment or with or without diabetes mellitus.

**Table 4: Ambulatory Blood Pressure Increases by Subject Sub-group**

Subgroups	Ambulatory Systolic Blood Pressure Change from Baseline (95% CI), mm Hg
With hypertensive treatment at baseline (n=49)	
4 Month	3.4 (1.0 to 5.9)
6 Month	3.1 (0.6 to 5.6)
Without hypertensive treatment at baseline (n=90)	
4 Month	0.7 (-1.0 to 2.4)
6 Month	1.0 (-0.7 to 2.8)
With diabetes at baseline (n=29)	
4 Month	3.0 (-0.2 to 6.2)
6 Month	3.4 (0.2 to 6.7)
Without diabetes at baseline (n=110)	
4 Month	1.3 (-0.2 to 2.9)
6 Month	1.3 (-0.3 to 2.9)

A total of 5 of 155 patients (3.2%) on KYZATREX in Study MRS-TU-2019EXT began taking new antihypertensive medications after study start. No patient had a dose increase in their antihypertensive medication by the end of treatment.

Of the 155 patients in Study MRS-TU-2019EXT who used KYZATREX, 4 patients (2.6%) were reported to have an adverse reaction of hypertension.

### **Cardiovascular Outcomes**

TRAVERSE was a randomized, double-blind, cardiovascular outcomes study to assess the cardiovascular (CV) safety of topical testosterone gel compared to placebo in 5198 hypogonadal men aged 45 to 80 years with a history of CV disease or with multiple CV risk factors. The primary outcome was the incidence of the composite endpoint of major adverse cardiovascular events (MACE), consisting of CV death, non-fatal myocardial infarction (MI), and non-fatal stroke.

The mean duration of therapy was approximately 22 months. The mean duration of follow-up was 33 months. Approximately 61% of all patients discontinued topical testosterone gel or placebo therapy.

The mean patient age ( $\pm$ SD) was 63.3 (7.9) years, with 2452 patients aged 65 years or more (47%); 2847 (about 55%) patients had pre-existing cardiovascular disease, whereas 2357 patients (about 45%) had an elevated cardiovascular risk at baseline, and mean BMI was 35kg/m<sup>2</sup>. Approximately 80% of patients were White, 17% were Black, and 3% were of other races or ethnic groups. Approximately 69%, 84%, and 93% had diabetes mellitus, hyperlipidemia, and hypertension, respectively.

The mean serum testosterone concentration at baseline in patients receiving topical testosterone gel was 220.4 ng/dL (n=2596). The mean serum testosterone concentrations at 12 months, 24 months, 36 months, and 48 months in patients receiving topical testosterone gel were 440.5 ng/dL (n=1683), 420.9 ng/dl (n=1125), 428.7 ng/dL (n=731), and 365.2 ng/dL (n=220), respectively.

For patients treated with topical testosterone gel, the incidence of MACE was 7.0% (n=182 events) and for those receiving placebo, the incidence of MACE was 7.3% (n=190 events). The study demonstrated non-inferiority of topical testosterone gel versus placebo because the upper bound of 95% CI was less than the pre-specified risk margin, of 1.5 for MACE (Hazard Ratio 0.96 [95% CI: 0.78, 1.17]).

### **Additional Adverse Reactions Reported in TRAVERSE**

Additional adverse reactions reported in TRAVERSE at an incidence rate >2% in either treatment group and greater in topical testosterone gel versus placebo included: nonfatal arrhythmias warranting intervention (5.2% vs 3.3%), atrial fibrillation (3.5% vs 2.4%), acute kidney injury (2.3% vs 1.5%) and bone fracture (3.5% vs 2.5%). For the adverse reaction of bone fracture, each event was adjudicated by clinical review.

#### **Heart Rate Increases**

KYZATREX increased mean (95%CI) 24-hour ambulatory heart rate by an average of 0.7 (-0.5 to 1.9) beats per minute (bpm) at 4 months and 1.9 (0.6 to 3.1) bpm at 6 months in Study MRS-TU-2019EXT. Changes in heart rate were similar between patients with or without hypertension or diabetes. Changes in heart rate with treatment were most prominent in the evening, 12 to 17 hours after the morning dose.

#### **Increases in Hemoglobin**

Increases in hemoglobin were reported in 7 out of 155 patients (4.5%) in Study MRS-TU-2019EXT. None of these increases led to premature discontinuation of KYZATREX.

Hematocrit was not assessed in this study.

#### **Headaches**

Headaches were reported in 3 of 155 patients (1.9%) receiving KYZATREX in Study MRS-TU-2019EXT.

#### **Increases in Serum PSA**

Four out of 155 patients (2.6%) receiving KYZATREX in Study MRS-TU-2019EXT had an increase in PSA from baseline greater than 1.4 ng/mL and two out of 155 patients (1.3%) had a PSA of at least 4.0 ng/mL during Study MRS-TU-2019EXT. The mean (SE) increase in PSA from baseline was 0.15 ( $\pm$ 0.04) ng/mL at 6 months (n=135).

## 6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of testosterone. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Cardiovascular Disorders: myocardial infarction, stroke*

*Vascular Disorders: Venous thromboembolism*

## 7 DRUG INTERACTIONS

### 7.1 Insulin

Changes in insulin sensitivity or glycemic control may occur in patients treated with androgens. In diabetic patients, the metabolic effects of androgens may decrease blood glucose and therefore necessitate a decrease in the dose of anti-diabetic medication.

### 7.2 Oral Vitamin K Antagonist Anticoagulants

Changes in anticoagulant activity may be seen with androgens; therefore, more frequent monitoring of international normalized ratio (INR) and prothrombin time are recommended in patients taking warfarin, especially at the initiation and termination of androgen therapy.

### 7.3 Corticosteroids

The concurrent use of testosterone with corticosteroids may result in increased fluid retention and requires careful monitoring particularly in patients with cardiac, renal, or hepatic disease.

### 7.4 Medications that May Also Increase Blood Pressure

Some prescription medications and nonprescription analgesic and cold medications contain drugs known to increase blood pressure. Concomitant administration of these medications with KYZATREX may lead to additional increases in blood pressure [see *Warnings and Precautions (5.4)*].

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

KYZATREX is contraindicated in pregnant women and not indicated for use in females [see *Contraindications (4)*]. Testosterone is teratogenic and may cause fetal harm when administered to a pregnant woman based on data from animal studies (see [Data](#)) and its mechanism of action [see *Clinical Pharmacology (12.1)*]. Exposure of a female fetus to androgens may result in varying degrees of virilization. In animal developmental studies, exposure to testosterone in utero resulted in hormonal and behavioral changes in offspring and structural impairments of reproductive tissues in female and male offspring. These studies do not meet current standards for nonclinical development toxicity studies.

#### Data

##### *Animal Data*

In developmental studies conducted in rats, rabbits, pigs, sheep, and rhesus monkeys, pregnant animals received intramuscular injections of testosterone during the period of organogenesis.

Testosterone treatment at doses that were comparable to those used for testosterone replacement therapy resulted in structural impairments in both female and male offspring. Structural impairments observed in females included increased anogenital distance, phallus development, empty scrotum, no external vagina, intrauterine growth retardation, reduced ovarian reserve, and increased ovarian follicular recruitment. Structural impairments seen in male offspring included increased testicular weight, larger seminal tubular lumen diameter, and higher frequency of occluded tubule lumen. Increased pituitary weight was seen in both sexes.

Testosterone exposure in utero also resulted in hormonal and behavioral changes in offspring. Hypertension was observed in pregnant female rats and their offspring exposed to doses approximately twice those used for testosterone replacement therapy.

## **8.2 Lactation**

### Risk Summary

KYZATREX is not indicated for use in females.

## **8.3 Females and Males of Reproductive Potential**

### Infertility

#### *Males*

During treatment with large doses of exogenous androgens, including KYZATREX, spermatogenesis may be suppressed through feedback inhibition of the hypothalamic-pituitary-testicular axis [see *Warnings and Precautions (5.7) and Impairment of Fertility (13.1)*], possibly leading to adverse effects on semen parameters including sperm count. Reduced fertility has been observed in some men taking testosterone replacement therapy. Testicular atrophy, subfertility, and infertility have also been reported in men who abuse anabolic androgenic steroids [see *Drug Abuse and Dependence (9.2)*]. With either type of use, the impact on fertility may be irreversible.

## **8.4 Pediatric Use**

The safety and efficacy of KYZATREX in pediatric patients less than 18 years old have not been established. KYZATREX is not recommended for use in patients less than 18 years of age because of the potential for acceleration of bone age and premature closure of epiphyses.

## **8.5 Geriatric Use**

Clinical studies of KYZATREX did not include any patients 65 years of age and older. Therefore, it cannot be determined whether these patients respond differently from younger adult patients. Additionally, there are insufficient long-term safety data in geriatric patients to assess the potentially increased risk of cardiovascular disease and prostate cancer.

Geriatric patients treated with androgens including KYZATREX may be at risk for worsening of signs and symptoms of BPH [see *Warnings and Precautions (5.3)*].

## **9 DRUG ABUSE AND DEPENDENCE**

### **9.1 Controlled Substance**

KYZATREX contains testosterone undecanoate, a Schedule III controlled substance.

### **9.2 Abuse**

Drug abuse is intentional non-therapeutic use of a drug, even once, for its rewarding psychological and physiological effects. Abuse and misuse of testosterone are seen in male and

female adults and adolescents. Testosterone, often in combination with other anabolic androgenic steroids, and not obtained by prescription through a pharmacy, may be abused by athletes and bodybuilders. There have been reports of misuse by men taking higher doses of legally obtained testosterone than prescribed and continuing testosterone despite adverse events or against medical advice.

#### Abuse-Related Adverse Reactions

Serious adverse reactions have been reported in individuals who abuse anabolic androgenic steroids and include cardiac arrest, myocardial infarction, hypertrophic cardiomyopathy, congestive heart failure, cerebrovascular accident, hepatotoxicity, and serious psychiatric manifestations, including major depression, mania, paranoia, psychosis, delusions, hallucinations, hostility, and aggression.

The following adverse reactions have also been reported in men: transient ischemic attacks, convulsions, hypomania, irritability, dyslipidemias, testicular atrophy, subfertility, and infertility.

The following additional adverse reactions have been reported in women: hirsutism, virilization, deepening of voice, clitoral enlargement, breast atrophy, male-pattern baldness, and menstrual irregularities.

The following adverse reactions have been reported in male and female adolescents: premature closure of bony epiphyses with termination of growth, and precocious puberty.

Because these reactions are reported voluntarily from a population of uncertain size and may include abuse of other agents, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

### 9.3 Dependence

#### Behaviors Associated with Addiction

Continued abuse of testosterone and other anabolic steroids leading to addiction is characterized by the following behaviors:

- Taking greater dosages than prescribed
- Continued drug use despite medical and social problems due to drug use
- Spending significant time to obtain the drug when supplies of the drug are interrupted
- Giving a higher priority to drug use than other obligations
- Having difficulty in discontinuing the drug despite desires and attempts to do so
- Experiencing withdrawal symptoms upon abrupt discontinuation of use

Physical dependence is characterized by withdrawal symptoms after abrupt drug discontinuation or a significant dose reduction of a drug. Individuals taking suprathreshold doses of testosterone may experience withdrawal symptoms lasting for weeks or months, which may include depressed mood, major depression, fatigue, craving, restlessness, irritability, anorexia, insomnia, decreased libido, and hypogonadotropic hypogonadism.

Drug dependence in individuals using approved doses of testosterone for approved indications has not been documented.

## 10 OVERDOSAGE

There is one report of acute overdose with use of an approved injectable testosterone product: this subject had serum testosterone levels of up to 11,400 ng/dL with a cerebrovascular accident.


Treatment of overdose consists of discontinuation of KYZATREX and appropriate symptomatic and supportive care.

## 11 DESCRIPTION

KYZATREX is provided as a gelatin capsule containing testosterone undecanoate, a fatty-acid ester of testosterone. Testosterone undecanoate is a white to off-white yellow crystalline powder. Testosterone, an androgen, is formed by cleavage of the ester side chain of testosterone undecanoate.

Testosterone undecanoate is chemically described as 17 $\beta$ -hydroxyandrost-4-en-3-one undecanoate. It has the empirical formula of C<sub>30</sub>H<sub>48</sub>O<sub>3</sub> and a molecular weight of 456.7 g/mol. The structural formula for testosterone undecanoate is presented in [Figure 1](#).

**Figure 1: Testosterone Undecanoate**



KYZATREX (testosterone undecanoate) capsules for oral use are available in three dosage strengths- 100 mg, 150 mg, and 200 mg. The 100 mg strength is an opaque, white capsule imprinted with “MP100” in red ink. The 150 mg strength is an opaque white capsule imprinted with “MP150” in red ink. The 200 mg strength is an opaque white capsule imprinted with “MP200” in red ink. All capsule strengths also contain DL-alpha-tocopheryl acetate (Vitamin E), phytosterol esters, polyoxyl 40 hydrogenated castor oil, and propylene glycol monolaurate as inactive ingredients.

Gelatin capsule shells are composed of the following inactive ingredients: gelatin, glycerin, purified water, sorbitol, and titanium dioxide.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Endogenous androgens, including testosterone and dihydrotestosterone (DHT), are responsible for the normal growth and development of the male sex organs and for maintenance of secondary sex characteristics. These effects include the growth and maturation of prostate, seminal vesicles, penis, and scrotum; the development of male hair distribution, such as facial, pubic, chest, and axillary hair; laryngeal enlargement; vocal cord thickening; alterations in body musculature; and fat distribution.

Male hypogonadism, a clinical syndrome resulting from insufficient secretion of testosterone, has two main etiologies. Primary hypogonadism is caused by defects of the gonads, such as Klinefelter syndrome or Leydig cell aplasia, whereas secondary hypogonadism (also known as

hypogonadotropic hypogonadism) is the failure of the hypothalamus (or pituitary gland) to produce sufficient gonadotropins (FSH, LH).

## 12.2 Pharmacodynamics

There is insufficient data to characterize an exposure-response relationship or time course of pharmacodynamics.

## 12.3 Pharmacokinetics

### Absorption

KYZATREX was taken orally at a starting dose of 200 mg twice per day with meals in a multicenter, open-label trial in hypogonadal males. The dose was adjusted, as needed, on Days 28 and 56 from a minimum dose of 100 mg (morning-only) to a maximum dose of 400 mg twice per day based on the plasma testosterone concentration obtained by a single blood draw collected 3 to 5 hours after the morning dose. The average daily NaF/EDTA plasma testosterone concentration was 393.3 ( $\pm$ 113.6) ng/dL after 90 days of treatment (normal eugonadal range in NaF/EDTA plasma: 222-800 ng/dL. Note that the titration scheme for use in clinical practice is based on serum total testosterone [see *Dosage and Administration (2.2)*].

KYZATREX is expected to produce testosterone concentrations that approximate normal concentrations seen in healthy men.

[Table 5](#) summarizes the pharmacokinetic (PK) parameters for plasma total testosterone in patients completing at least 90 days of KYZATREX treatment administered daily.


**Table 5: NaF-EDTA Plasma Testosterone  $C_{avg}$  and  $C_{max}$  at Day 90 Visit**

PK Parameter		Plasma (N=130)
$C_{avg}$ (ng/dL)	n	127
	Mean	393.3
	SD	113.6
$C_{max}$ (ng/dL)	n	130
	Mean	852.4
	SD	311.3

PK = pharmacokinetic;  $C_{avg}$  = 24-hour average concentration;  $C_{max}$  = maximum concentration

[Figure 2](#) summarizes the mean plasma total testosterone profile at the final PK visit.

**Figure 2: Mean ( $\pm$ SEM) Concentration-Time Profile for NaF-EDTA Plasma Total Testosterone in KYZATREX-Treated Patients at Day 90 Visit**



SEM = standard error of the mean

Testosterone normal ranges: plasma = 222-800 ng/dL

When KYZATREX was given with breakfast containing 16%, 33%, and 45% fat, the exposure ( $AUC_{0-24 \text{ hr}}$ ) of testosterone was increased by 37%, 87%, and 94%, respectively, compared to when given under fasted conditions. The primary efficacy and safety study was conducted under fed conditions regardless of the type of meals and the primary efficacy endpoint of achieving testosterone  $C_{\text{avg}}$  in normal testosterone range was met.

There was no effect on testosterone PK when KYZATREX was administered with 20% alcohol along with a high-fat meal versus a high-fat meal alone.

### **Distribution**

Circulating testosterone is primarily bound in serum to sex hormone-binding globulin (SHBG) and albumin. Approximately 40% of testosterone in plasma is bound to SHBG, 2% remains unbound (free), and the rest is loosely bound to albumin and other proteins.

### **Metabolism**

The androgenic activity of testosterone undecanoate occurs after the ester bond linking the testosterone to the undecanoic acid is cleaved by endogenous non-specific esterases.

Undecanoic acid is metabolized like all fatty acids via the beta-oxidation pathway.

Testosterone is metabolized to various 17-keto steroids through two different pathways. The major active metabolites of testosterone are dihydrotestosterone (DHT) and estradiol.

### **Excretion**

About 90% of a dose of testosterone given intramuscularly is excreted in the urine as glucuronic and sulfuric acid conjugates of testosterone and its metabolites. About 6% of a dose is excreted in the feces, mostly in the unconjugated form. Inactivation of testosterone occurs primarily in the liver.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### **Carcinogenesis**

Testosterone has been tested by subcutaneous injection and implantation in mice and rats. In mice, the implant induced cervical-uterine tumors, which metastasized in some cases. There is suggestive evidence that injection of testosterone into some strains of female mice increases their susceptibility to hepatoma. Testosterone is also known to increase the number of tumors and decrease the degree of differentiation of chemically induced carcinomas of the liver in rats.

#### **Mutagenesis**

Testosterone was negative in the in vitro Ames and in the in vivo mouse micronucleus assays.

#### **Impairment of Fertility**

The administration of exogenous testosterone suppressed spermatogenesis and impaired fertility in the rat, dog, and non-human primate, which was reversible on cessation of treatment.

A reproductive toxicology study was conducted in rats to evaluate functional effects of KYZATREX on male fertility. In untreated female rats mated with males receiving 2 times the maximum recommended human daily dose (MRHDD) of KYZATREX (based on mean AUC exposure to testosterone), the number of impregnated females was reduced, fertility was significantly lower, and pre-implantation loss was significantly higher compared to the control group. There was no impairment of fertility in males receiving an equivalent dose of KYZATREX to the MRHDD.

### **13.2 Animal Toxicology and/or Pharmacology**

A 3-month repeat-dose oral toxicity study in male eugonadal dogs was conducted to evaluate whether phytosterol esters present in the KYZATREX formulation influenced target organ toxicity due to their structural similarities to sex steroids like testosterone. KYZATREX doses 2 times the MRHDD (based on mean AUC exposure to testosterone) produced similar effects on androgen-responsive tissues as testosterone undecanoate without phytosterol esters. These included mild to marked effects on the testes (decreased size, germ cell depletion, Leydig cell atrophy), epididymides (aspermia), adrenal glands (vacuolation in the zona fasciculata) and prostate (increased size and glandular hypertrophy/hyperplasia). Following a 4-week treatment-free period, findings in the testes, epididymides, and adrenal glands were not fully reversible at doses of 2 times the MRHDD of KYZATREX as compared to treatment with the excipients

alone, including phytosterol esters. Reversibility was not assessed in testosterone undecanoate groups without phytosterol esters.

## 14 CLINICAL STUDIES

The efficacy and safety of KYZATREX were evaluated in Study MRS-TU-2019EXT (NCT04467697) a multi-center, open-label study of approximately 6 months of duration in 155 hypogonadal males.

Patients received KYZATREX at a starting dose of 200 mg twice daily with meals. The dosage was adjusted on Days 28 and 56 between a minimum dose of 100 mg (single morning dose) and a maximum dose of 800 mg (400 mg twice daily) based on plasma testosterone concentration from a single blood draw between 3 to 5 hours after the morning dose.

The primary efficacy endpoint was the percentage of KYZATREX-treated patients with mean plasma total testosterone concentration ( $C_{avg}$ ) over 24-hours within the normal range of 222-800 ng/dL on the final PK visit of the study at Day 90.

The efficacy population consisted of 139 hypogonadal, males with a median age of 50 years (range 22 to 66 years), 79% were White, 16% were Black, 3% were Asian, and 2% were American Indian, Alaskan Native or Other.

Primary efficacy results are summarized in [Table 6](#).

**Table 6: Proportion of Patients in Study MRS-TU-2019EXT with Average Plasma Total Testosterone in the Normal Range (222-800 ng/dL) on Day 90**

Parameter	N=139
Patients (%) with Testosterone, $C_{avg}$ (ng/dL), 222-800 ng/dL	122 (88%)
95% Confidence Interval	(82%, 93%)

$C_{avg}$  = 24-hour average concentration

Secondary endpoints were the percentage of patients with a maximum total testosterone concentration ( $C_{max}$ ) meeting three predetermined limits: less than or equal to 1.5 times the upper limit of normal range (ULN) (1200 ng/dL), between 1.8 and 2.5 times ULN (1440-2000 ng/dL), and greater than 2.5 times ULN (2000 ng/dL).

The percentage of patients who received KYZATREX and had testosterone  $C_{max}$  threshold less than or equal to 1200 ng/dL, between 1440 and 2000 ng/dL, and greater than 2000 ng/dL at the final PK visit were 88%, 4%, and 0%, respectively.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

KYZATREX capsules are available in three strengths of 100 mg, 150 mg, and 200 mg packaged as 60, 90 and 120 units in wide-mouth, round, white HDPE bottles with white, polypropylene, child resistant caps and induction-sealed liner.

100 mg: Oval, opaque, white capsules imprinted with “MP100” in red ink; NDC 80603-101-11 for 90 capsules and NDC 80603-101-22 for 120 capsules.

150 mg: Oblong, opaque, white capsules imprinted with “MP150” in red ink; NDC 80603-103-11 for 90 capsules and NDC 80603-103-22 for 120 capsules.

200 mg: Oblong, opaque, white capsules imprinted with “MP200” in red ink; NDC-105-33 for 60 capsules, NDC 80603-105-11 for 90 capsules and NDC 80603-105-22 for 120 capsules.

Store at 20°C to 25°C (68°F to 77°F), with excursions permitted between 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Store the capsules in a dry place avoiding exposure to excessive moisture and humid conditions.

Dispose of unused KYZATREX via a take-back option. If a take-back option is unavailable, follow FDA instructions at [www.fda.gov/drugdisposal](http://www.fda.gov/drugdisposal).

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

### Polycythemia

Advise patients that KYZATREX can cause an increase in hemoglobin/hematocrit levels that may increase the risk of thromboembolic events. Advise patients about the importance of completing laboratory testing as instructed by their health care provider while on KYZATREX [see *Warnings and Precautions (5.1)*].

### Venous Thromboembolism

Advise patients that KYZATREX can cause venous thromboembolism. Advise patients of the signs and symptoms of venous thromboembolism, which may include the following: lower limb pain, edema, or erythema; and dyspnea or chest pain. Advise patients to promptly report the signs and symptoms of venous thromboembolism, discontinue use of KYZATREX and seek urgent medical care.

Worsening of Benign Prostatic Hyperplasia (BPH) Advise patients that KYZATREX can cause increased symptoms of BPH. Advise patients to contact their health care provider if they have any prostate-related symptoms [see *Warnings and Precautions (5.3)*].

### Blood Pressure Increases

Inform patients that KYZATREX can increase blood pressure (BP) which can increase cardiovascular risk over time. Instruct patients about the importance of monitoring BP periodically while on KYZATREX. If BP increases while on KYZATREX, antihypertensive medications may need to be started, added, or adjusted to control BP, or KYZATREX may need to be discontinued [see *Warnings and Precautions (5.4)*].

### Edema

Advise patients that KYZATREX can cause edema in patients with preexisting cardiac, renal, or hepatic disease. Advise patients to notify their health care provider if edema develops or worsens [*see Warnings and Precautions (5.9)*].

### Sleep Apnea

Advise patients that KYZATREX can worsen sleep apnea especially in patients with risk factors such as obesity or chronic lung diseases [*see Warnings and Precautions (5.10)*].

### Gynecomastia

Advise patients that KYZATREX can cause gynecomastia. [*see Warnings and Precautions (5.11)*].

### Administration Instructions

Advise patients to take KYZATREX with food [*see Dosage and Administration (2.3)*].

Marketed by:

Marius Pharmaceuticals  
2301 Sugar Bush Road, Suite 510  
Raleigh, NC 27612

Issued 07/2025

## MEDICATION GUIDE

### KYZATREX® (ky-ZAH-treks) (testosterone undecanoate) capsules, for oral use, CIII

#### What is KYZATREX®?

- KYZATREX® is a prescription medicine that contains testosterone. KYZATREX® is used to treat adult men who have low or no testosterone due to certain medical conditions.
- It is not known if KYZATREX® is safe or effective in children younger than 18 years old. Improper use of KYZATREX® may affect bone growth in children.
- KYZATREX® is a controlled substance (CIII) because it contains testosterone that can be a target for people who abuse prescription medicines. Keep your KYZATREX® in a safe place to protect it. Never give your KYZATREX® to anyone else, even if they have the same symptoms you have. Selling or giving away this medicine may harm others and is against the law.
- KYZATREX® is not meant for use by women.

#### Do not take KYZATREX® if you:

- have breast cancer.
- have or might have prostate cancer.
- are a woman who is pregnant. KYZATREX may harm your unborn baby.
- are allergic to KYZATREX® or any ingredients in KYZATREX®. See the end of this Medication Guide for a complete list of ingredients in KYZATREX®.

#### Before you take KYZATREX®, tell your healthcare provider about all of your medical conditions, including if you:

- have high blood pressure or are being treated for high blood pressure.
- have a history of diabetes.
- have heart problems.
- have high red blood cell count (hematocrit) or high hemoglobin laboratory value.
- have urinary problems due to an enlarged prostate.
- have liver or kidney problems.
- have problems breathing while you sleep (sleep apnea).

**Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.** Taking KYZATREX® with certain other medicines can affect each other.

#### Especially, tell your healthcare provider if you take:

- insulin
- medicines that decrease blood clotting (blood thinners)
- corticosteroids
- medicines that increase blood pressure such as some cold medicine and pain medicines

Know the medicines you take. Ask your healthcare provider or pharmacist for a list of these medicines, if you are not sure. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

### How should I take KYZATREX®?

- Take KYZATREX® exactly as your healthcare provider tells you take it.
- Take KYZATREX® by mouth 2 times daily. Take 1 time in the morning and take 1 time in the evening. If your dose is 100 mg daily, take 1 time in the morning.
- Take KYZATREX® with food.
- Your healthcare provider may change your KYZATREX® dose. Do not change your KYZATREX® dose without talking to your healthcare provider.

### What are the possible side effects of KYZATREX®?

#### KYZATREX® may cause serious side effects including:

- See “**What is the most important information I should know about KYZATREX®?**”
- **Increase in red blood cell count (hematocrit) or hemoglobin.**
  - KYZATREX® increases red blood cell counts in some patients. High red blood cell counts increase the risk of blood clots, strokes, and heart attacks.
  - You may need to stop KYZATREX if your red blood cell count increases.
  - Your healthcare provider should check your red blood cell count and hemoglobin while you take KYZATREX®.
- **If you already have an enlarged prostate, your signs and symptoms may worsen while taking KYZATREX®.** These may include:
  - increased urination at night
  - trouble starting your urine stream
  - urinating many times during the day
  - urge to go to the bathroom right away
  - a urine accident
  - inability to pass urine or weak urine flow
- **Increased risk of prostate cancer.** Your healthcare provider should check you for prostate cancer or any other prostate problems before you start and while you take KYZATREX®
- **Blood clots in the legs or lungs.** Signs and symptoms of a blood clot in your leg can include pain, swelling or redness. Signs and symptoms of a blood clot in your lungs can include difficulty breathing or chest pain.
- **Increase in blood pressure.** KYZATREX can increase your blood pressure. Increases in blood pressure can increase the risk of heart attack or stroke over time. If your blood pressure increases while on KYZATREX, blood pressure medicines may need to be started. If you are taking blood pressure medicines, new blood pressure medicines may need to be added or your current blood pressure medicines may need to be adjusted to control your blood pressure. If your blood pressure cannot be controlled, KYZATREX may need to be stopped. Your healthcare provider will monitor your blood pressure while you are being treated with KYZATREX.
- **Abuse.** Testosterone can be abused, when taken at higher than prescribed doses and when used with other anabolic androgenic steroids. Abuse can cause serious heart and psychological side effects. Your healthcare provider should check you for signs of abuse before and during treatment with KYZATREX®.
- **In large doses KYZATREX® may lower your sperm count.**
- **Liver problems.** Symptoms of liver problems may include:
  - nausea or vomiting
  - yellowing of your skin or whites of your eyes
  - dark urine
  - pain on the right side of your stomach area (abdominal pain)
- **Swelling of your ankles, feet, or body (edema), with or without heart failure.**
- **Enlarged or painful breasts.**
- **Breathing problems while you sleep (sleep apnea).**

**Call your healthcare provider right away if you have any of the serious side effects listed above.**

**The most common side effect of KYZATREX® is high blood pressure.**

**Other side effects may include** headache, joint or back pain, diarrhea, increased red blood cell count, anxiety, constipation, swelling of the legs, and increased prostate specific antigen (PSA) levels.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of KYZATREX®. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store KYZATREX®?**

- Store KYZATREX® at room temperature between 68°F to 77°F (20°C to 25°C).
- Store KYZATREX® in a dry place.

**Keep KYZATREX and all medicines out of the reach of children.**

**How should I throw away (dispose of) KYZATREX®?**

- Throw away unused KYZATREX® via a take-back option.
- If a take-back option is unavailable, follow FDA instructions at [www.fda.gov/drugdisposal](http://www.fda.gov/drugdisposal) for properly throwing away medicine.

**General information about the safe and effective use of KYZATREX®**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use KYZATREX® for a condition for which it was not prescribed. Do not give KYZATREX® to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about KYZATREX® that is written for health professionals.

**What are the ingredients in KYZATREX®?**

**Active ingredient:** testosterone undecanoate

**Inactive ingredients:** DL-alpha-tocopheryl acetate (Vitamin E), phytosterol esters, polyoxyl 40 hydrogenated castor oil and propylene glycol monolaurate. The ingredients of the gelatin capsule shells are gelatin, glycerin, purified water, sorbitol, and titanium dioxide.

Marketed by:

Marius Pharmaceuticals  
Raleigh, NC 27612

For more information, go to [www.KYZATREX.com](http://www.KYZATREX.com) or call 1-83-949-5040

This Medication Guide has been approved by the U.S. Food and Drug Administration

Issued: 07/2025