

A | B | C | D | E | F | G | H

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- If you smoke, quit smoking.

- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.

000000000

May take 1 to 4 days for full effect

First 2 doses of a 14-day course of treatment

2 TABLETS

Melts in your mouth
Dissolves without water

Acid Reducer

Delayed Release

Orally Disintegrating Tablets 20 mg

Esomeprazole

Treats Frequent Heartburn!
24 HR

Compare to the active ingredient in Nexium® 24HR*

NDC 64230-112-05

MELTech™
Melts In Your Mouth

FREE SAMPLE

No Lacquer →

← No Lacquer

STANDART FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Bold Cond. Oblique
TITLE (continued): 7.5 pt Helvetica Bold Cond. Oblique
(continued) 7.5 pt Helvetica Medium Cond.
HEADINGS: 8 pt Helvetica Bold Cond. Oblique
SUBHEAD NGS: 6 pt Helvetica Bold Cond. Oblique
TEXT: 6 pt Helvetica Medium Cond.
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6.5 pt Helvetica Bold Cond.
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

D MENSIONS: 117 x 17 x 90 mm

Do not use if printed tablet blister unit is open or torn.

Safety Feature -

NOT FOR SALE

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

*The product is not manufactured or distributed by CS, distributor of Nexium® 24HR.

D-55

Batch No

EXP

Drug Facts	Drug Facts (continued)
<p>Active ingredient (in each tablet) Esomeprazole 20 mg (Each delayed-release orally disintegrating tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Purpose Acid reducer</p> <p>Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are: ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p>	<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect</p> <p>14-Day Course of Treatment ■ Take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ store in the original package</p> <p>Inactive ingredients acetone, alcohol, amino methacrylate copolymer, ascorbic acid, benzyl alcohol, cetyl alcohol, colloidal silicon dioxide, croscopolidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, flavor, hypromellose, hypromellose phthalate, isopropyl alcohol, mannitol, microcrystalline cellulose, modified starch, polysorbate 80, pregelatinized starch, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, taic, titanium dioxide, triacetin, triethyl citrate</p>

Drug Facts (continued)
[Questions or comments? 1-800-XXX-XXXX | weekdays XXX AM to XXX PM EST]

Made in Israel
Manufactured by:
Dexal Pharma Technologies Ltd.,
10 Hakdima St., Yokneam 2069200, Israel

A B C D E F G H

No Lacquer →

← No Lacquer

2 TABLETS

First 2 doses of a 14-day course of treatment
May take 1 to 4 days for full effect

Wildberry Flavor

Actual size

Melts in your mouth
Dissolves without water

Acid Reducer

Delayed Release

Esomeprazole

Orally Disintegrating Tablets 20 mg

Treats Frequent Heartburn!
24 HR

Compare to the active ingredient in Nexium® 24HR*

FREE SAMPLE

Melts In Your Mouth
MELTECH™

NDC 64239-1-12-10

000000000

- Tips for Managing Heartburn**
- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
 - Eat slowly and do not eat big meals.
 - Do not eat late at night or just before bedtime.
 - If you smoke, quit smoking.
 - Do not lie flat or bend over soon after eating.
 - Raise the head of your bed.
 - Wear loose-fitting clothing around your stomach.
 - If you are overweight, lose weight.

NOT FOR SALE

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

*This product is not manufactured or distributed by GSK, distributor of Nexium® 24HR

Safety Feature -
Do not use if printed tablet blister unit is open or torn.

Made in Israel
Manufactured by:
Develcal Pharma Technologies Ltd.
10 Hakdima St., Yokneam 2069200, Israel

Batch No
EXP

Drug Facts		Drug Facts (continued)
Active ingredient (in each tablet) Esomeprazole 20 mg (Each delayed-release orally disintegrating tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)	Purpose Acid reducer	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect		Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect 14-Day Course of Treatment ■ Take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.		Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ store in the original package
Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain		Inactive ingredients acetone, alcohol, amino methacrylate copolymer, ascorbic acid, benzyl alcohol, cetyl alcohol, colloidal silicon dioxide, croscopolidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, flavor, hypromellose, hypromellose phthalate, isopropyl alcohol, mannitol, microcrystalline cellulose, modified starch, polysorbate 80, pregelatinized starch, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, taic, titanium dioxide, triacetin, triethyl citrate
Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.		
Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain		

Drug Facts (Continued)
Questions or comments? 1-800-XXX-XXXX | weekdays XXX AM to XXX PM EST

STANDART FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Bold Cond. Oblique
TITLE (continued): 7.5 pt Helvetica Bold Cond. Oblique
(continued) 7.5 pt Helvetica Medium Cond.
HEADINGS: 8 pt Helvetica Bold Cond. Oblique
SUBHEAD NGS: 6 pt Helvetica Bold Cond. Oblique
TEXT: 6 pt Helvetica Medium Cond.
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6.5 pt Helvetica Bold Cond.
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

D MENSIONS: 117 x 17 x 90 mm

A B C D E F G H

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- If you smoke, quit smoking.

- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.

000000000

Actual size



Wildberry Flavor

May take 1 to 4 days for full effect

14 TABLETS

Melts in your mouth
Dissolves without water

Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

Treats Frequent Heartburn!
24 HR

Compare to the active ingredient in Nexium® 24HR*

Melts In Your Mouth
MELTech™

NDC 64229-112-01

No Lacquer →

← No Lacquer

STANDART FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Bold Cond. Oblique
TITLE (continued): 7.5 pt Helvetica Bold Cond. Oblique
(continued) 7.5 pt Helvetica Medium Cond.
HEADINGS: 8 pt Helvetica Bold Cond. Oblique
SUBHEAD NGS: 6 pt Helvetica Bold Cond. Oblique
TEXT: 6 pt Helvetica Medium Cond.
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6.5 pt Helvetica Bold Cond.
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

D MENSIONS: 117 x 17 x 90 mm

Do not use if printed tablet blister unit is open or torn.

Safety Feature -

NOT FOR RESALE

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

*This product is not manufactured or distributed by GSK, distributor of Nexium® 24HR.

D-55

Batch No

EXP

Drug Facts	Drug Facts (continued)
<p>Active ingredient (in each tablet) Esomeprazole 20 mg - Acid reducer (Each delayed-release orally disintegrating tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p>	<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect 14-Day Course of Treatment ■ Take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ store in the original package</p> <p>Inactive ingredients acetone, alcohol, amino methacrylate copolymer, ascorbic acid, benzyl alcohol, cetyl alcohol, colloidal silicon dioxide, croscopolidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, flavor, hypromellose, hypromellose phthalate, isopropyl alcohol, mannitol, microcrystalline cellulose, modified starch, polysorbate 80, pregelatinized starch, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, taic, titanium dioxide, triacetin, triethyl citrate</p>

Drug Facts (Continued)

Questions or comments? 1-800-XXX-XXXX | weekdays XXX AM to XXX PM EST

Made in Israel
 Manufactured by:
 Dexal Pharma Technologies Ltd.,
 10 Hakdima St., Yokneam 2069200, Israel

A B C D E F G H

0000000000

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- If you smoke, quit smoking.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.

Actual size

Wildberry Flavor

One 14-day course of treatment
May take 1 to 4 days for full effect

14 TABLETS

Melts in your mouth
Dissolves without water

Acid Reducer

Delayed Release
Orally Disintegrating Tablets 20 mg

Treats Frequent Heartburn
24 HR

Melts In Your Mouth
MELTECH™

Compare to the active ingredient
in Nexium® 24HR

NDC 64239-112-06

No Lacquer →

← No Lacquer

STANDART FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Bold Cond. Oblique
TITLE (continued): 7.5 pt Helvetica Bold Cond. Oblique
(continued) 7.5 pt Helvetica Medium Cond.
HEADINGS: 8 pt Helvetica Bold Cond. Oblique
SUBHEAD NGS: 6 pt Helvetica Bold Cond. Oblique
TEXT: 6 pt Helvetica Medium Cond.
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6.5 pt Helvetica Bold Cond.
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

D MENSIONS: 117 x 17 x 90 mm

Do not use if printed tablet blister unit is open or torn.

Safety Feature -

IMPORTANT INFORMATION
The product is not manufactured or distributed by CSK, distributor of Nexium® 24HR.

NOT FOR RESALE

KEEP THE CARTON. IT CONTAINS

D-55

<p>Made in Israel Manufactured by: Dexal Pharma Technologies Ltd. 10 Hakdima St., Yokneam 2069200, Israel</p>	<p>Drug Facts</p> <p>Active ingredient (in each tablet) Esomeprazole 20 mg</p> <p>Purpose Acid reducer (Each delayed-release orally disintegrating tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p>	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>
	<p>Uses</p> <ul style="list-style-type: none"> ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect <p>Warnings Allergy alert:</p> <ul style="list-style-type: none"> ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash ■ If an allergic reaction occurs, stop use and seek medical help right away. <p>Do not use if you have:</p> <ul style="list-style-type: none"> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain <p>These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs. <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain 	<p>Directions</p> <ul style="list-style-type: none"> ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ Take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ store in the original package <p>Inactive ingredients</p> <p>acetone, alcohol, amino methacrylate copolymer, ascorbic acid, benzyl alcohol, cetyl alcohol, colloidal silicon dioxide, croscopolidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, flavor, hypromellose, hypromellose phthalate, isopropyl alcohol, mannitol, microcrystalline cellulose, modified starch, polysorbate 80, pregelatinized starch, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, taic, titanium dioxide, triacetin, triethyl citrate</p>

Drug Facts (Continued)
Questions or comments? 1-800-XXX-XXXX | weekdays XXX AM to XXX PM EST

Batch No
EXP

A B C D E F G H

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- If you smoke, quit smoking.

- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.

0000000000

May take 1 to 4 days for full effect

14 TABLETS

One 14-day course of treatment

Melts in your mouth

Dissolves without water

Acid Reducer

Delayed Release

Esomeprazole

Orally Disintegrating Tablets 20 mg

Treats Frequent Heartburn!

24 HR

Compare to the active ingredient in Nexium® 24HR*

Melts In Your Mouth

MELTech™

ND C 64239-11-2-02

No Lacquer →

← No Lacquer

STANDART FORMAT

TYPE SIZES:
 TITLE: 9 pt Helvetica Bold Cond. Oblique
 TITLE (continued): 7.5 pt Helvetica Bold Cond. Oblique
 (continued) 7.5 pt Helvetica Medium Cond.
 HEADINGS: 8 pt Helvetica Bold Cond. Oblique
 SUBHEAD NGS: 6 pt Helvetica Bold Cond. Oblique
 TEXT: 6 pt Helvetica Medium Cond.
 LEADING: 6.5 pt
 BULLETS: 5 pt Square with 2 M'S spacing between statements
 TELEPHONE #: 6.5 pt Helvetica Bold Cond.
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt

D MENSIONS: 117 x 17 x 90 mm

Safety Feature -

Do not use if printed blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

*The product is not manufactured or distributed by GSK, distributor of Nexium® 24HR

D-55

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Esomeprazole 20 mg</p> <p>Purpose Acid reducer (Each delayed-release orally disintegrating tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are: ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p>	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect</p> <p>14-Day Course of Treatment ■ Take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ store in the original package</p> <p>Inactive ingredients acetone, alcohol, amino methacrylate copolymer, ascorbic acid, benzyl alcohol, cetyl alcohol, colloidal silicon dioxide, croscopolidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, flavor, hypromellose, hypromellose phthalate, isopropyl alcohol, mannitol, microcrystalline cellulose, modified starch, polysorbate 80, pregelatinized starch, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, taic, titanium dioxide, triacetin, triethyl citrate</p>
--	--

Made in Israel
 Manufactured by:
 Dexal Pharma Technologies Ltd.,
 10 Hakdima St., Yokneam 2069200, Israel

Batch No
 EXP

Drug Facts (Continued)
 (Questions or comments? 1-800-XXX-XXXX | weekdays XXX AM to XXX PM EST)

A | B | C | D | E | F | G | H


Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- If you smoke, quit smoking.

- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.

000000000

Actual size



Wildberry Flavor

One 14-day course of treatment
May take 1 to 4 days for full effect

14 TABLETS

Disolves without water

Melts in your mouth

Acid Reducer

Delayed Release
Orally Disintegrating Tablets 20 mg

Esomeprazole

Treats Frequent Heartburn!

24 HR

Melts In Your Mouth

MELTECH™

Compare to the active ingredient
in Nexium® 24HR*

NDC 64239-112-07

No Lacquer →

← No Lacquer

STANDART FORMAT


TYPE SIZES:
TITLE: 9 pt Helvetica Bold Cond. Oblique
TITLE (continued): 7.5 pt Helvetica Bold Cond. Oblique
(continued) 7.5 pt Helvetica Medium Cond.
HEADINGS: 8 pt Helvetica Bold Cond. Oblique
SUBHEAD NGS: 6 pt Helvetica Bold Cond. Oblique
TEXT: 6 pt Helvetica Medium Cond.
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6.5 pt Helvetica Bold Cond.
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

D MENSIONS: 117 x 17 x 90 mm

Safety Feature -
Do not use if printed tablet
blister unit is open or torn.

KEEP THE CARTON. IT CONTAINS
IMPORTANT INFORMATION.

*This product is not manufactured
or distributed by GSK, distributor of
Nexium® 24HR.



D-55

Batch No

EXP

Drug Facts	Drug Facts (continued)
<p>Active ingredient (in each tablet) Esomeprazole 20 mg (Each delayed-release orally disintegrating tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Purpose Acid reducer</p> <p>Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition.</p> <p>■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain</p>	<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect</p> <p>14-Day Course of Treatment ■ Take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water. ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol</p> <p>Repeated 14-Day Course (if needed) ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor</p> <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ store in the original package</p> <p>Inactive ingredients acetone, alcohol, amino methacrylate copolymer, ascorbic acid, benzyl alcohol, cetyl alcohol, colloidal silicon dioxide, croscopolidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, flavor, hypromellose, hypromellose phthalate, isopropyl alcohol, mannitol, microcrystalline cellulose, modified starch, polysorbate 80, pregelatinized starch, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, taic, titanium dioxide, triacetin, triethyl citrate</p>

Manufactured by:
Dexal Pharma Technologies Ltd.,
10 Hakdima St., Yokneam 2069200, Israel

Made in Israel

Drug Facts (Continued)
Questions or comments? 1-800-XXX-XXXX | weekdays XXX AM to XXX PM EST

A B C D E F G H

Tips for Managing Heartburn


- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Drug Facts (continued)

Inactive ingredients acetone, alcohol, amino methacrylate copolymer, ascorbic acid, benzyl alcohol, cetyl alcohol, colloidal silicon dioxide, crospovidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, flavor, hypromellose, hypromellose phthalate, isopropyl alcohol, mannitol, microcrystalline cellulose, modified starch, polysorbate 80, pregelatinized starch, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-XXX-XXXX weekdays XXX AM to XXX PM EST

Actual size



Wildberry Flavor

May take 1 to 4 days for full effect

28 TABLETS

Melts in your mouth
Dissolves without water

Acid Reducer

Delayed Release
Orally Disintegrating Tablets 20 mg

Esomeprazole

Treats Frequent Heartburn!
24 HR

Compare to the active ingredient in Nexium® 24HR

MELTech™
Melts In Your Mouth

Batch No
EXP


MDC 6429-1-12-08

**This product is not manufactured or distributed by GSK, distributor of Nexium® 24HR*

*Made in Israel
Manufactured by
Dexal Pharma Technologies Ltd,
10 Hakikima St. Yokneam 2069200, Israel*

Safe by Feature - Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON.
IT CONTAINS IMPORTANT INFORMATION.



Esomeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg
Acid Reducer

D-28

Drug Facts

Active Ingredient (in each tablet)	Purpose
Esomeprazole 20 mg (Each delayed-release orally disintegrating tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)	Acid reducer

Uses

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.

Warnings

Allergy alert:

- Do not use if you are allergic to esomeprazole.
- Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens

Drug Facts (continued)

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- take 1 tablet before eating in the morning
- do not crush or chew tablets
- place the tablet on tongue; tablet disintegrates, with or without water.

The tablets can also be swallowed whole with water.

- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

Repeated 14-Day Course (if needed).


- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F); keep product out of high heat and moisture
- store in the original package

No Lacquer →



← No Lacquer

STANDART FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Bold Cond. Oblique
TITLE (continued): 7.5 pt Helvetica Bold Cond. Oblique
(continued) 7.5 pt Helvetica Medium Cond.
HEADINGS: 8 pt Helvetica Bold Cond. Oblique
SUBHEADINGS: 6 pt Helvetica Bold Cond. Oblique
TEXT: 6 pt Helvetica Medium Cond.
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6.5 pt Helvetica Bold Cond.
HEAVY L NES: 1.5 pt
HAIRLINES: 0.5 pt

DIMENSIONS: 120 x 42 x 94 mm

A B C D E F G H

Tips for Managing Heartburn


- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Drug Facts (continued)

Inactive ingredients acetone, alcohol, amino methacrylate copolymer, ascorbic acid, benzyl alcohol, cetyl alcohol, colloidal silicon dioxide, crospovidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, flavor, hypromellose, hypromellose phthalate, isopropyl alcohol, mannitol, microcrystalline cellulose, modified starch, polysorbate 80, pregelatinized starch, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-XXX-XXXX weekdays XXX AM to XXX PM EST

Actual size



Wildberry Flavor

28 TABLETS

Two 14-day courses of treatment
May take 1 to 4 days for full effect

Melts in your mouth
Dissolves without water

Acid Reducer

Delayed Release

Esomeprazole

Orally Disintegrating Tablets 20 mg

Treats Frequent Heartburn! 24 HR

Compare to the active ingredient in Nexium® 24HR

MELTech™ Melts In Your Mouth

Batch No
EXP


MDC 6429-1-12-08

Made in Israel
Manufactured by
Dexal Pharma Technologies Ltd.
10 Hakdama St. Yokneam 206200, Israel

*This product is not manufactured or distributed by GSK, distributor of Nexium® 24HR

Safe by Feature - Do not use if printed tablet blister unit is open or torn.

KEEP THE CARTON.
IT CONTAINS IMPORTANT INFORMATION.



Acid Reducer

Delayed Release

Orally Disintegrating Tablets 20 mg

Esomeprazole

D-28

Drug Facts

Active ingredient (in each tablet)	Purpose
Esomeprazole 20 mg	Acid reducer
(Each delayed-release orally disintegrating tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)	

Uses

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.

Warnings

Allergy alert:

- Do not use if you are allergic to esomeprazole.
- Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
- If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens

Drug Facts (continued)

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- take 1 tablet before eating in the morning
- do not crush or chew tablets
- place the tablet on tongue; tablet disintegrates, with or without water.

The tablets can also be swallowed whole with water.

- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F); keep product out of high heat and moisture
- store in the original package

STANDART FORMAT

TYPE SIZES:
TITLE: 9 pt Helvetica Bold Cond. Oblique
TITLE (continued): 7.5 pt Helvetica Bold Cond. Oblique
(continued) 7.5 pt Helvetica Medium Cond.
HEADINGS: 8 pt Helvetica Bold Cond. Oblique
SUBHEADINGS: 6 pt Helvetica Bold Cond. Oblique
TEXT: 6 pt Helvetica Medium Cond.
LEADING: 6.5 pt
BULLETS: 5 pt Square with 2 M'S spacing between statements
TELEPHONE #: 6.5 pt Helvetica Bold Cond.
HEAVY LINES: 1.5 pt
HAIRLINES: 0.5 pt

DIMENSIONS: 120 x 42 x 94 mm

No Lacquer →

← No Lacquer

A B C D E F G H

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Drug Facts (continued)

Inactive ingredients acetone, alcohol, amino methacrylate copolymer, ascorbic acid, benzyl alcohol, cetyl alcohol, colloidal silicon dioxide, croscopvidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, flavor, hypromellose, hypromellose phthalate, isopropyl alcohol, mannitol, microcrystalline cellulose, modified starch, polysorbate 80, pregelatinized starch, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-XXX-XXXX weekdays XXX AM to XXX PM EST

Batch No
EXP

Made in Israel
Manufactured by
Deemed Pharma Technologies Ltd.
10 Hakidana St. Yokneam 2089200, Israel

Esomeprazole
Delayed Release
Orally Disintegrating Tablets 20 mg

Wildberry Flavor

Actual size



3 1/2

24 HR

Melts in your mouth
Dissolves without water

Three 14-day courses of treatment
May take 1 to 4 days for full effect

42 TABLETS

Treats Frequent Heartburn!

Compare to the active ingredient
in Nexium® 24HR*

MELTech™
Melts In Your Mouth

NDC 6429-112-04

Acid Reducer

Delayed Release
Orally Disintegrating Tablets 20 mg

Esomeprazole

Acid Reducer

D-42



KEEP THE CARTON
IT CONTAINS IMPORTANT INFORMATION.

Safety Feature - Do not use if printed
tablet blister units open or torn.

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Esomeprazole 20 mg</p> <p>Purpose Acid reducer (Each delayed-release orally disintegrating tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p>Uses ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</p> <p>Warnings Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.</p> <p>Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain</p> <p>Ask a doctor or pharmacist before use if you are: ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if: ■ your heartburn continues or worsens</p>	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p> <p>Directions</p> <ul style="list-style-type: none"> ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect <p>14-Day Course of Treatment</p> <ul style="list-style-type: none"> ■ take 1 tablet before eating in the morning ■ do not crush or chew tablets ■ place the tablet on tongue; tablet disintegrates, with or without water. <p>The tablets can also be swallowed whole with water.</p> <ul style="list-style-type: none"> ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ do not take this medicine with alcohol <p>Repeated 14-Day Course (if needed)</p> <ul style="list-style-type: none"> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor <p>■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p>Other information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20-25°C (68-77°F); keep product out of high heat and moisture ■ store in the original package
---	--

STANDART FORMAT

TYPE SIZES:
 TITLE: 9 pt Helvetica Bold Cond. Oblique
 TITLE (continued): 7.5 pt Helvetica Bold Cond. Oblique (continued)
 7.5 pt Helvetica Medium Cond.
 HEADINGS: 8 pt Helvetica Bold Cond. Oblique
 SUBHEADINGS: 6 pt Helvetica Bold Cond. Oblique
 TEXT: 6 pt Helvetica Medium Cond.
 LEADING: 6.5 pt
 BULLETS: 5 pt Square with 2 M'S spacing between statements
 TELEPHONE #: 6.5 pt Helvetica Bold Cond.
 HEAVY LINES: 1.5 pt
 HAIRLINES: 0.5 pt

DIMENSIONS: 120 x 58 x 94 mm

A B C D E F G H

Batch No

EXP

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.


Drug Facts (continued)

Inactive ingredients acetone, alcohol, amino methacrylate copolymer, ascorbic acid, benzyl alcohol, cetyl alcohol, colloidal silicon dioxide, crospovidone, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, flavor, hypromellose, hypromellose phthalate, isopropyl alcohol, mannitol, microcrystalline cellulose, modified starch, polysorbate 80, pregelatinized starch, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-XXX-XXXX weekdays XXX AM to XXX PM EST

This product is not manufactured or distributed by GSK, distributor of Nexium® 24HR

Actual size



42 TABLETS

Three 14-day courses of treatment

May take 1 to 4 days for full effect

Melts in your mouth

Dissolves without water

Acid Reducer

Delayed Release

Orally Disintegrating Tablets 20 mg

Esomeprazole

Treats Frequent Heartburn!

24 HR

Compare to the active ingredient in Nexium® 24HR*

Melts in Your Mouth

MELTECH™

NDC 64239-112-09

This product is not manufactured or distributed by GSK, distributor of Nexium® 24HR

Made in Israel
Manufactured by:
Chesed Pharmaceuticals Ltd,
10 Haldimida St., Yokneam 268200, Israel

D-42

KEEP THE CARTON.
IT CONTAINS IMPORTANT INFORMATION.

Safety Feature - Do not use if printed tablet blister unit is open or torn.

Drug Facts

Active ingredient (in each tablet)	Purpose
Esomeprazole 20 mg. (Each delayed-release orally disintegrating tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)	Acid reducer

Uses

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.

Warnings

Allergy alert:

- Do not use if you are allergic to esomeprazole.
- Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens

Drug Facts (continued)

- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- take 1 tablet before eating in the morning
- do not crush or chew tablets
- place the tablet on tongue; tablet disintegrates, with or without water.

The tablets can also be swallowed whole with water.

- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F); keep product out of high heat and moisture
- store in the original package

D-42

STANDART FORMAT

TYPE SIZES:

TITLE: 9 pt Helvetica Bold Cond. Oblique

TITLE (continued): 7.5 pt Helvetica Bold Cond. Oblique (continued) 7.5 pt Helvetica Medium Cond.

HEADINGS: 8 pt Helvetica Bold Cond. Oblique

SUBHEADINGS: 6 pt Helvetica Bold Cond. Oblique

TEXT: 6 pt Helvetica Medium Cond.

LEADING: 6.5 pt

BULLETS: 5 pt Square with 2 M'S spacing between statements

TELEPHONE #: 6.5 pt Helvetica Bold Cond.

HEAVY LINES: 1.5 pt

HAIRLINES: 0.5 pt

DIMENSIONS: 120 x 58 x 94 mm

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
05/11/2022 02:08:36 PM